

### 1.1 Introduction

We introduce this chapter with a brief discussion of the integers and close the same with a brief discussion of the field properties and absolute values of the real numbers. Integers as you know, look very simple and very unlikely to hide any interesting properties. However, this impression is far from reality. Based on the properties of integers, so far explored by Mathematicians both ancient and modern, a fascinating branch of Mathematics under the name “Number Theory” comes into existence. Almost every great mathematician has made contribution to the development of this branch which goes on expanding. Here in this chapter, two interesting topics from the elementary number theory and some fundamental properties of the real numbers will be discussed.

### 1.2 Euclid’s Division Lemma

About arithmetic, Carl Friedrich Gauss (1777-1855) a German mathematician, considered to be one of the greatest mathematicians in history once said, “Mathematics is the Queen of all Sciences and Arithmetic is the Queen of Mathematics”. And integers are introduced very early in the elementary arithmetic and thereupon you have studied many interesting properties of integers. You have learnt that the sum, difference and product of two integers are all integers. In the language of Modern Algebra these properties are expressed by saying that the set of integers  $Z$  is closed under addition, subtraction and multiplication. Can you now tell whether  $Z$  is closed under division or not? Certainly  $Z$  is not closed under division, for the result of division of one integer by another is not always an integer. For example, 36 divided by 7 yields  $\frac{36}{7}$  which is not an integer. Further, division of an integer by the integer zero is undefined. This may appear to be an obstacle in the discussion of properties of integers, but it is not so. Had  $Z$  been closed under division, Mathematics would have been deprived of one of its fascinating branches namely Number Theory.

Let us consider the division of 36 by 7. As usual it is carried out as follows:

$$\begin{array}{r} 7)36(5 \\ \underline{35} \\ 1 \end{array}$$

For such a long division sum you have seen that

$$\text{Dividend} = \text{Divisor} \times \text{Quotient} + \text{Remainder}$$

We now state Euclid’s algorithm for finding HCF of two given positive integers, stepwise as follows:

- Step 1.** Find the quotient and remainder of the division of the greater number by the smaller.
- Step 2.** If the remainder is zero, then the divisor is the HCF.
- Step 3.** Else, taking the previous remainder as the new divisor and the previous divisor as the new dividend, find the quotient and remainder.
- Step 4.** Continue the process till the remainder is zero. The last divisor is the required HCF.

Let  $a, b$  be two given positive integers where  $a > b$ . Then applying the successive steps of the above algorithm, we obtain a series of relations as follows :

$$a = bq + r_1 \quad , \quad 0 < r_1 < b \quad (1)$$

$$b = r_1q_1 + r_2 \quad , \quad 0 < r_2 < r_1 \quad (2)$$

$$r_1 = r_2q_2 + r_3 \quad , \quad 0 < r_3 < r_2 \quad (3)$$

... ..

$$r_{n-2} = r_{n-1}q_{n-1} + r_n \quad , \quad 0 < r_n < r_{n-1} \quad (n)$$

$$r_{n-1} = r_nq_n + 0 \quad (n+1)$$

Here,  $a > b > r_1 > r_2 > \dots > r_n$  and as shown in the  $(n+1)$ th relation, the last remainder is zero. The divisor at this stage i.e. the last divisor  $r_n$  is the HCF of  $a$  and  $b$ .

The set of equations (1) to  $(n + 1)$  is usually referred to as “Euclid’s algorithm for  $a$  and  $b$ ”. To find the HCF of two given positive integers, we first form Euclid’s algorithm for the integers and then conclude that the last divisor is the HCF. We shall use the symbol  $(x,y)$  to denote the HCF of the two integers  $x$  and  $y$ .

An important result from number theory, is given here in the form of the following theorem, the proof of which goes beyond the scope of this book.

**Theorem:** If  $a = bq + r$   
then  $(a,b) = (b, r)$ .

Applying this result successively to the relations (1) to  $(n+1)$ , it may be verified that

$$(a,b) = (b, r_1) = (r_1, r_2) = \dots = (r_{n-1}, r_n) = r_n.$$

**Example 1.** Using Euclid’s algorithm find the HCF of 378 and 735.

**Solution :** Euclid’s algorithm for the two integers comprises of the following three equalities:

**Example 5. Show that every odd integer is of the form  $4k + 1$  or  $4k - 1$ .**

**Solution :** Let  $a$  be any odd integer

Then ,  $a = 2q + 1$  for some integer  $q$ , even or odd.

If  $q$  is even, then  $q = 2k$  for some integer  $k$  and  $a = 2(2k) + 1 = 4k + 1$ .

If  $q$  is odd, then  $q = 2k - 1$  for some integer  $k$  and  $a = 2(2k - 1) + 1 = 4k - 1$ .

Thus, every odd integer  $a$  is of the form  $4k + 1$  or  $4k - 1$ .

**Example 6. Show that every square integer is of the form  $4k$  or  $4k + 1$ .**

**Solution :** Let  $a^2$  be a given square integer. The integer  $a$  is of the form  $2q$  or  $2q + 1$ . Also

$$(2q)^2 = 4q^2 = 4k, \text{ where } k = q^2 \text{ is an integer,}$$

$$\text{and } (2q+1)^2 = 4q^2 + 4q + 1$$

$$= 4k + 1, \text{ where } k = q^2 + q \text{ is an integer.}$$

$\therefore a^2$  is of the form  $4k$  or  $4k + 1$ .

**Example 7. Prove that one of every three consecutive integers is divisible by 3.**

**Solution :** Let  $a, a+1, a+2$  be three consecutive integers. The integer  $a$  is of the form  $3q, 3q+1$  or  $3q-1$ .

If  $a = 3q$ , the result holds trivially

If  $a = 3q+1$ , then  $a+2 = 3q+3 = 3(q+1)$  which is divisible by 3.

If  $a = 3q-1$ , then  $a+1 = 3q$  is divisible by 3

Hence, one of  $a, a+1$  and  $a+2$  is divisible by 3. This proves the result.

**Example 8. If  $a$  is an odd integer, show that  $a^2 + (a+2)^2 + (a+4)^2 + 1$  is divisible by 12.**

**Solution :** The odd integer  $a$  is of the form  $4k+1$  or  $4k-1$  for some integer  $k$ .

If  $a = 4k+1$ , then

$$a^2 + (a+2)^2 + (a+4)^2 + 1 = (4k+1)^2 + (4k+3)^2 + (4k+5)^2 + 1$$

$$= 48k^2 + 72k + 36$$

$$= 12(4k^2 + 6k + 3) \quad \text{which is divisible by 12.}$$

factors is a composite number. Thus a composite number is one which has at least one factor greater than 1 but less than the number itself. In lower classes, you have also studied factorisation of composite numbers by using factor tree or successive division. Let us consider some large number, say 291060 and factorise it into prime factors by means of successive division as follows:

$$\begin{array}{r}
 2 \overline{)291060} \\
 \underline{2 \ 145530} \\
 3 \overline{)72765} \\
 \underline{3 \ 24255} \\
 3 \overline{)8085} \\
 \underline{5 \ 2695} \\
 7 \overline{)539} \\
 \underline{7 \ 77} \\
 11
 \end{array}$$

$$\begin{aligned}
 \therefore 291060 &= 2 \times 2 \times 3 \times 3 \times 3 \times 5 \times 7 \times 7 \times 11 \\
 &= 2^2 \times 3^3 \times 5 \times 7^2 \times 11
 \end{aligned}$$

There are in all 9 prime factors in the factorisation and the distinct prime factors are 2,3,5,7 and 11. Also we see that, in the factorisation the factor 3 is repeated thrice, the factors 2 and 7 are each repeated twice while the factors 5 and 11 are non-repeating.

This factorisation of the composite number 291060 is unique except for the order in which the factors occur. Will such a factorisation exist for each composite number? This question is answered in the affirmative by the following theorem which looks so simple inspite of its basic importance.

### Theorem 1.2

#### [Fundamental Theorem of Arithmetic or Unique Factorisation Theorem]

**Every composite number can be expressed as a product of primes uniquely except for the order of the factors.**

This theorem is often stated in the following alternative form also.

Every integer  $n > 1$  can be expressed uniquely in the form

$$n = p_1^{a_1} p_2^{a_2} \dots p_k^{a_k} \quad (1)$$

where  $p_1, p_2, \dots, p_k$  are primes such that  $p_1 < p_2 < \dots < p_k$  and  $a_1, a_2, \dots, a_k$  are all positive integers.



**Example 10.** Using the prime factorisation method find the HCF and LCM of 308, 336 and 420.

**Solution :** Canonical decomposition of the numbers gives

$$308 = 2^2 \times 7 \times 11$$

$$336 = 2^4 \times 3 \times 7$$

$$420 = 2^2 \times 3 \times 5 \times 7$$

The primes common to all the numbers are 2, 7 and the minimum powers to which they are raised are 2, 1 respectively. Hence

$$(308, 336, 420) = 2^2 \times 7 = 28.$$

The primes which appear in the three numbers are 2, 3, 5, 7, 11 and the maximum powers to which they are raised are 4, 1, 1, 1, 1 respectively.

Hence

$$[308, 336, 420] = 2^4 \times 3 \times 5 \times 7 \times 11 = 18480.$$

Some other applications, direct or indirect of the unique factorisation theorem are illustrated in the following examples.

**Example 11.** For any natural number  $n$ , prove that the digital root of  $5^n$  cannot be a multiple of 3.

**Solution :** Since the only prime involved in the canonical form  $5^n$  is 5, therefore the only prime that divides  $5^n$  is 5. It means that  $5^n$  is not divisible by any prime other than 5. In particular  $5^n$  is not divisible by 3.

We also know that any number whose digital root is a multiple of 3, is divisible by 3.

It therefore follows that the digital root of  $5^n$  cannot be a multiple of 3 for any natural number  $n$ .

**Example 12.** Find any six consecutive composite numbers less than 300.

**Solution :** Consider the primes 2, 3, 5, 7 whose product is 210.

Clearly the six consecutive numbers

$$2 \times 3 \times 5 \times 7 + 2 = 212$$

$$2 \times 3 \times 5 \times 7 + 3 = 213$$

$$2 \times 3 \times 5 \times 7 + 4 = 214$$

$$2 \times 3 \times 5 \times 7 + 5 = 215$$

$$2 \times 3 \times 5 \times 7 + 6 = 216$$

$$2 \times 3 \times 5 \times 7 + 7 = 217$$

are composite, the first, third and fifth being divisible by 2 and the second, fourth and sixth by 3, 5 and 7 respectively.

**EXERCISE 1.2**

- Find the canonical decomposition of the numbers :  
(i) 1914      (ii) 2332      (iii) 4284  
(iv) 190575      (v) 133848      (vi) 217350
- Find  $(a, b)$ ,  $[a, b]$  and verify that  $(a, b)[a, b] = ab$  for each of the following pairs of integers :  
(i)  $a = 429, b = 715$       (ii)  $a = 756, b = 1044$       (iii)  $a = 576, b = 2520$
- Find the HCF and LCM of the following integers by prime factorisation method  
(i) 204, 1020, 1190      (ii) 126, 882, 1617  
(iii) 504, 2394, 4725      (iv) 1260, 1800, 3780, 7560
- The HCF and LCM of two numbers are 27 and 29295 respectively. If one number is 837, find the other.
- Show that for any natural number  $n$ , the digit in units place of  $3^n$  cannot be even.
- Find any five consecutive composite numbers.
- Find any four consecutive odd composite numbers.
- Find the least number which when divided by 24, 36 and 60 will leave in each case the same remainder 7.
- Find the least number which when divided by 7, 8, and 12 leaves the same remainder 5 in each case.
- Find the least multiple of 13 which when divided by 5, 8 and 12 leaves the same remainder 2 in each case.
- By what prime numbers may 319 be divided so that the remainder is 4?
- By what numbers may 27 be divided so that the remainder is 3.?

**ANSWER**

- |  |  |   |
|--|--|---|
| 1. (i) $2 \times 3 \times 11 \times 29$    | (ii) $2^2 \times 11 \times 53$             | (iii) $2^2 \times 3^2 \times 7 \times 17$         |
| (iv) $3^2 \times 5^2 \times 7 \times 11^2$ | (v) $2^3 \times 3^2 \times 11 \times 13^2$ | (vi) $2 \times 3^3 \times 5^2 \times 7 \times 23$ |
| 2. (i) 143, 2145                           | (ii) 36, 21924                             | (iii) 72, 20160                                   |
| 3. (i) 34, 7140                            | (ii) 21, 9702                              | (iii) 63, 718200                                  |
| (iv) 180, 37800                            |  |   |
| 4. 945                                     | 6. 32, 33, 34, 35, 36                      |   |
| 7. 213, 215, 217, 219                      | 8. 367                                     |   |
| 9. 173                                     | 10. 962                                    |   |
| 11. 5, 7                                   | 12. 4, 6, 8, 12, 24                        |   |

**1.4 Field Properties of Real Numbers**

**Proof :**  $x + y = x + z$  .....(1)

Since,  $x \in \mathbb{R}$ ,  $-x \in \mathbb{R}$  (existence of additive inverse). Now adding  $-x$  to both sides of (1) we get

$$(-x) + (x + y) = (-x) + (x + z)$$

$$\Rightarrow (-x + x) + y = (-x + x) + z \quad (\text{by associativity of addition})$$

$$\Rightarrow 0 + y = 0 + z \quad (\text{by property of additive inverse})$$

$$\Rightarrow y = z \quad (\text{by property of additive identity})$$

**Corollary 2. (Cancellation law for multiplication)**

If  $x, y, z \in \mathbb{R}$ ,  $x \neq 0$  and

$xy = xz$ , then  $y = z$ .

**Proof :** Since  $x \neq 0$ , by existence of multiplicative inverse,  $\frac{1}{x} \in \mathbb{R}$

Thus  $xy = xz$

$$\Rightarrow \frac{1}{x}(xy) = \frac{1}{x}(xz)$$

$$\Rightarrow \left(\frac{1}{x} \times x\right)y = \left(\frac{1}{x} \times x\right)z \quad (\text{by associativity of multiplication})$$

$$\Rightarrow 1.y = 1.z \quad (\text{by property of multiplicative inverse})$$

$$\Rightarrow y = z \quad (\text{by property of multiplicative identity})$$

**Corollary 3. For any  $x \in \mathbb{R}$ ,  $x.0 = 0$**

**Proof :**  $0 + 0 = 0$  (by property of additive identity)

$$\Rightarrow x.(0 + 0) = x.0$$

$$\Rightarrow x.0 + x.0 = x.0 \quad (\text{by distributive law})$$

$$\Rightarrow x.0 + x.0 = x.0 + 0 \quad (\text{by property of additive identity})$$

$$\Rightarrow x.0 = 0 \quad (\text{by cancellation law})$$

**Corollary 4. For  $x, y \in \mathbb{R}$ ,  $x(-y) = -xy$**

**Definition :** The absolute value or modulus of a real number  $x$ , denoted by

$|x|$  is defined by

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Thus  $|0| = 0$ ,  $|3| = 3$ ,  $|-3| = 3$ ,  $|\sqrt{5}| = \sqrt{5}$ ,  $|-\sqrt{5}| = \sqrt{5}$  etc.

**Observe the following facts :**

- (i) If  $x = 0$ , then  $x = -x = 0$
- (ii) If  $x > 0$ , then  $-x < 0$  and  $x > -x$
- (iii) If  $x < 0$ , then  $-x > 0$  and  $-x > x$

Keeping these facts in mind, we can also define  $|x|$  as follows:

$$|x| = \begin{cases} 0, & \text{if } x = 0 \\ \text{the greater of } x \text{ and } -x, & \text{if } x \neq 0 \end{cases}$$

It is readily seen from the last definition that, for any  $x \in R$

$$|x| \geq x \text{ and } |x| \geq -x$$

Some fundamental properties concerning absolute values of real numbers are given below. In the following,  $x, y$  are any real numbers and  $\delta$ , a positive real number.

- (1)  $|x| \geq 0$
- (2)  $|-x| = |x|$
- (3)  $|xy| = |x||y|$
- (4)  $|x + y| \leq |x| + |y|$
- (5)  $|x - y| \geq |x| - |y|$  and  $|x - y| \geq |y| - |x|$
- (6)  $|x - y| < \delta$  if and only if  $y - \delta < x < y + \delta$

**Proof:** (1)  $|x| \geq 0$

**Case I.** If  $x = 0$ , then  $|x| = 0$

**Case II.** If  $x > 0$ , then  $|x| = x > 0$

**Case III.** If  $x < 0$ , then  $-x > 0$  and  $|x| = -x > 0$

Thus in all cases,  $|x| \geq 0$  holds.

$$(5) \quad |x - y| \geq |x| - |y| \text{ and } |x - y| \geq |y| - |x|$$

We have  $|x| = |(x - y) + y| \leq |x - y| + |y|$ , by (4)

$$\Rightarrow |x| - |y| \leq |x - y|$$

$$\text{i.e. } |x - y| \geq |x| - |y|$$

Similarly,  $|y - x| \geq |y| - |x|$

$$\Rightarrow |x - y| \geq |y| - |x| \quad \therefore |y - x| = |x - y| \text{ by (2)}$$

$$(6) \quad |x - y| < \delta \text{ if and only if } y - \delta < x < y + \delta$$

Suppose  $|x - y| < \delta$

$$\text{Then } x - y \leq |x - y| < \delta$$

$$\Rightarrow x - y < \delta$$

$$\Rightarrow x < y + \delta \text{ -----(i)}$$

$$\text{and } -(x - y) \leq |x - y| < \delta$$

$$\Rightarrow -x + y < \delta$$

$$\Rightarrow y < x + \delta$$

$$\Rightarrow y - \delta < x \text{ -----(ii)}$$

Combining (i) and (ii),  $y - \delta < x < y + \delta$

Conversely suppose  $y - \delta < x < y + \delta$

$$\text{Then } y - \delta < x \Rightarrow y - x < \delta$$

$$\Rightarrow -(x - y) < \delta$$

$$\text{And } x < y + \delta \Rightarrow x - y < \delta$$

Hence  $|x - y| =$  the greater of  $x - y$  and  $-(x - y) < \delta$

as both  $x - y$  and  $-(x - y)$  are less than  $\delta$

$$\text{i.e. } |x - y| < \delta$$

$\therefore |a| + |b|$  and  $|a| - |b|$  are either both positive or both negative.

But  $|a| + |b|$  being the sum of non-negative numbers cannot be negative.

Hence both  $|a| + |b|$  and  $|a| - |b|$  should be positive. In particular

$$|a| - |b| > 0$$

$$\therefore |a| > |b|$$

### EXERCISE 1.3

1. Examine whether the following statements are true or false:

- (i) The reciprocal of an irrational number is irrational.
- (ii) The set of natural numbers contains additive identity.
- (iii) The set of integers has multiplicative identity.
- (iv) The reciprocal of a non-zero rational number is rational.
- (v) The operation of subtraction in  $\mathbb{R}$  is commutative.
- (vi) The operation of division in  $\mathbb{R}$  is associative.

2. (a) Is there any real number  $x$  such that  $\frac{1}{x} \notin \mathbb{R}$ ?
- (b) Is there any  $x \in \mathbb{R}$  such that  $-x \notin \mathbb{R}$ ?
- (c) Is there any  $x \in \mathbb{R}$  such that  $x^2$  is not positive?
- (d) Is there any  $x \in \mathbb{R}$  such that  $x^2$  is negative?
- (e) Can two different real numbers have the same absolute value?

3. If  $|a| = |b|$ , find all possible relations between  $a$  and  $b$ .

4. Give any three values of  $x$  satisfying  $|x-3| < 1$ .

5. Find  $x$  if

(i)  $|x-2| = 0$     (ii)  $|x-2| = 3$     (iii)  $|x-3| = \sqrt{2}$     (iv)  $|x-2| = x$

6. If  $a^2 + b^2 = 0$ , prove that  $a = 0$  and  $b = 0$ .

7. Identify on the number line, the points  $x$  satisfying :

- (i)  $|x| \leq 3$     (ii)  $|x| < 3$     (iii)  $|x| \geq 3$
- (iv)  $|x| > 3$     (v)  $|x-2| \leq 3$

3. The Fundamental Theorem of Arithmetic or Unique Factorisation Theorem:

Every composite number can be expressed as a product of primes, uniquely except for the order of the factors or equivalently

Every integer  $n > 1$  can be expressed uniquely in the form

$$n = p_1^{a_1} p_2^{a_2} \dots p_k^{a_k}$$

where  $p_1, p_2, \dots, p_k$  are primes such that  $p_1 < p_2 < \dots < p_k$  and  $a_1, a_2, \dots, a_k$  are all positive integers.

4. Eleven properties of real numbers associated with the operations of addition and multiplication, known by the name 'Field properties of real numbers'.

5. The properties concerning absolute values of real numbers as listed below:

(1)  $|x| \geq 0$

(2)  $|-x| = |x|$

(3)  $|xy| = |x||y|$

(4)  $|x + y| \leq |x| + |y|$

(5)  $|x - y| \geq |x| - |y|$  and  $|x - y| \geq |y| - |x|$

(6)  $|x - y| < \delta$  if and only if  $y - \delta < x < y + \delta$ .

\*\*\*\*\*

©BOSEM Not to be republished



To start with, let us try when the divisor is a monomial. For example, let us divide the polynomial  $5x^3 - 2x^2 + x$  by the monomial  $x$ .

We have ,

$$\begin{aligned}(5x^3 - 2x^2 + x) \div x &= \frac{5x^3 - 2x^2 + x}{x} \\ &= \frac{5x^3}{x} - \frac{2x^2}{x} + \frac{x}{x} \\ &= 5x^2 - 2x + 1\end{aligned}$$

Then we express this fact as  $5x^3 - 2x^2 + x = x(5x^2 - 2x + 1) + 0$  and we say that  $x$  and  $5x^2 - 2x + 1$  are factors of  $5x^3 - 2x^2 + x$  and  $5x^3 - 2x^2 + x$  is a multiple of  $x$  as well as a multiple of  $5x^2 - 2x + 1$ .

Again, let us divide  $2x^3 + 4x + 5$  by  $2x$

$$\text{Here, } (2x^3 + 4x + 5) \div 2x = x^2 + 2 + \frac{5}{2x}$$

We see that  $\frac{5}{2x}$  cannot be a term of any polynomial. So, in this case we stop the process of division keeping 5 as the remainder.

Therefore,

$$2x^3 + 4x + 5 = 2x(x^2 + 2) + 5.$$

i.e. Dividend = Divisor  $\times$  Quotient + Remainder.

Observe that the degree of the remainder 5 is 0, which is less than that of the divisor  $2x$ , which is 1.

Again consider the division of  $7x + 3x^2 - 4$  by  $3 + x$ .

**We carry out the division by means of the following steps.**

**Step 1:** We write the dividend and the divisor after arranging the terms in the descending order of their degrees. Then the dividend is  $3x^2 + 7x - 4$  and divisor is  $x + 3$ .

**Step 2 :** We divide the first term of the dividend by the first term of the divisor i.e., we divide  $3x^2$  by  $x$  and we get  $3x$ . This gives the first term of the quotient.

**Step 3 :** We multiply the divisor  $x + 3$  by  $3x$  (the first term of the quotient) and obtain the product as  $3x^2 + 9x$ . Subtract this product from the dividend  $3x^2 + 7x - 4$  and we get the remainder as  $-2x - 4$ .

$x+3 \overline{) 3x^2+7x-4}$	$\frac{3x-2}{x}$	First term of quotient
$\underline{3x^2+9x}$	$= \frac{3x^2}{x} = 3x$	
$\underline{-2x-4}$	$\frac{-2x-6}{x}$	Second term of quotient
$\underline{-2x-6}$	$= \frac{-2x}{x} = -2$	
$\underline{2}$		

$$p(x) = d(x) \times q(x) + r(x)$$

where either  $r(x) = 0$  or  $\text{degree of } r(x) < \text{degree of } d(x)$ .

This result is known as the Division Algorithm for polynomials.

**Example 1.** Divide  $3x^2 - x^3 + 2x + 1$  by  $2x + 1 - x^2$ , and verify the division algorithm.

**Solution:** To carry out the division process, we first write the terms of both the dividend and divisor in descending order of their degrees. So, dividend  $= -x^3 + 3x^2 + 2x + 1$  and divisor  $= -x^2 + 2x + 1$ . Division process is shown below:

$$\begin{array}{r} x-1 \\ -x^2+2x+1 \overline{) -x^3+3x^2+2x+1} \\ \underline{-x^3+2x^2+x} \phantom{+1} \\ x^2+x+1 \\ \underline{x^2-2x-1} \\ 3x+2 \end{array}$$

We see that the degree of  $3x+2$  is less than the degree of the divisor  $-x^2 + 2x + 1$ . So, we cannot continue the division any further.

Then, quotient  $= x - 1$  and remainder  $= 3x + 2$

Now, Divisor  $\times$  Quotient + Remainder

$$\begin{aligned} &= (-x^2 + 2x + 1)(x - 1) + (3x + 2) \\ &= -x^3 + 2x^2 + x + x^2 - 2x - 1 + 3x + 2 \\ &= -x^3 + 3x^2 + 2x + 1 \\ &= \text{Dividend} \end{aligned}$$

Hence verified.

## EXERCISE 2.1

1. Divide the polynomial  $p(x)$  by the polynomial  $d(x)$  and find the quotient and remainder, and verify division algorithm in each of the following:

(i)  $p(x) = 2x^3 + x^2 - x$ ,  $d(x) = x$

(ii)  $p(x) = 3x^2 - x - 1$ ,  $d(x) = -x$

(iii)  $p(x) = 3x^2 + 2x + 1$ ,  $d(x) = x + 1$

(iv)  $p(x) = x^3 - 1$ ,  $d(x) = x - 1$

(v)  $p(x) = 2x^2 + 3x + 1$ ,  $d(x) = 2 + x$

(vi)  $p(x) = x^3 - 3x^2 + 5x - 3$ ,  $d(x) = x^2 - 2$

(vii)  $p(x) = 4x^3 + 4x^2 - x + 1$ ,  $d(x) = x^2 + 2x$

(viii)  $p(x) = 3x^2 - x^3 - 3x + 5$ ,  $d(x) = x - 1 - x^2$

### 2.3 Remainder Theorem

From the previous article, we know that if  $p(x)$  and  $d(x)$  are two polynomials with  $d(x) \neq 0$ , then we can divide  $p(x)$  by  $d(x)$  to get a polynomial  $q(x)$  as quotient and a remainder  $r(x)$ . The degree of  $r(x)$ , if it is non-zero, is definitely less than the degree of the divisor  $d(x)$ . Further,

$$\text{Dividend} = \text{Divisor} \times \text{Quotient} + \text{Remainder}$$

$$\text{i.e. } p(x) = d(x) \times q(x) + r(x) \text{ where } r(x) = 0 \text{ or degree of } r(x) \text{ is less than that of } d(x).$$

This relation is true for all values of  $x$ .

**Note: In case degree of the dividend  $p(x)$  is less than that of the divisor, then we take  $q(x) = 0$  and  $r(x) = p(x)$ .**

If the divisor  $d(x)$  is a linear polynomial i.e. degree of  $d(x)$  is 1, then  $r(x)$  will be a constant. Let it be denoted by  $R$ .

Then the relation takes the form :

$$p(x) = d(x) \times q(x) + R$$

To illustrate the above discussion, let us divide the polynomial  $p(x) = 5x^2 + 2x - 3$  by the linear polynomial  $x - 2$ . By long division, we have :

$$\begin{array}{r} 5x + 12 \\ x - 2 \overline{) 5x^2 + 2x - 3} \\ \underline{5x^2 - 10x} \phantom{- 3} \\ 12x - 3 \\ \underline{12x - 24} \\ 21 \end{array}$$

Hence, we see that the remainder is 21, which is a constant. Now, let us investigate whether there is any link between the constant remainder and value of the dividend corresponding to certain value of  $x$ .

The zero of the divisor  $x - 2$  is 2. If we replace  $x$  by 2 in  $p(x) = 5x^2 + 2x - 3$ , we have

$$\begin{aligned} p(2) &= 5 \times 2^2 + 2 \times 2 - 3 \\ &= 20 + 4 - 3 \\ &= 21, \text{ which is the remainder.} \end{aligned}$$

**Example 2.** Find the remainder when  $x^4 + 2x^3 - 3x^2 - 5x + 4$  is divided by  $x-3$ , without actual division.

**Solution :** Let  $p(x) = x^4 + 2x^3 - 3x^2 - 5x + 4$

The zero of  $x-3$  is 3.

Hence, by Remainder Theorem, the remainder when  $p(x)$  is divided by  $x-3$

$$\begin{aligned} &= p(3) \\ &= 3^4 + 2 \times 3^3 - 3 \times 3^2 - 5 \times 3 + 4 \\ &= 81 + 54 - 27 - 15 + 4 \\ &= 97. \end{aligned}$$

**Example 3.** Check whether the polynomial  $4x^3 + 4x^2 - x - 1$  is a multiple of  $2x + 1$ .

**Solution :** Let  $p(x) = 4x^3 + 4x^2 - x - 1$

The zero of  $2x + 1$  is  $-\frac{1}{2}$

So, the remainder when  $p(x)$  is divided by  $2x + 1$

$$\begin{aligned} &= p\left(-\frac{1}{2}\right) \\ &= 4 \times \left(-\frac{1}{2}\right)^3 + 4 \times \left(-\frac{1}{2}\right)^2 - \left(-\frac{1}{2}\right) - 1 \\ &= -\frac{1}{2} + 1 + \frac{1}{2} - 1 = 0 \end{aligned}$$

Hence,  $4x^3 + 4x^2 - x - 1$  is a multiple of  $2x + 1$ .

## 2.4 Factor Theorem

Factor Theorem can be deduced from the Remainder Theorem as a particular case.

**Statement :** If  $p(x)$  is a polynomial of degree  $\geq 1$  and  $a$  is any real number, then  $x - a$  is a factor of  $p(x)$  if and only if  $p(a) = 0$ .

**Proof :** Let us suppose that  $p(a) = 0$ .

By Remainder Theorem, we know that  $p(a)$  is the remainder when  $p(x)$  is divided by  $x - a$ .

$$\begin{aligned} \therefore p(x) &= (x - a)q(x) + p(a) \\ &= (x - a)q(x) \quad [\because p(a) = 0] \end{aligned}$$

Hence  $x - a$  is a factor of  $p(x)$ .

Conversely, let us suppose that  $x - a$  is a factor of  $p(x)$ .

## EXERCISE 2.2

1. Find without actual division, the remainder when  $x^3+3x^2+3x+1$  is divided by
- (i)  $x - 1$       (ii)  $x+1$       (iii)  $x - \frac{1}{2}$       (iv)  $2x+1$
2. Determine whether  $x + 1$  is a factor of :
- (i)  $x^3 + x^2 + x + 1$       (ii)  $x^4 + x^3 + x^2 + x + 1$
- (iii)  $x^4 + 3x^3 + 3x^2 + x + 1$       (iv)  $x^3 + 3x^2 + (2 + \sqrt{2})x + \sqrt{2}$
3. Use Factor Theorem to determine whether  $q(x)$  is a factor of  $p(x)$  in each of the following cases:
- (i)  $p(x) = x^4 - 81$ ,       $q(x) = x + 3$
- (ii)  $p(x) = 2x^3 + x^2 - 2x - 1$ ,       $q(x) = x + 1$
- (iii)  $p(x) = x^3 - 3x^2 - 3x + 1$ ,       $q(x) = x - 1$
- (iv)  $p(x) = x^3 - 4x^2 + x + 6$ ,       $q(x) = x - 4$
4. Using Factor Theorem, show that
- (i)  $x - 1$  is a factor of  $3x^5 - 2x^2 - 6x + 5$
- (ii)  $x + 1$  is a factor of  $2x^4 - 3x^2 + 6x + 7$
- (iii)  $x - 2$  is a factor of  $x^3 - 9x^2 + 26x - 24$
- (iv)  $x + y, y + z, z + x$  are factors of  $(x + y + z)^3 - (x^3 + y^3 + z^3)$ .
- (v)  $x - y, y - z, z - x$  are factors of  $x^3(y^2 - z^2) + y^3(z^2 - x^2) + z^3(x^2 - y^2)$ .
5. Find the value of  $k$ , if  $x-1$  is a factor of  $p(x)$  in each of the following cases :
- (i)  $p(x) = x^2 - 2x + k$       (ii)  $p(x) = \sqrt{2}x^2 + kx - 1$
- (iii)  $p(x) = kx^2 - \sqrt{2}x + 1$       (iv)  $p(x) = kx^2 - 3x + 2k$
6. If  $x^2 + px + q$  and  $x^2 + lx + m$  are both divisible by  $x + a$ , show that
- $$a = \frac{m - q}{l - p}.$$

(ii) Let  $p(x) = x^3 - 2x^2 - x + 2$ .

The factors of 2 (the constant term in  $p(x)$ ) are  $\pm 1, \pm 2$ .

Now,  $p(1) = 1^3 - 2 \times 1^2 - 1 + 2 = 0$

So,  $x - 1$  is a factor of  $p(x)$ .

Again,  $p(-1) = (-1)^3 - 2 \times (-1)^2 - (-1) + 2 = 0$

$\therefore x + 1$  is also a factor of  $p(x)$

Dividing  $p(x)$  by the product of the two factors i.e.  $(x - 1)(x + 1)$  or  $x^2 - 1$ , we have :

$$\begin{array}{r} x-2 \\ x^2-1 \overline{) x^3-2x^2-x+2} \\ \underline{x^3 \phantom{-2x^2} -x} \phantom{+2} \\ -2x^2 + 2 \\ \underline{-2x^2 + 2} \\ 0 \end{array}$$

Hence,  $x^3 - 2x^2 - x + 2 = (x^2 - 1)(x - 2)$   
 $= (x - 1)(x + 1)(x - 2)$ .

(iii) Let  $p(x) = x^4 - 6x^3 + 13x^2 - 12x + 4$ .

The factors of 4 are  $\pm 1, \pm 2, \pm 4$ .

By trial, we find that  $p(1) = 0$ . So,  $x - 1$  is a factor of  $p(x)$ . Similarly, we can find that  $x - 2$  is also a factor of  $p(x)$ .

Then, we have two factors of  $p(x)$ , namely  $x - 1$  and  $x - 2$ . Dividing  $p(x)$  by the product of these two factors i.e.  $(x - 1)(x - 2) = x^2 - 3x + 2$ , we have :

$$\begin{array}{r} x^2-3x+2 \\ x^2-3x+2 \overline{) x^4-6x^3+13x^2-12x+4} \\ \underline{x^4-3x^3+2x^2} \phantom{+4} \\ -3x^3+11x^2-12x \phantom{+4} \\ \underline{-3x^3+9x^2-6x} \phantom{+4} \\ 2x^2-6x+4 \\ \underline{2x^2-6x+4} \\ 0 \end{array}$$

$\therefore x^4 - 6x^3 + 13x^2 - 12x + 4 = (x^2 - 3x + 2)(x^2 - 3x + 2)$

**ANSWER**

- |                           |                          |                       |
|---------------------------|--------------------------|-----------------------|
| 1. (i) $(x-1)(x-3)$       | (ii) $(x-2)(x-4)$        | (iii) $(x+3)(x+5)$    |
| (iv) $(x+1)(x-7)$         | (v) $(x+5)(x-2)$         | (vi) $(x-5)(x+1)^2$   |
| (vii) $(x+1)(x+2)(x+10)$  | (viii) $(x-1)(x-2)(x+3)$ |                       |
| (ix) $(x+1)(x+2)(x+4)$    | (x) $(x+1)(x+3)(x-4)$    | (xi) $(x+1)^2(x+2)^2$ |
| (xii) $(x-1)(x+1)(x^2+1)$ |                          |                       |
| 2. $(x+y)(y+z)(z+x)$      |                          |                       |

**SUMMARY**

**Main points studied in this chapter are given below :**

1. The division algorithm for polynomials states that if  $p(x)$  and  $d(x)$  are any two polynomials with  $d(x) \neq 0$ , then we can find polynomials  $q(x)$  and  $r(x)$  such that  $p(x) = d(x) \times q(x) + r(x)$ , where either  $r(x) = 0$  or degree of  $r(x) <$  degree of  $d(x)$ .
2. Remainder Theorem : If  $p(x)$  is any polynomial of degree greater than or equal to one and  $p(x)$  is divided by the linear polynomial  $x - a$ , then the remainder is  $p(a)$ .
3. Factor Theorem : If  $p(x)$  is a polynomial of degree  $\geq 1$  and  $a$  is any real number, then  $x - a$  is a factor of  $p(x)$  if and only if  $p(a) = 0$ .

\*\*\*\*\*

©BOSEM Not to be republished



### 3.3 Factorisation of Cyclic Expressions

In many cases, cyclic expressions can be factorised by using the following steps:

- (i) Write the terms of the expression according to the ascending or descending powers of one of the letters involved in the expression.
- (ii) Take out the factor(s) common to each coefficient.
- (iii) Write the terms of the other factor according to the ascending or descending powers of any letters other than the previous.
- (iv) Repeat the process till the factorisation is complete.

Now, let us illustrate the process by considering some cyclic expressions.

#### 3.4 To Factorise $a^2(b+c)+b^2(c+a)+c^2(a+b)+2abc$

$$\begin{aligned}
 & a^2(b+c)+b^2(c+a)+c^2(a+b)+2abc \\
 &= a^2(b+c)+b^2c+ab^2+c^2a+bc^2+2abc \\
 &= a^2(b+c)+a(b^2+c^2+2bc)+(b^2c+bc^2) \text{ [arranging in descending powers of } a] \\
 &= a^2(b+c)+a(b+c)^2+bc(b+c) \\
 &= (b+c) [a^2+a(b+c)+bc] \\
 &= (b+c)[b(c+a)+a(c+a)] \text{ [arranging in descending powers of } b] \\
 &= (b+c)(c+a)(b+a) \\
 &= (a+b)(b+c)(c+a)
 \end{aligned}$$

#### 3.5 To Factorise $a^2(b-c)+b^2(c-a)+c^2(a-b)$

$$\begin{aligned}
 & a^2(b-c)+b^2(c-a)+c^2(a-b) \\
 &= a^2(b-c)+b^2c-ab^2+c^2a-bc^2 \\
 &= a^2(b-c)-a(b^2-c^2)+(b^2c-bc^2) \text{ [arranging in descending powers of } a] \\
 &= a^2(b-c)-a(b+c)(b-c)+bc(b-c) \\
 &= (b-c)[a^2-a(b+c)+bc] \\
 &= (b-c)[b(c-a)-a(c-a)] \text{ [arranging in descending powers of } b] \\
 &= (b-c)(c-a)(b-a) \\
 &= -(a-b)(b-c)(c-a)
 \end{aligned}$$

**3.8 To Factorise**  $a^2(b+c)+b^2(c+a)+c^2(a+b)+3abc$ .

$$\begin{aligned} & a^2(b+c)+b^2(c+a)+c^2(a+b)+3abc \\ &= a^2b+ca^2+b^2c+ab^2+c^2a+bc^2+3abc \\ &= (a^2b+ab^2+abc)+(abc+b^2c+bc^2)+(ca^2+abc+c^2a) \\ &= ab(a+b+c)+bc(a+b+c)+ca(a+b+c) \\ &= (a+b+c)(ab+bc+ca) \end{aligned}$$

**Note :**

$$\begin{aligned} & a^2(b+c)+b^2(c+a)+c^2(a+b) \\ &= bc(b+c)+ca(c+a)+ab(a+b)\dots\dots\dots(1) \\ &= a(b^2+c^2)+b(c^2+a^2)+c(a^2+b^2)\dots\dots\dots(2) \end{aligned}$$

Therefore,  $a^2(b+c)+b^2(c+a)+c^2(a+b)$  can be replaced by either (1) or (2).

**3.9 To Factorise**  $a^3+b^3+c^3-3abc$ .

$$\begin{aligned} & a^3+b^3+c^3-3abc \\ &= (a+b)^3-3ab(a+b)+c^3-3abc [\because a^3+b^3=(a+b)^3-3ab(a+b)] \\ &= (a+b)^3+c^3-\{3ab(a+b)+3abc\} \\ &= (a+b+c)\{(a+b)^2-(a+b)c+c^2\}-3ab(a+b+c) \\ & \qquad \qquad \qquad [\because x^3+y^3=(x+y)(x^2-xy+y^2)] \\ &= (a+b+c)(a^2+2ab+b^2-ca-bc+c^2-3ab) \\ &= (a+b+c)(a^2+b^2+c^2-ab-bc-ca)\dots\dots\dots(1) \\ &= \frac{1}{2}(a+b+c)(2a^2+2b^2+2c^2-2ab-2bc-2ca) \\ &= \frac{1}{2}(a+b+c)\{(a^2-2ab+b^2)+(b^2-2bc+c^2)+(c^2-2ca+a^2)\} \\ &= \frac{1}{2}(a+b+c)\{(a-b)^2+(b-c)^2+(c-a)^2\}\dots\dots\dots(2) \end{aligned}$$

Hence,

$$\begin{aligned} & a^3+b^3+c^3-3abc \\ &= (a+b+c)(a^2+b^2+c^2-ab-bc-ca) \\ &= \frac{1}{2}(a+b+c)\{(a-b)^2+(b-c)^2+(c-a)^2\}. \end{aligned}$$

**3.10 To Factorise**  $2b^2c^2+2c^2a^2+2a^2b^2-a^4-b^4-c^4$ .

We have,  $(a+b+c)^2 = a^2+b^2+c^2+2ab+2bc+2ca$ .

$\therefore$  The given expression

3. Prove that  $(x - y)^3 + (y - z)^3 + (z - x)^3 - 3(x - y)(y - z)(z - x) = 0$
4. If  $a^3 + b^3 + c^3 = 3abc$ , prove that either  $a + b + c = 0$  or  $a = b = c$ .
5. If  $x + y + z = 9$ ,  $xy + yz + zx = 26$  and  $xyz = 24$ , find the value of  $x^2(y + z) + y^2(z + x) + z^2(x + y)$ .
6. If  $x + y - z = 2$ ,  $y + z - x = 4$  and  $z + x - y = 6$ , find the value of  $2y^2z^2 + 2z^2x^2 + 2x^2y^2 - x^4 - y^4 - z^4$ .
7. If  $x + y + z = 12$  and  $x^2 + y^2 + z^2 = 44$ , find the value of  $(x + y + z)^3 - x^3 - y^3 - z^3 + 3xyz$ .
8. Find the value of  $xy(x + y) + yz(y + z) + zx(z + x) + 3xyz$ , when  $x = a(b - c)$ ,  $y = b(c - a)$ ,  $z = c(a - b)$ .

## ANSWER

1.
  - (i)  $(x + y - z)(x^2 + y^2 + z^2 - xy + yz + zx)$
  - (ii)  $(a - b + 3)(a^2 + b^2 + 9 + ab + 3b - 3a)$
  - (iii)  $(2a + 3b + 4c)(4a^2 + 9b^2 + 16c^2 - 6ab - 12bc - 8ca)$
  - (iv)  $(x - y - 5z)(x^2 + y^2 + 25z^2 + xy - 5yz + 5zx)$
  - (v)  $(a^2 - a + 2)(a^4 + a^3 - a^2 + 2a + 4)$
  - (vi)  $(x^2 - x + 3)(x^4 + x^3 - 2x^2 + 3x + 9)$
2.
  - (i)  $-(x - y)(y - z)(z - x)$
  - (ii)  $(x + y)(y + z)(z + x)$
  - (iii)  $(a + b + c)(ab + bc + ca)$
  - (iv)  $(x + y)(y + z)(z + x)$
  - (v)  $(a + b + c)(ab + bc + ca)$
  - (vi)  $-(x - y)(y - z)(z - x)(x^2 + y^2 + z^2 + xy + yz + zx)$
  - (vii)  $-(x - y)(y - z)(z - x)(x^2 + y^2 + z^2 + xy + yz + zx)$
  - (viii)  $-(a + b)(b + c)(c + a)(a - b)(b - c)(c - a)$
  - (ix)  $-(x - y)(y - z)(z - x)(xy + yz + zx)$
  - (x)  $3(x + z)(y + 2z - x)(z - y)$

### 4.1 Introduction

In class IX, you have learnt about linear equations in two variables and their solutions. Recall that the general form of a linear equation in two variables is  $ax + by + c = 0$ , where  $a, b, c$  are constants and  $a^2 + b^2 \neq 0$ , i.e.  $a$  and  $b$  are not zero simultaneously. An ordered pair  $(x_1, y_1)$  which satisfies this equation is a solution of the equation and a linear equation in two variables has infinitely many solutions. You also know that the graph of a linear equation in two variables is a straight line whose points make up the aggregate of solutions of the equation. In this chapter, we shall discuss about the solution of a pair of linear equations in two variables.

### 4.2 Solution of a Pair of Linear Equations in Two Variables

Let us consider the statement “The sum of two numbers is 9 and their difference is 5”. If we denote the two numbers by  $x$  and  $y$  ( $x > y$ ), then the statement can be expressed as  $x + y = 9$  and  $x - y = 5$ .

Thus, the statement gives a pair of linear equations in two variables ( $x$  and  $y$ ). Some other examples of pair of linear equations in two variables are

$$3x + 2y - 1 = 0 \text{ and } 2x + 3y + 4 = 0;$$

$$x = 5y \text{ and } -2x + 4y + 7 = 0;$$

$$x - y = 3 \text{ and } 3x - y - 11 = 0 \text{ etc.}$$

The general form of a pair of linear equations in two variables  $x$  and  $y$  is

$$a_1x + b_1y + c_1 = 0$$

$$\text{and } a_2x + b_2y + c_2 = 0$$

where  $a_1, b_1, c_1, a_2, b_2, c_2$  are all real numbers and  $a_1^2 + b_1^2 \neq 0, a_2^2 + b_2^2 \neq 0$ . If  $x = x_1, y = y_1$  satisfy both the linear equations, then the ordered pair  $(x_1, y_1)$  is called a solution of the pair of linear equations.

$x$	0	5	-1
$y$	5	0	6

Table (I)

Similarly, some of the values of  $x$  and  $y$  satisfying equation (2) are given in the Table (II) below.

$x$	0	4	-2
$y$	6	0	9

Table (II)

The points given by Table (I) namely  $(0,5)$ ,  $(5,0)$  and  $(-1, 6)$  are plotted in the Cartesian plane.

Joining these points, we get the straight line  $AB$ . This is the graph of the equation  $x+y=5$  (Fig. 4.1).

Again, in the same Cartesian plane and taking the same units the points given by Table (II) i.e.,  $(0,6)$ ,  $(4,0)$  and  $(-2, 9)$  are plotted. Joining these points we get the straight line  $CD$ . This is the graph of the equation  $3x + 2y = 12$  (Fig. 4.1).




Fig. 4.1

It is seen from the Fig. 4.1 that the two straight lines  $AB$  and  $CD$  intersect at the point  $P(2,3)$ . This is a point common to both the straight lines  $AB$  and  $CD$ . Therefore, the coordinates of  $P$  will satisfy both the equations (1) and (2).

Hence, the solution of the given pair of equations is given by

$$x = 2 \text{ and } y = 3.$$

**Example 3.** Solve graphically  $2x + 3y = 6$   
and  $2x + 3y = 12$ .

**Solution :**  $2x + 3y = 6$  .....(1)  
 $2x + 3y = 12$  .....(2)

Some values of  $x$  and  $y$  satisfying equation (1) are given in Table (I).


$x$	0	3	-3
$y$	2	0	4

Table (I)

Some values of  $x$  and  $y$  satisfying equation (2) are given in Table (II)

$x$	6	0	3
$y$	0	4	2

Table (II)



When we draw the graph of equations (1) and (2) with the points given by Table (I) and (II) we get two parallel straight lines AB and CD (Fig. 4.3). That is, the points lying on AB are completely different from those lying on CD. In other words, the two lines AB and CD have no point in common. Hence, the given pair of equations has no solution.

Let us now write the pair of linear equations in Examples 1, 2, and 3 in the general form

$$a_1x + b_1y + c_1 = 0$$

and  $a_2x + b_2y + c_2 = 0$

and compare the values of  $\frac{a_1}{a_2}, \frac{b_1}{b_2}$  and  $\frac{c_1}{c_2}$  in each.

(iii) is inconsistent if  $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$ .

**Example 4. Examine whether the lines representing the pair of linear equations**

$$5x - 4y + 8 = 0 \text{ and}$$

$$7x + 6y - 9 = 0, \text{ intersect at a point, are parallel or coincident.}$$

**Solution :** Comparing the given pair of equations with the general form, we get

$$a_1 = 5, \quad b_1 = -4, \quad c_1 = 8$$

$$\text{and } a_2 = 7, \quad b_2 = 6, \quad c_2 = -9.$$

$$\therefore \frac{a_1}{a_2} = \frac{5}{7}, \quad \frac{b_1}{b_2} = \frac{-4}{6}, \quad \frac{c_1}{c_2} = \frac{8}{-9}$$

$$\text{We see that, } \frac{a_1}{a_2} \neq \frac{b_1}{b_2}.$$

Hence, the two lines representing the given pair of equations intersect at a point.

**Example 5. Examine whether the pair of linear equations**

$$\frac{4}{3}x + 2y = 9$$

$$\text{and } 2x + 3y = 12 \text{ is consistent or not.}$$

**Solution :** Writing the given pair of equations in the general form, we get

$$\frac{4}{3}x + 2y - 9 = 0$$

$$2x + 3y - 12 = 0$$

$$\text{Here, } \frac{a_1}{a_2} = \frac{\frac{4}{3}}{2} = \frac{2}{3}, \quad \frac{b_1}{b_2} = \frac{2}{3} \text{ and } \frac{c_1}{c_2} = \frac{-9}{-12} = \frac{3}{4}.$$

$$\text{We see that, } \frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}.$$

Hence the given pair of equations is inconsistent.

## EXERCISE 4.1

1. By comparing the ratios of the coefficients, find out whether the lines representing the following pair of linear equations intersect at a point, are parallel or coincident:

(i)  $3x + 2y - 5 = 0$                       (ii)  $9x + 3y + 4 = 0$   
 $4x - 3y + 2 = 0$                        $18x + 6y + 8 = 0$

(iii)  $2x + y = 3$                       (iv)  $6x - 2y + 5 = 0$   
 $3x - 2y + 6 = 0$                        $3x - y + 9 = 0$

(v)  $\frac{3}{2}x + \frac{5}{3}y = 7$                       (vi)  $x + y = 5$   
 $9x + 10y = 14$                        $3x - 5y = -9$

2. By comparing the ratios of the coefficients, find out whether the following pairs of linear equations are consistent or not:

(i)  $2x + y - 3 = 0$                       (ii)  $3x + 4y = 12$   
 $3x + 4y + 6 = 0$                        $6x + 8y = 24$

(iii)  $3x - y + 5 = 0$                       (iv)  $2x - 3y = 8$                       (v)  $\frac{2}{3}x + 2y = 8$   
 $4x + 3y = 11$                        $4x - 6y = 9$                        $x + 3y = 2$

3. Solve the following pair of equations graphically:

(i)  $x - 3y + 6 = 0$                       (ii)  $2x - y = 2$                       (iii)  $2x + 3y = 5$   
 $x - 3y - 12 = 0$                        $3x + 2y = 17$                        $5x - 4y + 22 = 0$

(iv)  $y = x$                       (v)  $x + 4y = 2$                       (vi)  $4x + 6y = 18$   
 $5x - 2y = 9$                        $3x + 12y = 6$                        $9 - 2x - 3y = 0$

(vii)  $2x + y = 7$                       (viii)  $x + 2y = 8$                       (ix)  $3x + y = 9$   
 $3x + 2y = 11$                        $3x - y = 3$                        $2x - 3y + 16 = 0$

(x)  $\frac{x}{2} + \frac{y}{5} = 1$                       (xi)  $2x + 3y = 12$                       (xii)  $3x + 2y = 4$   
 $5x + 2y = 10$                        $2x = 3y$                        $6x + 4y = 13$



**4.4 Algebraic Method of Solving a Pair of Linear Equations**

In the previous section, we learnt how to solve a pair of linear equations in two variables graphically. This method is not convenient in cases when the point representing the solution of

the pair of equations has non-integral co-ordinates like  $\left(\frac{1}{11}, \frac{2}{13}\right), (2.73, -3.41), (-\sqrt{2}, \sqrt{5})$

etc. as there is difficulty in reading such co-ordinates accurately from the graph. To avoid such disadvantages of graphical method, we shall now discuss algebraic methods.

**4.4.1 Substitution Method**

To find the solution of a pair of linear equations in two variables by substitution method, we are to follow the following steps:

**Step 1.** From either equation, whichever is convenient, find the value of one variable, say  $y$  in terms of the other variable i.e.  $x$ .

**Step 2.** Substitute the value of  $y$  thus obtained in Step 1 in the other equation and reduce it to an equation in only one variable  $x$ , which can be solved; and hence obtain the value of  $x$ .

Sometimes after substitution you may get an equality relation with no variable. If this relation is true, you can conclude that the pair of linear equations has infinitely many solutions. If the relation is false, then the pair of linear equation is inconsistent.

**Step 3.** Substitute the value of  $x$  obtained in Step 2 in the equations of Step 1 and obtain the value of  $y$ .

This method is illustrated in the following examples.

**Example 8. Solve the following pair of equations by substitution method:**

$$2x + 3y = 32$$

$$11y - 9x = 3$$

**Solution :** Rewriting the given equations, we have

$$2x + 3y = 32 \dots\dots\dots(1)$$

$$11y - 9x = 3 \dots\dots\dots(2)$$

**Step 1.** From (1), we have

$$y = \frac{32 - 2x}{3} \dots\dots\dots(3)$$

$$x = \frac{3y + 5}{2}$$

Substituting this value of  $x$  in (2), we have

$$4\left(\frac{3y + 5}{2}\right) - 6y = 10$$

$$\Rightarrow 2(3y + 5) - 6y = 10$$

$$\Rightarrow 6y + 10 - 6y = 10$$

$$\Rightarrow 10 = 10, \text{ which is true.}$$

Hence the given pair of equations has infinitely many solutions.

**4.4.2 Elimination Method**

To solve a pair of linear equations in two variables by elimination method, we are to follow the following steps:

- Step 1.** Multiply or divide both the equations by suitable non-zero constants so that the coefficients of one variable (either  $x$  or  $y$ ) become numerically equal.
- Step 2.** Then add one equation to the other or subtract one from the other so that one variable gets eliminated. If you get an equation in one variable, go to Step 3.  
 If we obtain a true equality relation involving no variable, then the original pair of equations has infinitely many solutions.  
 If we obtain a false relation involving no variable, then the original pair of equations has no solution.
- Step 3.** Solve the equation obtained in step 2 and obtain the value of the variable which is not eliminated.
- Step 4.** Substitute the value of the variable obtained in Step 3 in any of the given equations to get the value of the other variable.

The following examples will illucidate the process.

**Example 10.** Solve the following pair of equations by elimination method:

$$3x + 2y = 13$$

$$7x - 5y = 11$$

**Solution :** Rewriting the given equations , we have

$$3x + 2y = 13 \dots\dots\dots(1)$$

$$7x - 5y = 11 \dots\dots\dots(2)$$

Multiplying (1) by  $b_2$  and (2) by  $b_1$ , and then subtracting, we get

$$(a_1b_2 - a_2b_1)x + (c_1b_2 - c_2b_1) = 0$$

$$\Rightarrow (a_1b_2 - a_2b_1)x = b_1c_2 - b_2c_1$$

$$\Rightarrow x = \frac{b_1c_2 - b_2c_1}{a_1b_2 - a_2b_1}, \text{ provided } a_1b_2 - a_2b_1 \neq 0 \quad \dots\dots\dots(3)$$

Again, multiplying (1) by  $a_2$  and (2) by  $a_1$ , and then subtracting, we get

$$(b_1a_2 - b_2a_1)y + (c_1a_2 - c_2a_1) = 0$$


$$\Rightarrow (a_1b_2 - a_2b_1)y = c_1a_2 - c_2a_1$$

$$\Rightarrow y = \frac{c_1a_2 - c_2a_1}{a_1b_2 - a_2b_1}, \text{ provided } a_1b_2 - a_2b_1 \neq 0 \quad \dots\dots\dots(4)$$

Hence, from (3) and (4), we have

$$\frac{x}{b_1c_2 - b_2c_1} = \frac{y}{c_1a_2 - c_2a_1} = \frac{1}{a_1b_2 - a_2b_1} \quad \dots\dots\dots(5)$$

The above result can be remembered from the following diagram :



The arrow between two numbers indicates that they are to be multiplied. In finding the corresponding denominator, the numbers with downward arrow are multiplied first, and then from their product, the product of the numbers with upward arrow is to be subtracted.

To find the solution of a pair of linear equations in two variables, we will follow the following steps :

- Step 1.** Write the given equations in the general form.
- Step 2.** Taking the help of the above diagram, write equations as given in (5).
- Step 3.** From the equations obtained in Step 2, find the values of  $x$  and  $y$ , provided

$$a_1b_2 - a_2b_1 \neq 0.$$

**Solution :** The given equations are written as

$$ax + by - c = 0$$

$$ax^2 + b^2y - c^2 = 0$$

Then by cross-multiplication, we have

$$\frac{x}{b(-c^2) - b^2(-c)} = \frac{y}{(-c)a^2 - (-c^2)a} = \frac{1}{ab^2 - a^2b}$$

$$\Rightarrow \frac{x}{bc(b-c)} = \frac{y}{ca(c-a)} = \frac{1}{ab(b-a)}$$

$$\Rightarrow x = \frac{c(b-c)}{a(b-a)}, y = \frac{c(c-a)}{b(b-a)}$$

**Example 14.** Solve the following pair of equations by reducing them to a pair of linear equations:

$$\frac{14}{x+y} + \frac{3}{x-y} = 5$$

$$\frac{21}{x+y} - \frac{1}{x-y} = 2$$

**Solution :** Writing  $u = \frac{1}{x+y}$  and  $v = \frac{1}{x-y}$ , the given equations are

$$\begin{cases} 14u + 3v = 5 \\ 21u - v = 2 \end{cases}$$

i.e.  $\begin{cases} 14u + 3v - 5 = 0 \\ 21u - v - 2 = 0 \end{cases}$

Then by cross-multiplication, we get

$$\frac{u}{3 \cdot (-2) - (-1) \cdot (-5)} = \frac{v}{(-5) \cdot 21 - (-2) \cdot 14} = \frac{1}{14 \cdot (-1) - 21 \cdot 3}$$

$$\Rightarrow \frac{u}{-6 - 5} = \frac{v}{-105 + 28} = \frac{1}{-14 - 63}$$

$$\Rightarrow \frac{u}{-11} = \frac{v}{-77} = \frac{1}{-77}$$

$$\therefore u = \frac{1}{7}$$

(vii)  $3x + 5y = 7$       (viii)  $4x - 3y = 10$       (ix)  $5x + 2y = 4$   
 $12x - 13y = -5$        $-5x + 4y = -13$        $10x + 4y = 8$

(x)  $2x - 5y = 6$       (xi)  $\frac{x}{2} + \frac{2y}{3} = -1$       (xii)  $3x - 5y + 2 = 0$   
 $4x - 10y = 9$        $\frac{x}{3} - \frac{y}{9} = 1$        $9x = 2y + 7$

3. Examine whether the following pairs of linear equations have unique solution, no solution or infinitely many solutions or Not? In case there is a unique solution, find it by using cross-multiplication method :

(i)  $2x - 3y - 4 = 0$       (ii)  $4x - 3y = 5$       (iii)  $2x + y - 5 = 0$   
 $4x - 6y + 5 = 0$        $3x - 5y = 1$        $3x + 2y - 8 = 0$

(iv)  $x - 2y + 3 = 0$       (v)  $x - 3y = 7$       (vi)  $ax + by = c^2$   
 $3x - 6y + 9 = 0$        $x - y = 5$        $\frac{x+a}{b} - \frac{y+b}{a} = 0$

(vii)  $7x - 5y = 11$       (viii)  $2x + 3y - 8 = 0$       (ix)  $\frac{x}{a} + \frac{y}{b} = 2$   
 $3x + 2y = 13$        $3x - 4y + 5 = 0$        $ax - by = a^2 - b^2$

(x)  $ax + by = a - b$       (xi)  $\frac{x}{a} + \frac{y}{b} = a + b$       (xii)  $\frac{x}{a} - \frac{y}{b} = 0$   
 $bx - ay = a + b$        $\frac{x}{a^2} + \frac{y}{b^2} = 2$        $ax + by = a^2 + b^2$

4. Solve the following pair of equations by reducing them to a pair of linear equations (by any algebraic method):

(i)  $\frac{6}{x} + \frac{8}{y} = 5$       (ii)  $\frac{6}{x} + \frac{10}{y} = 7$       (iii)  $\frac{3}{y} - \frac{1}{x} = 1$

$\frac{8}{x} + \frac{12}{y} = 7$        $\frac{2}{x} + \frac{3}{y} = \frac{13}{6}$        $\frac{2}{5x} + \frac{5}{2y} = 7$

(iv)  $\frac{a}{x} + \frac{b}{y} = p$       (v)  $\frac{2}{\sqrt{x}} + \frac{3}{\sqrt{y}} = 2$

$\frac{b}{x} + \frac{a}{y} = q$ , where  $\frac{a}{b} \neq \frac{p}{q} \neq \frac{q}{p}$        $\frac{4}{\sqrt{x}} - \frac{9}{\sqrt{y}} = -1$

4. (i)  $x = 2, y = 4$                       (ii)  $x = 3, y = 2$                       (iii)  $x = \frac{1}{5}, y = \frac{1}{2}$

(iv)  $x = \frac{a^2 - b^2}{ap - bq}, y = \frac{a^2 - b^2}{aq - bp}$                       (v)  $x = 4, y = 9$

(vi)  $x = 1, y = 1$                       (vii)  $x = 4, y = 5$                       (viii)  $x = 1, y = 2$

(ix)  $x = 3, y = 2$                       (x)  $x = 1, y = 1$                       (xi)  $x = 12, y = 4$

(xii)  $x = 4, y = 2$

**4.5 Problems Involving Pair of Linear Equations in Two Variables**

Pair of linear equations in two variables can be used to solve a wide variety of problems. We discuss a few problems below.

**Example 15. Two numbers are in the ratio 2:3. If 6 is added to each number, the sums are in the ratio 3:4. Find the numbers.**

**Solution :** Let  $x$  and  $y$  be the two numbers.  
Now, from the given conditions of the problem, we get

$$\frac{x}{y} = \frac{2}{3}$$

$$\Rightarrow 3x = 2y \dots\dots\dots(1)$$

and  $\frac{x+6}{y+6} = \frac{3}{4}$

$$\Rightarrow 4x - 3y + 6 = 0 \dots\dots\dots(2)$$

Solving for  $x$  and  $y$ , we get  
 $x = 12$   
and  $y = 18$   
Hence, the required numbers are 12 and 18.

**Example 16. A number of two digits is equal to four times the sum of its digits. If 18 is added to the number, the digits are reversed; find the number.**

**Solution :** Let  $x$  be the digit in the unit's place and  $y$  the digit in the ten's place of the number.  
Then the number =  $10y + x \dots\dots\dots(1)$   
The number formed by interchanging the places of the two digits =  $10x + y$ .  
The sum of the digits =  $x + y$   
From the given conditions of the problem, we get

Solving (1) and (2) for  $x$  and  $y$ , we get  $x = 60$  and  $y = 28$

Hence, the present age of the father is 60 years and that of the son is 28 years.

**Example 19.** A man purchased a cycle and a watch for ₹ 2,500. He then sold the cycle at a profit of 20 % and the watch at a loss of 10%. He then made a profit of 8% on the whole. What was the cost price of the cycle ?

**Solution :** Let ₹  $x$  and ₹  $y$  be the prices of the cycle and the watch respectively.

$$\text{Then, } x + y = 2500 \dots(1)$$

$$\text{Total selling price} = (x + 20\% \text{ of } x) + (y - 10\% \text{ of } y)$$

$$= \frac{6x}{5} + \frac{9y}{10} = \frac{12x + 9y}{10}$$

$$\text{Actual gain} = \text{selling price} - \text{cost price}$$

$$= \frac{12x + 9y}{10} - (x + y) = \frac{2x - y}{10}$$

But gain in the whole transaction is 8% of the total cost price.

$$\therefore \frac{2x - y}{10} = 8\% \text{ of } (x + y) = \frac{8}{100}(x + y)$$

$$\Rightarrow 2x = 3y \quad \dots(2)$$

Solving (1) and (2) for  $x$  and  $y$ , we have

$$x = 1500$$

$$\text{and } y = 1000$$

Hence, the cost price of the cycle is Rs. 1500.

**Example 20.** Chaoba and Tomba are at two stations on a highway 27 km. apart. They meet each other in 9 hours, if they walk in the same direction, but in 3 hours, if they walk in opposite directions. Find their rates of walking if Chaoba can walk faster than Tomba.

**Solution :** Let  $x$  km/hr and  $y$  km/hr be the walking speeds of Chaoba and Tomba respectively so that  $x > y$ . When they walk in opposite directions their relative speed =  $(x + y)$  km/hr and the relative speed is  $(x - y)$  km/hr when they walk in the same direction. But, distance travelled with uniform speed = speed  $\times$  time.

$$\therefore 27 = 3(x + y)$$

$$\Rightarrow x + y = 9 \quad \dots(1)$$

12. Two stations A and B on a highway are 90km. apart. A car starts from A and another car starts from B at the same time. If they travel in the same direction they meet in 9 hours, but if they travel towards each other they meet in 1 hour after start. Find the speeds of the two cars, the car from A moving faster.
13. 90% and 95% pure mustard oils are mixed to obtain 20 litres of a 92% pure mustard oil. How many litres of each kind of mustard oil are needed?
14. A steamer goes 50 km. downstream and 45 km. upstream in 5 hours. In 5 hours 8 minutes it can go 50 km. upstream and 45 km. downstream. Find the speed of the stream and that of the steamer in still water.
15. In a rectangle, if the length is reduced by 5cm. and the breadth is increased by 2cm, the area is reduced by 40 sq. cm. If however, the length is increased by 2cm. and the breadth by 4cm, the area is increased by 92 sq. cm. Find the length and breadth of the rectangle.
16. The area of a rectangular garden is increased by 55 sq. m. if its length is reduced by 2 m. and the breadth is increased by 5 m. If the length is increased by 3m. and the breadth by 2 m. the area is increased by 70 sq. m. Find the area of the rectangular garden.
17. A rectangle is of the same area as another which is 6 m longer and 4 m narrower. It is also of the same area as a third rectangle which is 8 m longer and 5 m. narrower. Find the area of the rectangle.
18. An annual income of ₹ 1,200 is derived from two sums invested, one at 4% and the other at 6% per annum simple interest. If the rates of interest are changed to 5% and 7% per annum simple interest respectively, the annual income derived from the investments is ₹ 1450 ; find the sums invested.
19. A man invested ₹ 36,000, a part of it at 12% and the rest at 15% per annum simple interest. If he received a total annual interest of ₹ 4,890, how much did he invest at each rate?
20. In a classroom there are a number of benches. If 4 students sit on each bench, five benches are left vacant; and if 3 students sit on each bench, 4 students are left standing. Find the number of benches and students in the class room.
21. The sum of a two-digit number and the number obtained by reversing the digits is 110. If the difference of the digits of the number is 4, find the number. How many such numbers are there?
22. The ratio of incomes of two persons is 9:7 and the ratio of their expenditures is 4:3. If each of them saves ₹ 2000 per month, find their monthly incomes.
23. A two - digit number is obtained either by multiplying the sum of the digits by 8 and adding 1, or by multiplying the difference of the digits by 13 and adding 2. Find the number. How many such numbers are there?



©BOSEM Not to be republished

### 3.3 Factorisation of Cyclic Expressions

In many cases, cyclic expressions can be factorised by using the following steps:

- (i) Write the terms of the expression according to the ascending or descending powers of one of the letters involved in the expression.
- (ii) Take out the factor(s) common to each coefficient.
- (iii) Write the terms of the other factor according to the ascending or descending powers of any letters other than the previous.
- (iv) Repeat the process till the factorisation is complete.

Now, let us illustrate the process by considering some cyclic expressions.

#### 3.4 To Factorise $a^2(b+c)+b^2(c+a)+c^2(a+b)+2abc$

$$\begin{aligned}
 & a^2(b+c)+b^2(c+a)+c^2(a+b)+2abc \\
 &= a^2(b+c)+b^2c+ab^2+c^2a+bc^2+2abc \\
 &= a^2(b+c)+a(b^2+c^2+2bc)+(b^2c+bc^2) \text{ [arranging in descending powers of } a] \\
 &= a^2(b+c)+a(b+c)^2+bc(b+c) \\
 &= (b+c) [a^2+a(b+c)+bc] \\
 &= (b+c)[b(c+a)+a(c+a)] \text{ [arranging in descending powers of } b] \\
 &= (b+c)(c+a)(b+a) \\
 &= (a+b)(b+c)(c+a)
 \end{aligned}$$

#### 3.5 To Factorise $a^2(b-c)+b^2(c-a)+c^2(a-b)$

$$\begin{aligned}
 & a^2(b-c)+b^2(c-a)+c^2(a-b) \\
 &= a^2(b-c)+b^2c-ab^2+c^2a-bc^2 \\
 &= a^2(b-c)-a(b^2-c^2)+(b^2c-bc^2) \text{ [arranging in descending powers of } a] \\
 &= a^2(b-c)-a(b+c)(b-c)+bc(b-c) \\
 &= (b-c)[a^2-a(b+c)+bc] \\
 &= (b-c)[b(c-a)-a(c-a)] \text{ [arranging in descending powers of } b] \\
 &= (b-c)(c-a)(b-a) \\
 &= -(a-b)(b-c)(c-a)
 \end{aligned}$$

**3.8 To Factorise**  $a^2(b+c)+b^2(c+a)+c^2(a+b)+3abc$ .

$$\begin{aligned} & a^2(b+c)+b^2(c+a)+c^2(a+b)+3abc \\ &= a^2b+ca^2+b^2c+ab^2+c^2a+bc^2+3abc \\ &= (a^2b+ab^2+abc)+(abc+b^2c+bc^2)+(ca^2+abc+c^2a) \\ &= ab(a+b+c)+bc(a+b+c)+ca(a+b+c) \\ &= (a+b+c)(ab+bc+ca) \end{aligned}$$

**Note :**

$$\begin{aligned} & a^2(b+c)+b^2(c+a)+c^2(a+b) \\ &= bc(b+c)+ca(c+a)+ab(a+b).....(1) \\ &= a(b^2+c^2)+b(c^2+a^2)+c(a^2+b^2).....(2) \end{aligned}$$

Therefore,  $a^2(b+c)+b^2(c+a)+c^2(a+b)$  can be replaced by either (1) or (2).

**3.9 To Factorise**  $a^3+b^3+c^3-3abc$ .

$$\begin{aligned} & a^3+b^3+c^3-3abc \\ &= (a+b)^3-3ab(a+b)+c^3-3abc [\because a^3+b^3=(a+b)^3-3ab(a+b)] \\ &= (a+b)^3+c^3-\{3ab(a+b)+3abc\} \\ &= (a+b+c)\{(a+b)^2-(a+b)c+c^2\}-3ab(a+b+c) \\ & \qquad \qquad \qquad [\because x^3+y^3=(x+y)(x^2-xy+y^2)] \\ &= (a+b+c)(a^2+2ab+b^2-ca-bc+c^2-3ab) \\ &= (a+b+c)(a^2+b^2+c^2-ab-bc-ca).....(1) \\ &= \frac{1}{2}(a+b+c)(2a^2+2b^2+2c^2-2ab-2bc-2ca) \\ &= \frac{1}{2}(a+b+c)\{(a^2-2ab+b^2)+(b^2-2bc+c^2)+(c^2-2ca+a^2)\} \\ &= \frac{1}{2}(a+b+c)\{(a-b)^2+(b-c)^2+(c-a)^2\}.....(2) \end{aligned}$$

Hence,

$$\begin{aligned} & a^3+b^3+c^3-3abc \\ &= (a+b+c)(a^2+b^2+c^2-ab-bc-ca) \\ &= \frac{1}{2}(a+b+c)\{(a-b)^2+(b-c)^2+(c-a)^2\}. \end{aligned}$$

**3.10 To Factorise**  $2b^2c^2+2c^2a^2+2a^2b^2-a^4-b^4-c^4$ .

We have,  $(a+b+c)^2 = a^2+b^2+c^2+2ab+2bc+2ca$ .

$\therefore$  The given expression

3. Prove that  $(x - y)^3 + (y - z)^3 + (z - x)^3 - 3(x - y)(y - z)(z - x) = 0$
4. If  $a^3 + b^3 + c^3 = 3abc$ , prove that either  $a + b + c = 0$  or  $a = b = c$ .
5. If  $x + y + z = 9$ ,  $xy + yz + zx = 26$  and  $xyz = 24$ , find the value of  $x^2(y + z) + y^2(z + x) + z^2(x + y)$ .
6. If  $x + y - z = 2$ ,  $y + z - x = 4$  and  $z + x - y = 6$ , find the value of  $2y^2z^2 + 2z^2x^2 + 2x^2y^2 - x^4 - y^4 - z^4$ .
7. If  $x + y + z = 12$  and  $x^2 + y^2 + z^2 = 44$ , find the value of  $(x + y + z)^3 - x^3 - y^3 - z^3 + 3xyz$ .
8. Find the value of  $xy(x + y) + yz(y + z) + zx(z + x) + 3xyz$ , when  $x = a(b - c)$ ,  $y = b(c - a)$ ,  $z = c(a - b)$ .

## ANSWER

1.
  - (i)  $(x + y - z)(x^2 + y^2 + z^2 - xy + yz + zx)$
  - (ii)  $(a - b + 3)(a^2 + b^2 + 9 + ab + 3b - 3a)$
  - (iii)  $(2a + 3b + 4c)(4a^2 + 9b^2 + 16c^2 - 6ab - 12bc - 8ca)$
  - (iv)  $(x - y - 5z)(x^2 + y^2 + 25z^2 + xy - 5yz + 5zx)$
  - (v)  $(a^2 - a + 2)(a^4 + a^3 - a^2 + 2a + 4)$
  - (vi)  $(x^2 - x + 3)(x^4 + x^3 - 2x^2 + 3x + 9)$
2.
  - (i)  $-(x - y)(y - z)(z - x)$
  - (ii)  $(x + y)(y + z)(z + x)$
  - (iii)  $(a + b + c)(ab + bc + ca)$
  - (iv)  $(x + y)(y + z)(z + x)$
  - (v)  $(a + b + c)(ab + bc + ca)$
  - (vi)  $-(x - y)(y - z)(z - x)(x^2 + y^2 + z^2 + xy + yz + zx)$
  - (vii)  $-(x - y)(y - z)(z - x)(x^2 + y^2 + z^2 + xy + yz + zx)$
  - (viii)  $-(a + b)(b + c)(c + a)(a - b)(b - c)(c - a)$
  - (ix)  $-(x - y)(y - z)(z - x)(xy + yz + zx)$
  - (x)  $3(x + z)(y + 2z - x)(z - y)$

©BOSEM Not to be republished

A succession of numbers formed according to a specific rule is called a sequence.

The numbers in a sequence are called **terms** of the sequence. The terms are labelled as first term, second term, third term, fourth term etc. which are usually denoted by  $a_1, a_2, a_3, a_4, \dots$  respectively.

For example, in (A), we have

$$a_1 = 10000, a_2 = 20000, a_3 = 40000, a_4 = 80000, \text{ etc.}$$

The number of terms of a sequence may be finite or infinite and accordingly the sequence is called finite or infinite. Usually in an infinite sequence, only the first few terms are given followed by dots. But the terms of a sequence follow a specific rule and this enables us to express the  $n^{\text{th}}$  term (or the general term)  $a_n$  in terms of  $n$  where  $n \in \mathbb{N}$ . For example, for sequence (A), we have

$$\begin{aligned} a_1 &= 10000 = 10000 \times 1 = 10000 \times 2^0 = 10000 \times 2^{1-1} \\ a_2 &= 20000 = 10000 \times 2 = 10000 \times 2^1 = 10000 \times 2^{2-1} \\ a_3 &= 40000 = 10000 \times 4 = 10000 \times 2^2 = 10000 \times 2^{3-1} \\ a_4 &= 80000 = 10000 \times 8 = 10000 \times 2^3 = 10000 \times 2^{4-1} \\ \dots & \dots \dots \dots \dots \dots \dots \dots \dots \\ \dots & \dots \dots \dots \dots \dots \dots \dots \dots \end{aligned}$$

Following the pattern, we see that

$$a_n = 10000 \times 2^{n-1}$$

Once we know  $a_n$  (in terms of  $n$ ) of a sequence, we can find any term of the sequence.

Thus, we have

$$\begin{aligned} a_6 &= 10000 \times 2^{6-1} = 320000, \\ a_{11} &= 10000 \times 2^{11-1} = 10000 \times 2^{10} \text{ etc.} \end{aligned}$$

Also, we have the answer to one of the three questions.

Now, the maturity amount after 25 years

$$= a_{25} = 10000 \times 2^{25-1} = ₹ 10000 \times 2^{24}$$

[Can you find the  $n^{\text{th}}$  term for the sequences (B) & (C)?]

### 6.3 The $n^{\text{th}}$ term of an AP

Let  $a$  be the first term and  $d$  be the common difference of an AP. Then, the AP is  $a, a+d, a+2d, a+3d, \dots$

$$\begin{aligned} \text{Here, } a_1 &= a = a + (1-1)d \\ a_2 &= a + d = a + (2-1)d \\ a_3 &= a + 2d = a + (3-1)d \\ a_4 &= a + 3d = a + (4-1)d \\ \dots & \dots \dots \dots \dots \\ \dots & \dots \dots \dots \dots \end{aligned}$$

Looking at the pattern, we can say that

$$a_n = a + (n-1)d.$$

Thus, for an AP whose first term is  $a$  and the common difference is  $d$ ,

**the  $n^{\text{th}}$  term (or the general term)  $a_n = a+(n-1)d$**

**Example 1.** Which of the following sequences are AP. In case of an AP, find the first term  $a$  and the common difference  $d$  and write the next three terms.

- (i)  $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$       (ii)  $4, 2, 0, -2, -4, \dots$
- (iii)  $\frac{1}{10}, \frac{1}{5}, \frac{3}{10}, \frac{2}{5}, \frac{1}{2}, \dots$       (iv)  $1, 2, 3, 5, 7, \dots$

**Solution :** (i) We have,  $a_1 = 1, a_2 = \frac{1}{2}, a_3 = \frac{1}{3}, a_4 = \frac{1}{4}, a_5 = \frac{1}{5}, \dots$

$$\text{Now, } a_2 - a_1 = \frac{1}{2} - 1 = -\frac{1}{2}$$

$$a_3 - a_2 = \frac{1}{3} - \frac{1}{2} = \frac{2-3}{6} = -\frac{1}{6}$$

As,  $a_3 - a_2 \neq a_2 - a_1$ , the given sequence is not an AP.

(ii) We have,  $a_1 = 4, a_2 = 2, a_3 = 0, a_4 = -2, a_5 = -4, \dots$

$$\text{Now, } a_2 - a_1 = 2 - 4 = -2$$

$$a_3 - a_2 = 0 - 2 = -2$$

(iv) We have ,  $a_1 = 1, a_2 = 2, a_3 = 3, a_4 = 5, a_5 = 7, \dots$

$$\text{Now, } a_2 - a_1 = 2 - 1 = 1$$

$$a_3 - a_2 = 3 - 2 = 1$$

$$a_4 - a_3 = 5 - 3 = 2$$

As,  $a_4 - a_3 \neq a_3 - a_2$ , the given sequence is not an AP.

**Example 2.** Write the  $n^{\text{th}}$  term and also the first four terms of the AP whose first term  $a$  and the common difference 'd' are given as follows :

(i)  $a = 5, d = 3$                       (ii)  $a = 0, d = -2$

(iii)  $a = -10, d = 0.5$               (iv)  $a = 6, d = \frac{1}{5}$

**Solution :** (i) We have,  $a = 5, d = 3$

$$\therefore a_n = a + (n-1)d$$

$$= 5 + (n-1) \times 3$$

$$= 3n + 2$$

So,  $a_1 = 3 \times 1 + 2 = 5, a_2 = 3 \times 2 + 2 = 8, a_3 = 3 \times 3 + 2 = 11$  and

$$a_4 = 3 \times 4 + 2 = 14.$$

Thus, the first four terms are 5, 8, 11 and 14.

[The first four terms can also be obtained as

$$a_1 = a = 5$$

$$a_2 = a + d = 5 + 3 = 8$$

$$a_3 = a + 2d = 5 + 2 \times 3 = 11$$

$$\text{and } a_4 = a + 3d = 5 + 3 \times 3 = 14]$$

(ii) We have,

$$a = 0, d = -2$$

$$\therefore a_n = a + (n-1)d$$

$$= 0 + (n-1) \times (-2)$$

$$= 2 - 2n$$

So,  $a_1 = 2 - 2 \times 1 = 0, a_2 = 2 - 2 \times 2 = -2,$

$$a_3 = 2 - 2 \times 3 = -4, a_4 = 2 - 2 \times 4 = -6.$$

Thus, the first four terms are 0, -2, -4 and -6.



**Example 4.** Is 72 a term of the AP : 7, 10, 13, 16,.....?

**Solution :** The given AP is 7, 10, 13, 16, ..... ?  
Here,  $a = 7$  and  $d = 10 - 7 = 3$   
If possible, let 72 be the  $n^{\text{th}}$  term of the AP.  
Then,  $72 = a + (n-1)d$

$$\Rightarrow 72 = 7 + (n-1) \times 3$$

$$\Rightarrow 72 = 3n + 4$$

$$\Rightarrow 3n = 72 - 4 = 68$$

$$\Rightarrow n = \frac{68}{3}, \text{ which is not a natural number.}$$

Therefore, 72 is not a term of the given AP.

**Note:** For 72 to be a term of the AP,  $n$  must be a natural number because there

is no term such as  $\left(\frac{68}{3}\right)^{\text{th}}$  term

**Example 5.** The  $n^{\text{th}}$  term of a sequence is given by  $a_n = 4n + 5$ . Show that the sequence is an AP.

**Solution :** We have,  $a_n = 4n + 5$

$$\therefore a_{n+1} = 4(n+1) + 5 = 4n + 9$$

$$\text{Now, } a_{n+1} - a_n = (4n + 9) - (4n + 5) = 4$$

Thus,  $a_{n+1} - a_n = \text{constant for all } n \in \mathbb{N}$ .

$\therefore$  the given sequence is an AP with common difference 4.

**Example 6.** The 6<sup>th</sup> and the 17<sup>th</sup> terms of an AP are respectively 19 and 41. Find the first term and the common difference. Also, find the 25<sup>th</sup> term.

**Solution :** Let  $a$  be the first term and  $d$ , the common difference.

$$\text{Here, } a_6 = 19$$

$$\Rightarrow a + 5d = 19 \quad \dots(1)$$

$$\text{and } a_{17} = 41$$

$$\Rightarrow a + 16d = 41 \quad \dots(2)$$

Solving (1) and (2), we get

$$a = 9 \text{ and } d = 2$$

$\therefore$  first term is 9 and common difference is 2.

$$\begin{aligned}
 \therefore n^{\text{th}} \text{ term} = a_n &= a + (n-1)d \\
 &= (p+q-1) + (n-1) \times (-1) \\
 &= p+q-1-n+1 \\
 &= p+q-n.
 \end{aligned}$$

**Example 9.** Ramesh got a government job and started work with a monthly salary of ₹ 8000 and enjoying an annual increment of ₹ 400. Find his annual income in the 10<sup>th</sup> year of service.

**Solution :** We have, the monthly salary (in ₹) of Ramesh in the first year, second year, third year, fourth year etc. are 8000, 8000 + 400, 8000 + 400 + 400, 8000 + 400 + 400 + 400 etc. i.e., 8000, 8400, 8800, 9200 etc. which form an AP.

Here,  $a = 8000$  and  $d = 400$

Now, the monthly salary (in ₹) in the tenth year

$$\begin{aligned}
 a_{10} &= a + (10-1)d \\
 &= 8000 + 9 \times 400 \\
 &= 8000 + 3600 \\
 &= 11600
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{The annual income in the tenth year of service} \\
 &= ₹ 11600 \times 12 \\
 &= ₹ 139200.
 \end{aligned}$$

### EXERCISE 6.1

1. Which of the following sequences are AP? In case of AP, find the first term  $a$  and the common difference  $d$  and write the next three terms.

(i)  $1^2, 2^2, 3^2, 4^2, 5^2, \dots$                       (ii)  $8, 5, 2, -1, -4, \dots$

(iii)  $2, \frac{5}{2}, 3, \frac{7}{2}, 4, \dots$                       (iv)  $0.6, 1.7, 2.8, 3.9, 5, \dots$

(v)  $3, 5, 7, 8, 10, \dots$                       (vi)  $-6, -1, 4, 9, 14, \dots$

2. Write the  $n^{\text{th}}$  term and also the first four terms of the AP whose first term  $a$  and the common difference  $d$  are given as follows :

(i)  $a = 2, d = 2$                                       (ii)  $a = 6, d = -4$

(iii)  $a = -10, d = -3$                               (iv)  $a = \frac{3}{2}, d = -\frac{1}{2}$

## ANSWER

1. (i) The sequence is not an AP.  
(ii) The sequence is an AP ;  $a = 8, d = -3; -7, -10, -13$   
(iii) The sequence is an AP ;  $a = 2, d = \frac{1}{2}; \frac{9}{2}, 5, \frac{11}{2}$   
(iv) The sequence is an AP;  $a = 0.6, d = 1.1; 6.1, 7.2, 8.3$   
(v) The sequence is not an AP  
(vi) The sequence is an AP ;  $a = -6, d = 5; 19, 24, 29$
2. (i)  $a_n = 2n$ ;  $a_1 = 2, a_2 = 4, a_3 = 6, a_4 = 8$   
(ii)  $a_n = 10 - 4n$ ;  $a_1 = 6, a_2 = 2, a_3 = -2, a_4 = -6$   
(iii)  $a_n = -7 - 3n$ ;  $a_1 = -10, a_2 = -13, a_3 = -16, a_4 = -19$   
(iv)  $a_n = \frac{4-n}{2}$ ;  $a_1 = \frac{3}{2}, a_2 = 1, a_3 = \frac{1}{2}, a_4 = 0$   
(v)  $a_n = 4 + \frac{n}{2}$ ;  $a_1 = 4.5, a_2 = 5, a_3 = 5.5, a_4 = 6$   
(vi)  $a_n = \frac{n-6}{3}$ ;  $a_1 = -\frac{5}{3}, a_2 = -\frac{4}{3}, a_3 = -1, a_4 = -\frac{2}{3}$
3. (i) 32            (ii) -38            (iii)  $\frac{21}{2}$             (iv)  $35\sqrt{2}$
4. (i) No            (ii) No
5. (i) 25<sup>th</sup> term    (ii) 22<sup>nd</sup> term
6. (i) 13            (ii) 31
7. The sequence is an AP; first term = -3, common difference = -5
9. 67
10. First term = 5, common difference =  $\frac{1}{2}$
11. 123
12.  $-\frac{8}{5}$
13. 30
16. ₹ 7500
17. 15

And, if the AP is finite with only  $n$  terms,

then  $a_n = l = \text{last term}$

By then,  $S_n = \frac{n}{2}[a + l]$

**Example 10.** Find the sum of the first 25 terms of the AP : 20, 17, 14, 11, 8, .....

**Solution :** Here,  $a = 20$  and  $d = 17 - 20 = -3$

And, we have  $S_n = \frac{n}{2}[2a + (n-1)d]$

Putting  $n = 25$ , we have

$$\begin{aligned} \text{the required sum} = S_{25} &= \frac{25}{2}[2 \times 20 + (25-1) \times (-3)] \\ &= \frac{25}{2}(40 - 72) \\ &= \frac{25}{2}(-32) \\ &= 25 \times (-16) \\ &= -400 \end{aligned}$$

**Example 11.** Find the sum of all the terms of the finite AP : 2, 5, 8, ....., 179, 182.

**Solution :** Here,  $a = 2$  and  $d = 5 - 2 = 3$   
Let there be  $n$  terms in the AP.

Then,  $a_n = 182$

$$\Rightarrow a + (n-1)d = 182$$

$$\Rightarrow 2 + (n-1) \times 3 = 182$$

$$\Rightarrow (n-1) \times 3 = 180$$

$$\Rightarrow (n-1) = \frac{180}{3} = 60$$

$$\Rightarrow n = 61$$

$$\begin{aligned} \therefore \text{the required sum} = S_{61} &= \frac{61}{2}[2 + 182] \quad \left[ \begin{array}{l} \text{using the formula,} \\ S_n = \frac{n}{2}[a + l] \text{ for} \\ \text{a finite AP} \end{array} \right] \\ &= \frac{61}{2} \times 184 \\ &= 61 \times 92 = 5612 \end{aligned}$$

**Example 14. Find the sum of (i) the first  $n$  odd natural numbers**

**(ii) the first  $n$  even natural numbers**

**Solution :** (i) The sequence of the odd natural numbers is : 1, 3, 5, 7, 9, 11 ..., which is an AP.

Here,  $a = 1$  and  $d = 3 - 1 = 2$

$$\begin{aligned}\therefore \text{ the required sum} = S_n &= \frac{n}{2}[2a + (n-1)d] \\ &= \frac{n}{2}[2 \times 1 + (n-1) \times 2] \\ &= \frac{n}{2} \times 2n = n^2\end{aligned}$$

(ii) The sequence of the even natural numbers is : 2, 4, 6, 8, 10, ..., which is an AP.

Here,  $a = 2$  and  $d = 2$ .

$$\begin{aligned}\therefore \text{ the required sum} = S_n &= \frac{n}{2}[2a + (n-1)d] \\ &= \frac{n}{2}[2 \times 2 + (n-1) \times 2] \\ &= \frac{n}{2}[2n + 2] \\ &= \frac{n}{2} \times 2(n+1) \\ &= n(n+1).\end{aligned}$$

**Example 15. Show that the sequence whose  $n^{\text{th}}$  term is given by  $a_n = 5 - 3n$ , is an AP. And, find the sum of the first 15 terms of the AP.**

**Solution :** Here,  $a_n = 5 - 3n$

$$\begin{aligned}\therefore a_{n+1} &= 5 - 3(n+1) \\ &= 5 - 3n - 3 = 2 - 3n\end{aligned}$$

Now,  $a_{n+1} - a_n = (2 - 3n) - (5 - 3n)$

$$= -3 \text{ which is constant for all } n \in \mathbb{N}$$

$\therefore$  the sequence is an AP.

Then,  $a = a_1 = 5 - 3 \times 1 = 2$

and  $d = -3$

$$\Rightarrow \frac{a}{p} = \frac{1}{2} [2A + (p-1)D] \quad \dots(i)$$

also,  $b = S_q$

$$\Rightarrow b = \frac{q}{2} [2A + (q-1)D]$$

$$\Rightarrow \frac{b}{q} = \frac{1}{2} [2A + (q-1)D] \quad \dots(ii)$$

and  $c = S_r$

$$\Rightarrow c = \frac{r}{2} [2A + (r-1)D]$$

$$\Rightarrow \frac{c}{r} = \frac{1}{2} [2A + (r-1)D] \quad \dots(iii)$$

Multiplying (i), (ii) and (iii) by  $(q-r)$ ,  $(r-p)$  and  $(p-q)$  respectively and adding, we get

$$\begin{aligned} & \frac{a}{p}(q-r) + \frac{b}{q}(r-p) + \frac{c}{r}(p-q) \\ &= \frac{1}{2} [2A + (p-1)D](q-r) + \frac{1}{2} [2A + (q-1)D](r-p) + \frac{1}{2} [2A + (r-1)D](p-q) \\ &= \frac{1}{2} [2A(q-r+r-p+p-q) + D\{(p-1)(q-r) + (q-1)(r-p) + (r-1)(p-q)\}] \\ &= \frac{1}{2} [2A \times 0 + D \times 0] \\ &= \frac{1}{2} \times 0 \\ &= 0 \end{aligned}$$

Hence shown.

**Example 18.** Anita put ₹ 50 into her daughter's money box on her first birthday and increased the amount by ₹ 10 every next birthday. Find the total amount in the money box by the time the daughter turns 25.

**Solution :** We have, the amount of money (in ₹) put in the money box on the first birthday, second birthday, third birthday etc. are respectively 50, 60, 70 etc. which form an AP.

Here,  $a = 50$  and  $d = 10$

Now, the required amount of money (in ₹)

6. Find the sum of all odd numbers between 100 and 200.
7. Find the sum of the first 25 multiples of 6.
8. Find the sum of all the multiples of 7 between 100 and 300.
9. If the 12<sup>th</sup> term of an AP is  $-13$  and the sum of the first four terms is 24, find the sum of the first 15 terms.
10. Find the sum of the first 22 terms of an AP whose 8<sup>th</sup> and 16<sup>th</sup> terms are respectively 37 and 85.
11. The sum of the first 7 terms of an AP is 10 and that of the next 7 terms is 17. Find the first term and the common difference of the AP.
12. Find the sum of the first 50 terms of an AP whose  $n^{\text{th}}$  term is  $3 - 2n$ .
13. Find the first term and the common difference of an AP if the sum of the first  $n$  terms is  $\frac{n(5n + 7)}{12}$ .
14. In an AP, if the sum of the first  $m$  terms is equal to  $n$  and that of the first  $n$  terms is equal to  $m$ , then prove that the sum of the first  $(m+n)$  terms is  $-(m+n)$ .
15. The sums of the first  $n$ ,  $2n$ ,  $3n$  terms of an AP are  $S_1$ ,  $S_2$ ,  $S_3$  respectively. Show that  $S_3 = 3(S_2 - S_1)$ .
16. A man repays a debt of ₹ 4860 by paying ₹ 30 in the first month and then increases the amount by ₹ 15 every month. How long will it take him to clear the debt (Assume that no interest is charged) ?
17. A manufacturer of radio sets produced 600 units in the third year and 700 units in the seventh year. Assuming that the production increases uniformly by a fixed number every year, find the production in the first year and also the total production in 8 years.

### ANSWER

1. (i) 180      (ii) 366      (iii) 100      (iv) 44.8
2. (i) 735      (ii) 5612      (iii) 408      (iv)  $-50$
3. 8

4. 6 or 11 *Explanation:* Since  $d$  is negative, the terms go on diminishing and the 9<sup>th</sup> term becomes zero after which the terms become negative. Thus, the positive and the negative terms from the 7<sup>th</sup> term to the 11<sup>th</sup> term cancel out each other.


### 7.1 Introduction.

In earlier classes, you have studied about triangles and many of their properties. In Class IX, you have studied about the congruence of triangles. Recall that two plane figures are congruent if they have the same shape and the same size. In our day-to-day life, we often come across figures of the same shape but of different sizes, for example bangle of different sizes, different size photographs from the same negative etc. To construct a bridge, engineers usually draw a picture (called blue print) of a model, much smaller in size than the actual bridge. The bridge and the model are of the same shape but of different sizes. Two figures having the same shape but not necessarily the same size are called similar figures. In this chapter, we shall first discuss about the similar figures, particularly, the similar triangles and then apply the knowledge of the similarity of triangles to obtain a simple proof of the Pythagoras Theorem which you have studied in the earlier class.

### 7.2. Similar Figures

In class IX, you have learnt that two line segments of equal length are congruent and two angles of equal measure are congruent. You have also learnt that two equilateral triangles with the same side lengths are congruent and two circles with the same radii are congruent.

The two circles (**Fig 7.1**) are of the same shape but of different radii. So, they are similar but not congruent.



By the same reasoning, the two equilateral triangles in figure 7.2 are similar but not congruent.






Fig. 7.2

From these examples, it is clear that all congruent figures are always similar but similar figures need not be congruent.

Our experience of solid objects enables us to recognise quite easily when two articles of different sizes have, as we say, ‘the same shape’. We shall now investigate what is involved in the question of shape. Draw a quadrilateral ABCD (of no special shape) on paper. Suppose we wish to make an enlarged version of the quadrilateral say , exactly two times in size. Let us take a suitable centre say O (Fig. 7.3) and scale factor (side ratio) 2. The enlargement is done by pushing out radially from the centre O: e.g. A' is on the line OA such that  $OA' = 2 \times OA$ .




Fig 7.3

In a similar way, take the points B', C', D'. Then, A'B'C'D' is the enlarged copy of ABCD. Compare A'B'C'D' with ABCD. You will observe that  $A'B' = 2AB$ ,  $B'C' = 2BC$ ,  $C'D' = 2CD$ ,  $D'A' = 2DA$  and  $\angle A = \angle A'$ ,  $\angle B = \angle B'$ ,  $\angle C = \angle C'$ ,  $\angle D = \angle D'$  i.e. not only are the sides in the larger figure exactly two times the


original sides but also all the corresponding angles are equal in both the figures. This is the essence of the similarity of two quadrilaterals and in general of two polygons. Thus:

**Two polygons having the same number of sides are similar, if (i) their corresponding angles are equal and (ii) their corresponding sides are in the same ratio.**

Now, to understand the similarity of two polygons clearly, let us perform the following activity:

**Activity :**

Draw a quadrilateral ABCD on a paper and take two photocopies of this paper, one enlarged and other reduced in size. Name the enlarged quadrilateral as  $A'B'C'D'$  and the reduced one as  $A''B''C''D''$  (Fig. 7.4)



**Fig. 7.4**

Observe that ABCD,  $A'B'C'D'$  and  $A''B''C''D''$  have the same shape but are of different sizes i.e., they are similar. Compare ABCD with  $A'B'C'D'$ . Note that the vertex  $A'$  corresponds to vertex A, vertex  $B'$  corresponds to vertex B, vertex  $C'$  corresponds to vertex C and vertex  $D'$  corresponds to vertex D. Measuring the angles and the sides of the two quadrilaterals, you will get

$$(i) \quad \angle A = \angle A', \angle B = \angle B', \angle C = \angle C', \angle D = \angle D' \text{ and}$$

$$(ii) \quad \frac{AB}{A'B'} = \frac{BC}{B'C'} = \frac{CD}{C'D'} = \frac{DA}{D'A'}$$


Again, comparing ABCD with  $A''B''C''D''$ . You will get

$$(i) \quad \angle A = \angle A'', \angle B = \angle B'', \angle C = \angle C'', \angle D = \angle D'' \text{ and}$$


$$(ii) \quad \frac{AB}{A''B''} = \frac{BC}{B''C''} = \frac{CD}{C''D''} = \frac{DA}{D''A''}$$

This result emphasises the requirements for two polygons having the same number of sides to be similar i.e., (i) the corresponding angles should be equal and (ii) their corresponding sides should be in the same ratio. Thus, to examine the similarity of two polygons, we have to examine three things (1) whether the two polygons have the same


number of sides or not, (2) whether the corresponding angles are equal or not and (3) whether the corresponding sides are in the same ratio or not. The polygons in Fig 7.5 are not similar because they do not have the same number of sides.



The quadrilaterals in Fig 7.6 are not similar as the corresponding angles are not equal.



The rectangles in Fig 7.7 are not similar as the corresponding sides are not in the same ratio.



### EXERCISE 7.1

1. Fill in the blanks using the correct word given in brackets :

- (i) All circles are ... (congruent, similar).
- (ii) All circles of same radii are ... (congruent, not congruent).
- (iii) All squares are ... (similar, congruent).
- (iv) All ... triangles are similar (isosceles, equilateral).
- (v) The reduced and enlarged photographs of an object made from the same negative are ... (similar, congruent).

**2. State true or false :**

- (i) All similar figures are congruent.
- (ii) All congruent figures are similar.
- (iii) All triangles are similar.
- (iv) All equilateral triangles are similar.
- (v) All rectangles are similar.
- (vi) All squares are not similar.
- (vii) Two photographs of the same size of the same person, one at the age of 10 years and the other at the age of 60 years are similar.

**ANSWER**

- |    |             |                |               |                  |
|----|-------------|----------------|---------------|------------------|
| 1. | (i) similar | (ii) congruent | (iii) similar | (iv) equilateral |
|    | (v) similar |                |               |                  |
| 2. | (i) False   | (ii) True      | (iii) False   | (iv) True        |
|    | (v) False   | (vi) False     | (vii) False.  |                  |

**7.3 Similarity of triangles**

A triangle is a polygon having three sides. So, according to the definition of similarity of polygons, two triangles will be similar if


- (i) Their corresponding sides are proportional and
- (ii) Their corresponding angles are equal.

A famous Greek mathematician Thales (600 B.C) stated that 'the ratio of any two corresponding sides in equiangular triangles is always the same irrespective of their sizes'. To get this result, it is believed that he used a theorem called the Basic Proportionality Theorem or Thales Theorem. In view of this result, the two conditions mentioned in the definition of similar triangles are found to be dependent on each other. It means, if one condition holds good, the other holds good automatically. In other words, if any of the two conditions are fulfilled i.e, either the corresponding angles of two triangles are equal or their corresponding sides are proportional, then the triangles are similar.

Let us perform the following activity.

**Activity :**

Draw an angle  $\angle XAY$  and mark points  $P_1, P_2, P_3, D, P_4$  and  $B$  on its arm  $AX$  such that  $AP_1 = P_1P_2 = P_2P_3 = P_3D = DP_4 = P_4B = 1$  unit (Fig. 7.8). Through  $B$ , draw a line intersecting arm  $AY$  at a point say,  $C$ . Through  $D$ , draw a line parallel to  $BC$  to intersect  $AC$  at  $E$ .



Now,  $AD = AP_1 + P_1P_2 + P_2P_3 + P_3D = 4$  units

and  $DB = DP_4 + P_4B = 2$  units

$$\therefore \frac{AD}{DB} = \frac{4}{2} = 2$$

Measure  $AE$  and  $EC$ . You will find that  $AE = 2EC$  i.e.  $\frac{AE}{EC} = 2$ . Here, we have observed that

in  $\triangle ABC$ ; if  $DE \parallel BC$ , then we have  $\frac{AD}{DB} = \frac{AE}{EC}$

We state this result as a theorem known as **Basic Proportionality Theorem** or **Thale's Theorem** as given below.

**Theorem 7.1** If a line is drawn parallel to one side of a triangle intersecting the other two sides, then it divides the other two sides in the same ratio.

**Given :**  $\triangle ABC$  and a line  $DE \parallel BC$  intersecting  $AB$  at  $D$  and  $AC$  at  $E$  (Fig 7.9).

**To prove :**  $\frac{AD}{DB} = \frac{AE}{EC}$

**Construction :** Join  $BE, CD$  and draw  $CF \perp AB$ ,  $BG \perp AC$  (Fig 7.9)

**Proof :** In fig 7.9, area of  $\triangle DBC =$  area of  $\triangle EBC$  ... (1) [being  $\triangle^s$  standing on the same base and between the same parallels]

$$\therefore \text{area of } \triangle ABC - \text{area of } \triangle DBC$$

$$= \text{area of } \triangle ABC - \text{area of } \triangle EBC$$

ie., area of  $\triangle ADC$  = area of  $\triangle AEB$  .....(2)

From (1) & (2), we have

$$\frac{\text{area of } \triangle ADC}{\text{area of } \triangle DBC} = \frac{\text{area of } \triangle AEB}{\text{area of } \triangle EBC}$$

$$\frac{\frac{1}{2} AD \cdot CF}{\frac{1}{2} DB \cdot CF} = \frac{\frac{1}{2} AE \cdot BG}{\frac{1}{2} EC \cdot BG}$$

$$\therefore \frac{AD}{DB} = \frac{AE}{EC}$$




Fig 7.9

Is the converse of this theorem true? That is, if a line divides any two sides of a triangle in the same ratio, is the line parallel to the third side? To examine this, let us perform the following activity.

Activity: Draw an angle  $\angle XAY$  and mark points  $B_1, B_2, B_3, B$  on AX such that  $AB_1 = B_1B_2 = B_2B_3 = B_3B$  (Fig. 7.10). Also mark points  $C_1, C_2, C_3$  on AY such that  $AC_1 = C_1C_2 = C_2C_3 = C_3C$ . Join BC and  $B_3C_3$ .




Fig . 7.10

Now,  $AB_3 = AB_1 + B_1B_2 + B_2B_3 = 3B_3B$

$$\text{i.e., } \frac{AB_3}{B_3B} = \frac{3}{1}$$

Again,  $AC_3 = AC_1 + C_1C_2 + C_2C_3 = 3C_3C$

$$\text{i.e., } \frac{AC_3}{C_3C} = \frac{3}{1}$$

$$\therefore \frac{AB_3}{B_3B} = \frac{AC_3}{C_3C}$$

Compare the angles  $\angle B_3C_3$  and  $\angle ABC$ . You will find that they are equal thereby showing  $B_3C_3 \parallel BC$ . You have observed that if a line divides two sides of a triangle in the same ratio, then the line is parallel to the third side. This fact is stated and proved as a theorem given below and it is the converse of the basic proportionality theorem.

**Theorem 7.2** If a line divides any two sides of a triangle in the same ratio, then the line is parallel to the third side.

**Given :** A line  $DE$  dividing the sides  $AB, AC$  of a  $\triangle ABC$  at  $D, E$  respectively such that

$$\frac{AD}{DB} = \frac{AE}{EC} \text{ (Fig. 7.11).}$$

**To prove :**  $DE \parallel BC$

**Construction :** Through  $D$ , draw  $DF \parallel BC$  meeting  $AC$  at  $F$  (Fig 7.11).

**Proof :** By Basic Proportionality Theorem,

$$\frac{AD}{DB} = \frac{AF}{FC}$$

But, it is given that

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\therefore \frac{AF}{FC} = \frac{AE}{EC}$$

$$\Rightarrow \frac{AF}{FC} + 1 = \frac{AE}{EC} + 1$$

$$\Rightarrow \frac{AF+FC}{FC} = \frac{AE+EC}{EC}$$

$$\Rightarrow \frac{AC}{FC} = \frac{AC}{EC}$$

$$\Rightarrow FC = EC$$

It shows that  $F$  coincides with  $E$ . But  $DF \parallel BC$  by construction.

Hence  $DF \parallel BC$ .




Fig. 7.11

**Example 1.** If a line segment  $DE$  is drawn parallel to the side  $BC$  of a  $\triangle ABC$

meeting  $AB$  at  $D$  and  $AC$  at  $E$ , then prove that  $\frac{AD}{AB} = \frac{AE}{AC}$  (Fig 7.12)

**Solution :**

**Given :**  $DE \parallel BC$  in Fig 7.12

**To Prove :**  $\frac{AD}{AB} = \frac{AE}{AC}$

**Proof :** It is given that  $DE \parallel BC$ .  
 $\therefore$  by Basic Proportionality Theorem,

$$\begin{aligned} \frac{AD}{DB} &= \frac{AE}{EC} \\ \Rightarrow \frac{DB}{AD} &= \frac{EC}{AE} \\ \Rightarrow 1 + \frac{DB}{AD} &= 1 + \frac{EC}{AE} \\ \Rightarrow \frac{AD+DB}{AD} &= \frac{AE+EC}{AE} \\ \Rightarrow \frac{AB}{AD} &= \frac{AC}{AE} \\ \therefore \frac{AD}{AB} &= \frac{AE}{AC} \end{aligned}$$




Fig. 7.12

**Example 2.** In Fig 7.13 , if  $EF \parallel BC$  and  $FG \parallel CD$ , prove that

$$\frac{AG}{AD} = \frac{AE}{AB}$$

**Solution :** **Given :**  $EF \parallel BC$  and  $FG \parallel CD$

**To prove :**  $\frac{AG}{AD} = \frac{AE}{AB}$

**Proof:** In  $\triangle ABC$ ,  
 $EF \parallel BC$  (Given)

$\therefore$  by Basic Proportionality Theorem,(vide Example 1.)

$$\frac{AE}{AB} = \frac{AF}{AC} \quad \dots(i)$$

Again, in  $\triangle ACD$ ,

$FG \parallel CD$

$$\therefore \frac{AF}{AC} = \frac{AG}{AD} \quad \dots(ii)$$

From (i) and (ii), we have




Fig. 7.13



$$\frac{AE}{AB} = \frac{AG}{AD} \text{ or } \frac{AG}{AD} = \frac{AE}{AB}$$

**Example 3.** Prove that the line joining the mid - points of two sides of a triangle is parallel to the third side.

**Solution:** **Given :** The line DE joining the mid point D of AB and the mid point E of AC of  $\triangle ABC$  (Fig 7.14).

**To prove :**  $DE \parallel BC$

**Proof :** Since D is the mid point of AB,


$$\therefore AD = DB$$

$$\Rightarrow \frac{AD}{DB} = 1$$

$$\text{Similarly, } \frac{AE}{EC} = 1$$


$$\therefore \frac{AD}{DB} = \frac{AE}{EC}$$

So, by the converse of the Basic Proportionality Theorem,  $DE \parallel BC$ .



**Example 4.** Prove that the line segments, joining the mid-points of the adjacent sides of a quadrilateral form a parallelogram.

**Solution :** **Given:** The line segments EF, FG, GH, HE formed by joining the mid points E, F, G, H respectively of AB, BC, CD, DA of a quadrilateral ABCD (Fig 7.15).



**To prove :** EFGH is a parallelogram.

**Construction :** Join BD

**Proof :** In  $\triangle ABD$ , E and H are the mid - points of AB and AD respectively.

$$\therefore EH \parallel BD \quad \dots\dots\dots(1)$$

Again, in  $\triangle CDB$ , F and G are the mid-points of CB and CD respectively.

$$\therefore FG \parallel BD \quad \dots\dots\dots(2)$$

From (1) and (2), we have

$$EH \parallel FG$$

Similarly,  $EH \parallel FG$   
 $\therefore EFGH$  is a parallelogram.


**Example 5 .** The diagonals of a quadrilateral  $ABCD$  intersect each other at the point  $O$  such that  $\frac{AO}{BO} = \frac{CO}{DO}$ . Show that  $ABCD$  is a trapezium.

**Solution :** Draw  $XO \parallel AB$  meeting  $AD$  at  $X$ . In  $\triangle DAB$ ,  $XO \parallel AB$

$\therefore$  by Basic Proportionality Theorem.

$$\frac{DX}{XA} = \frac{DO}{OB}$$

$$\Rightarrow \frac{XA}{DX} = \frac{OB}{DO} \dots\dots\dots(i)$$



But, it is given that  $\frac{AO}{BO} = \frac{CO}{DO}$

or  $\frac{AO}{CO} = \frac{BO}{DO}$

or  $\frac{AO}{CO} = \frac{OB}{DO} \dots\dots\dots(ii)$

From (i) and (ii),  $\frac{XA}{DX} = \frac{AO}{CO}$

$\therefore$  by the converse of the Basic Proportionality Theorem,  
 $XO \parallel DC$ .

But, by construction  $XO \parallel AB$ .

$\therefore AB \parallel DC$

Hence,  $ABCD$  is a Trapezium.

**EXERCISE 7.2**

1. Prove that any line drawn parallel to the parallel sides of a trapezium divides the oblique sides proportionally.
2. Prove that the line drawn from the mid-point of one side of a triangle parallel to another side bisects the third side.
3. Any point  $X$  inside a  $\triangle ABC$  is joined to the vertices. From a point  $P$  on  $AX$ ,  $PQ$  is drawn parallel to  $AB$  meeting  $XB$  at  $Q$  and  $QR$  is drawn parallel to  $BC$  meeting  $XC$  at  $R$ . Prove that  $PR \parallel AC$ .

4. Let ABC be a triangle and D, E be two points on side AB such that AD=BE. If DP||BC and EQ || AC, where P and Q lie on AC and BC respectively, Prove that PQ || AB.
5. ABCD is a quadrilateral. P,Q,R and S are the points on the sides AB, BC, CD and DA respectively such that AP:PB=AS:SD=CQ:QB=CR:RD. Prove that PQRS is a parallelogram.
6. On three line segments OA, OB and OC, points L,M,N respectively are so chosen that LM || AB and MN || BC but neither L, M, N nor A, B, C are collinear. Show that LN || AC.
7. If three or more parallel lines are intersected by two transversals, prove that the intercepts made by them on the transversals are proportional.
8. **ABCD is a parallelogram and P is a point on the side BC. DP when produced meets AB produced at L, prove that**
- (i)  $\frac{DP}{PL} = \frac{DC}{BL}$       (ii)  $\frac{DL}{DP} = \frac{AL}{DC}$
9. In a  $\triangle ABC$ , D and E are points on the sides AB and AC respectively, such that  $AD \times EC = AE \times DB$ . Prove that  $DE \parallel BC$ .
10. The side BC of a triangle ABC is bisected at D, O is any point on AD. BO and CO produced meet AC and AB at E and F respectively and AD is produced to X so that D is the mid point of OX. Prove that  $AO:AX=AF:AB$  and show that  $FE \parallel BC$ .
11. P is the mid-point of the side BC of a triangle ABC and Q is the mid-point of AP. If BQ, when produced, meets AC at L, prove that  $LA = \frac{1}{3}CA$ .
12. Two triangles ABC and DBC lie on the same side of BC. From a point P on BC, PQ is drawn parallel to BA and meeting AC at Q. PR is also drawn parallel to BD meeting CD at R. Prove that  $QR \parallel AD$ .

#### 7.4 Internal and External bisectors of an angle of a triangle.

You are already familiar with the notion of the internal and external bisectors of an angle of a triangle. In this section, we will derive some properties of internal and external bisectors of an angle of a triangle. First, let us learn the notion of internal and external division of a line segment. A point C on the segment AB is said to divide AB internally in the

ratio m:n if  $\frac{AC}{CB} = \frac{m}{n}$  i.e., if  $AC:CB = m:n$  (Fig 7.17).




Fig. 7.17

Whereas, a point C on the line AB, not lying between A and B, is said to divide AB externally in the ratio  $m : n$  if  $\frac{AC}{CB} = \frac{m}{n}$  (Fig 7.18). In the later case the point C , lies on AB produced or BA produced according as  $m > n$  or  $m < n$ .




Fig. 7.18

Now, to examine the property of internal bisector of an angle of a triangle , let us perform the following activity.

**Activity :**

Draw an angle  $\angle XAY$  and mark points  $P_1, P_2, P_3,$  and B on the arm AX such that  $AP_1 = P_1P_2 = P_2P_3 = P_3B = 1$  unit (Fig 7.19). Also, mark points  $Q_1, Q_2, Q_3, Q_4, Q_5$  and C on the arm AY such that  $AQ_1 = Q_1Q_2 = Q_2Q_3 = Q_3Q_4 = Q_4Q_5 = Q_5C = 1$  unit.

Join BC. Here, we have

$$\frac{AB}{AC} = \frac{4}{6} = \frac{2}{3}$$

Draw bisector of  $\angle XAY$  to intersect BC at D (Fig. 7.19). Measure lengths BD and DC and compute  $\frac{BD}{DC}$ .

You will find that  $\frac{BD}{DC} = \frac{2}{3}$

$$\therefore \frac{AB}{AC} = \frac{BD}{DC}$$




Fig 7.19

It means that the internal bisector of  $\angle A$  of  $\triangle ABC$  divides the opposite side BC internally in the ratio  $AB:AC$ . This result is stated and proved as a theorem given below.

**Theorem 7.3** The internal bisector of an angle of a triangle divides the opposite side internally in the ratio of the other two sides.

**Given:** Internal bisector AD of the  $\angle A$  of  $\triangle ABC$  meeting BC at D (Fig 7.20).

**To prove :**  $\frac{BD}{DC} = \frac{AB}{AC}$

**Construction :** Draw CE parallel to DA meeting BA produced at E (Fig 7.20).

**Proof :** By construction,

$$DA \parallel CE$$

$\therefore$  by Basic Proportionality theorem,

$$\frac{BD}{DC} = \frac{BA}{AE} \text{ ---(1)}$$

DA  $\parallel$  CE and BE is a transversal.

$$\therefore \angle BAD = \angle AEC \text{ ---(2)}$$

(corresponding angles)

Again, DA  $\parallel$  CE and AC is a transversal.

$$\therefore \angle DAC = \angle ACE \text{ ---(3) (alternative angles)}$$

But, AD bisects  $\angle BAC$

$$\therefore \angle BAD = \angle DAC$$

$\therefore$  from (2) and (3)


$$\angle AEC = \angle ACE$$

$$\Rightarrow AC = AE \text{ (sides opposite to equal angles of a triangle)}$$

Putting this value in (1), we have

$$\frac{BD}{DC} = \frac{BA}{AC}$$

Hence,  $\frac{BD}{DC} = \frac{AB}{AC}$ .



**Example 6.** If the diagonal AC of a quadrilateral ABCD bisects both  $\angle A$  and  $\angle C$ ,

show that  $\frac{AB}{AD} = \frac{BC}{CD}$ .

**Solution :** **Given :** Diagonal AC of a quadrilateral ABCD bisecting both  $\angle A$  and  $\angle C$  (Fig. 7.21)

**To prove :**  $\frac{AB}{AD} = \frac{BC}{CD}$

**Construction :** Join BD intersecting AC at O.

**Proof :** In  $\triangle ABD$ , AO is the internal bisector of  $\angle A$ .


$$\begin{aligned} \therefore \frac{BO}{OD} &= \frac{AB}{AD} \\ \Rightarrow \frac{OB}{OD} &= \frac{AB}{AD} \quad \text{---(1)} \end{aligned}$$

In  $\triangle BCD$ ,  $CO$  is the internal bisector of  $\angle C$

$$\begin{aligned} \therefore \frac{BO}{OD} &= \frac{BC}{CD} \\ \Rightarrow \frac{OB}{OD} &= \frac{BC}{CD} \quad \text{---(2)} \end{aligned}$$

From (1) and (2), we have

$$\frac{AB}{AD} = \frac{BC}{CD}.$$



**Example 7.**  $O$  is any point inside a triangle  $ABC$ . The bisector of  $\angle AOB$ ,  $\angle BOC$  and  $\angle COA$  meet the sides  $AB$ ,  $BC$  and  $CA$  at points  $D$ ,  $E$  and  $F$  respectively. Show that  $AD \times BE \times CF = DB \times EC \times FA$ .

**Solution :** In  $\triangle AOB$ ,  $OD$  is the internal bisector of  $\angle AOB$

$$\therefore \frac{AD}{DB} = \frac{OA}{OB} \quad \text{---(1)}$$

In  $\triangle BOC$ ,  $OE$  is the internal bisector of  $\angle BOC$ .


$$\therefore \frac{BE}{EC} = \frac{OB}{OC} \quad \text{---(2)}$$

Again, in  $\triangle AOC$ ,  $OF$  is the internal bisector of  $\angle AOC$ .

$$\therefore \frac{CF}{FA} = \frac{OC}{OA}$$

From (1), (2), (3), we have

$$\begin{aligned} \Rightarrow \frac{AD}{DB} \times \frac{BE}{EC} \times \frac{CF}{FA} &= \frac{OA}{OB} \times \frac{OB}{OC} \times \frac{OC}{OA} \\ \Rightarrow \frac{AD}{DB} \times \frac{BE}{EC} \times \frac{CF}{FA} &= 1 \\ \Rightarrow AD \times BE \times CF &= DB \times EC \times FA. \end{aligned}$$



### EXERCISE 7.3

- AD is a median of  $\triangle ABC$ . The bisector of  $\angle ADB$  and  $\angle ADC$  meet AB and AC at E and F respectively. Prove that  $EF \parallel BC$ .
- If the bisector of an angle of a triangle bisects the opposite sides, prove that the triangle is isosceles.
- In  $\triangle ABC$ , the bisector of  $\angle B$  meets AC at D. A line PQ is drawn parallel to AC meeting AB, BC and BD at P, Q and R respectively. Show that  
(i)  $PR \times BQ = QR \times BP$  (ii)  $AB \times CQ = BC \times AP$
- If the medians of a triangle are the bisectors of the corresponding angles of the triangle, prove that the triangle is equilateral.
- The bisectors of the angles  $\angle B$  and  $\angle C$  of a triangle ABC meet the opposite sides at D and E respectively. If  $ED \parallel BC$ , prove that the triangle is isosceles.

#### 7.5 Criteria for similarity of triangles

In section 7.3, we have already stated that two triangles are similar if

- their corresponding angles are equal and
- their corresponding sides are in the same ratio.

In other words, two triangles ABC and DEF are similar, if

- $\angle A = \angle D$ ,  $\angle B = \angle E$ ,  $\angle C = \angle F$  and
- $\frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD}$





Fig. 7.23

Here, we see that A corresponds to D, B corresponds to E and C corresponds to F. At this situation we write  $\triangle ABC \parallel \triangle DEF$  [read as “triangle ABC is similar to triangle DEF”]. As done in the case of congruency, for the similarity of the triangles also, correct correspondence of their vertices should be maintained. For example, in figure 7.25, we write  $\triangle ABC \parallel \triangle DEF$  [but neither as  $\triangle ABC \parallel \triangle EDF$  nor  $\triangle ABC \parallel \triangle EFD$  nor  $\triangle ABC \parallel \triangle FED$  nor  $\triangle ABC \parallel \triangle FDE$  nor  $\triangle ABC \parallel \triangle DFE$ ].

In this section, using the theorems discussed in earlier sections, we shall derive some criteria for similarity of triangles which in turn will imply that either of the above two conditions is sufficient to define the similarity of two triangles. Now, let us perform the following activity.

**Activity:**

Draw two line segments BC and EF of lengths say, 4cm and 6cm respectively. At B and C, construct angles XBC and YCB of some measures say,  $60^\circ$  and  $45^\circ$  respectively.



*Fig. 7.24*

Also, construct angles ZEF and WFE of  $60^\circ$  and  $45^\circ$  respectively. Suppose rays BX and CY intersect each other at A and rays EZ and FW intersect each other at D (Fig 7.24)

Now, in  $\triangle ABC$ , and  $\triangle DEF$

$$\angle B = \angle E, \angle C = \angle F, \text{ and so } \angle A = \angle D$$

i.e. corresponding angles are equal

Observe that 
$$\frac{BC}{EF} = \frac{4}{6} = \frac{2}{3}$$

Now, measure AB, DE, CA and FD. You will find that

$$\frac{AB}{DE} = \frac{2}{3} = \frac{CA}{FD}$$

$$\therefore \frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD}$$

i.e. the corresponding sides are in the same ratio. It follows that  $\triangle ABC \sim \triangle DEF$ .

Repeating this activity by constructing several pairs of triangles having their corresponding angles equal, you will get the same result. Thus, we have the following criterion for the similarity of two triangles.



**Theorem 7.4 (AAA Similarity):** If the corresponding angles of two triangles are equal, then the triangles are similar.

**Given :** Two triangles ABC and DEF such that  $\angle A = \angle D$ ,  $\angle B = \angle E$ ,  $\angle C = \angle F$ .

**To prove :**  $\triangle ABC \sim \triangle DEF$ .

**Construction :** Take points M and N on  $\overline{AB}$  and  $\overline{AC}$  respectively such that  $AM=DE$  and  $AN=DF$ . Join MN.




Fig. 7.25

**Proof :** Three cases arise :

**Case I**  $AB > DE$ .

Here, M lies in AB {Fig. 7.25(i)}.

In  $\Delta^s$  AMN and DEF

$AM = DE, AN = DF$  and  $\angle A = \angle D$ .

$\therefore$  by SAS axiom

$\Delta AMN \cong \Delta DEF$

It gives  $\angle AMN = \angle DEF$

But  $\angle DEF = \angle ABC$  (given)

$\therefore \angle AMN = \angle ABC$

$\Rightarrow MN \parallel BC$

$\Rightarrow \frac{AM}{AN} = \frac{DM}{DN}$  [by a deduction from the Basic Proportionality Theorem]

$\Rightarrow \frac{AB}{DE} = \frac{AC}{DF}$

Similarly, it can be shown that

$$\frac{AB}{DE} = \frac{BC}{EF}$$

$$\therefore \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

**Case II**  $AB = DE$

Here, M coincides with B [fig. 7.25(ii)]

Then, by ASA congruence theorem,

$\Delta ABC \cong \Delta DEF$

$\therefore AB = DE, BC = EF, AC = DF$

$$\therefore \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

**Case III**  $AB < DE$

Here, M lies on AB produced [fig 7.25 (iii)]

As in case I, it can be shown that  $BC \parallel MN$  and ultimately, we obtain

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

Thus, in all possible cases we get

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

i.e. the corresponding sides of  $\triangle ABC$  and  $\triangle DEF$  are proportional. As the corresponding angles are given to be equal, we can conclude that  $\triangle ABC \sim \triangle DEF$ .

**Cor. (AA similarity)**

If two angles of one triangle are respectively equal to two angles of another triangle, then the two triangles are similar.


The proof follows immediately after applying the angle sum property of a triangle.

In the proof of the AAA similarity, we have seen that if the corresponding angles of two triangles are equal, then the corresponding sides are in the same ratio.

Is the converse of this result true? That is, if the sides of a triangle are respectively, proportional to the sides of another triangle, are the corresponding angles equal? To examine it, let us perform the following activity.

**Activity :**

Draw two triangles ABC and DEF such that  
 $AB = 4\text{cm}$ ,  $BC = 5\text{cm}$ ,  $CA = 6\text{cm}$  and  
 $DE = 6\text{cm}$ ,  $EF = 7.5\text{cm}$ ,  $FD = 9\text{cm}$  (Fig. 7.26)



**Fig. 7.26**

Then, we have  $\frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD} = \frac{2}{3}$  Measure  $\angle A, \angle D, \angle B, \angle E, \angle C, \angle F$ .

Observe that  $\angle A = \angle D, \angle B = \angle E$  and  $\angle C = \angle F$  i.e. the corresponding angles are equal. Repeating this activity by taking corresponding sides of two triangles in the same ratio, we will find that the corresponding angles are equal and hence they are similar. Let us now prove this result as a criterion of similarity of two triangles.

**Theorem 7.5 (SSS similarity)**

**If the corresponding sides of two triangles are in the same ratio, then the triangles are similar.**

**Given :** Two triangles ABC and DEF such that

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} \quad (\text{Fig 7.25})$$

**To prove :**  $\Delta ABC \sim \Delta DEF$

**Construction :** Take points M and N on  $\overline{AB}$  and  $\overline{AC}$  respectively such that  $AM=DE$  and  $AN=DF$ . Join MN(Fig 7.25).

**Proof :** Three cases arise.

**Case I**  $AB > DE$

Here, M lies in AB (Fig 7.25(i)).

Now, 
$$\frac{AB}{DE} = \frac{AC}{DF}$$

$$\Rightarrow \frac{AB}{AM} = \frac{AC}{AN}$$

$$\Rightarrow \frac{AB}{AM} - 1 = \frac{AC}{AN} - 1$$

$$\Rightarrow \frac{AB - AM}{AM} = \frac{AC - AN}{AN}$$

$$\Rightarrow \frac{MB}{AM} = \frac{NC}{AN}$$

$$\Rightarrow \frac{AM}{MB} = \frac{AN}{NC}$$

$\therefore$  by the converse of the Basic Proportionality Theorem, we have,  
 $MN \parallel BC$

$\therefore \angle AMN = \angle ABC$  and  $\angle ANM = \angle ACB$  (corresponding angles)

So, by AA similarity

$$\Delta ABC \sim \Delta AMN$$

$$\therefore \frac{AB}{AM} = \frac{BC}{MN}$$

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{MN} \quad \text{---(1)}$$

But, 
$$\frac{AB}{DE} = \frac{BC}{EF} \quad \text{---(2)}$$

$$\text{From (1) and (2), } \frac{BC}{MN} = \frac{BC}{EF}$$

$$\therefore MN = EF$$

So, by SSS congruence Theorem,

$$\triangle AMN \cong \triangle DEF$$

Therefore,  $\triangle AMN$  and  $\triangle DEF$  are equiangular i.e., the corresponding angles are same.

But,  $\triangle ABC \sim \triangle AMN$  and so, they are equiangular. Therefore,  $\triangle ABC$  and  $\triangle DEF$  are equiangular. It is also given that the corresponding sides of  $\triangle ABC$  and  $\triangle DEF$  are in the same ratio. Hence  $\triangle ABC \sim \triangle DEF$ .

**Case II**  $AB=DE$ .

Here M coincides with B (Fig 7.25(ii))

$$\text{Now, } 1 = \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

$$AB = DE, BC = EF, AC = DF.$$

So, by SSS congruence theorem,

$$\triangle ABC \cong \triangle DEF$$

Hence,  $\triangle ABC \sim \triangle DEF$ .

**Case III**  $AB < DE$ .

Here, M lies on AB produced (Fig 7.25(iii)). As in the case of (i), we can show that  $MN \parallel BC$  and ultimately we can obtain  $\triangle ABC \sim \triangle DEF$ .

Thus, in all possible cases,  $\triangle ABC \sim \triangle DEF$ .

In view of the above two theorems, the following definitions of the similarity of two triangles can be given.

**Definition 1.** Two triangles are similar if the corresponding angles are equal.


**Definition 2.** Two triangles are similar if the corresponding sides are in the same ratio.

In class IX, we have learnt about various criteria for congruency of two triangles. Observe that, corresponding to the SSS congruency criterion there is SSS similarity criterion. Again, there is SAS congruency criterion for congruency of two triangles. It is natural to look for a similarity criterion corresponding to SAS congruency criterion of triangles. To check the existence of such criterion, let us perform the following activity.

**Activity :**

Draw two triangles ABC and DEF such that

$AB = 8\text{ cm}$ ,  $\angle B = 30^\circ$ ,  $BC = 4\text{ cm}$  and  $DE = 6\text{ cm}$ ,  $\angle E = 30^\circ$ ,  $EF = 3\text{ cm}$ . (Fig 7.27).




**Fig. 7.27**

Then,  $\frac{AB}{DE} = \frac{4}{3} = \frac{BC}{EF}$  i.e.  $\frac{AB}{BC} = \frac{DE}{EF}$  and  $\angle B = \angle E$ . That is, one angle of  $\triangle ABC$  equals one angle of  $\triangle DEF$  and the sides including these angles are in the same ratio. Measure  $\angle A$ ,  $\angle C$ ,  $\angle D$  and  $\angle F$ . Observe that  $\angle A = \angle D$  and  $\angle C = \angle F$ . So, by AA similarity,  $\triangle ABC \sim \triangle DEF$ .

This observation is stated and proved as a theorem given below.

**Theorem 7.6 (SAS similarity)**

**If one angle of a triangle is equal to one angle of another triangle and the sides including these angles are in the same ratio, the triangles are similar.**



**Fig. 7.28**

**Given :**  $\triangle ABC$  and  $DEF$  in which  $\angle A = \angle D$  and  $\frac{AB}{DE} = \frac{AC}{DF}$  (Fig 7.28).

**To prove :**  $\triangle ABC \sim \triangle DEF$ .

**Proof :** If  $AB=DE$ , then  $\frac{AC}{DF} = \frac{AB}{DE} = 1$

$$\Rightarrow AC = DF$$

$\therefore$  by SAS axiom,  $\triangle ABC \cong \triangle DEF$

So,  $\triangle ABC \sim \triangle DEF$

If  $AB \neq DE$ , then one is greater than the other. Without loss of generality, we can take  $AB > DE$ . Take points M and N on AB and AC respectively such that  $AM=DE$  and  $AN=DF$ . Join MN (Fig 7.28)

In  $\triangle AMN$  and  $DEF$

$$AM = DE, AN = DF \text{ and } \angle A = \angle D .$$

$\therefore$  by SAS congruence axiom,

$$\triangle AMN \cong \triangle DEF$$

$\therefore \angle AMN = \angle DEF, \angle ANM = \angle DFE$  ----(1)

$$\text{Now, } \frac{AB}{DE} = \frac{AC}{DF} \quad (\text{given})$$

$$\therefore \frac{AB}{AM} = \frac{AC}{AN} \quad (\text{by construction } AM=DE, AN=DF)$$

$$\Rightarrow \frac{AM}{AB} = \frac{AN}{AC}$$

So,  $MN \parallel BC$ . (by converse of the Basic proportionality theorem)

$\therefore \angle AMN = \angle ABC$  and  $\angle ANM = \angle ACB$

ie.  $\angle DEF = \angle ABC$  and  $\angle DFE = \angle ACB$  (using (1))

So, by AA similarity,

$$\triangle DEF \sim \triangle ABC$$

Hence,  $\triangle ABC \sim \triangle DEF$ .

Using the above criteria of similarity of two triangles, we can have the following theorem.

### Theorem 7.7

**If a perpendicular is drawn from the vertex of the right angle of a right triangle to the hypotenuse, the triangles on each side of the perpendicular are similar to the whole triangle and to each other.**

**Given :** Right  $\triangle ABC$  right angled at A and  $AD \perp BC$  (Fig 7.29)

- To prove :** (i)  $\triangle DBA \sim \triangle ABC$   
 (ii)  $\triangle DAC \sim \triangle ABC$   
 (iii)  $\triangle DBA \sim \triangle DAC$




Fig. 7.29

**Proof :** (i) In  $\triangle DBA$  and  $\triangle ABC$ ,  
 $\angle DBA = \angle ABC$  (same angle)  
 $\angle ADB = \angle CAB = 90^\circ$   
 $\therefore$  by AA similarity  $\triangle DBA \sim \triangle ABC$

(ii) In  $\triangle DAC$  and  $\triangle ABC$   
 $\angle DCA = \angle ACB$  (same angle)  
 $\angle ADC = \angle BAC = 90^\circ$   
 $\therefore$  by AA similarity  $\triangle DAC \sim \triangle ABC$ .

(iii) In  $\triangle ADB$ ,

$$\angle DBA + \angle DAB = 90^\circ \left[ \because \angle ADB = 90^\circ \right]$$

$$\text{But } \angle DBA + \angle DCA = 90^\circ \left[ \text{in } \triangle ABC, \angle BAC = 90^\circ \right]$$

$$\therefore \angle DAB = \angle DCA$$

$$\text{Also, } \angle ADB = \angle CDA = 90^\circ$$

$\therefore$  by AA similarity  
 $\triangle DBA \sim \triangle DAC$ .

**Example 8.** D is a point on the side BC of a  $\triangle ABC$  such that  $\angle ADC = \angle BAC$ .  
 Prove that  $CA^2 = CB \times CD$ .

**Solution :** **Given :** The point D on BC of  $\triangle ABC$  such that  $\angle ADC = \angle BAC$ .

**To prove :**  $CA^2 = CB \times CD$ .

**Proof :**  $\triangle ABC$  and  $\triangle DAC$ ,  
 $\angle ACB = \angle DCA$  (same angle)  
 $\angle BAC = \angle ADC$  (given)  
 $\therefore$  by AA similarity  
 $\triangle ABC \sim \triangle DAC$

$$\therefore \frac{AB}{DA} = \frac{BC}{AC} = \frac{CA}{CD}$$

$$\Rightarrow \frac{BC}{AC} = \frac{CA}{CD}$$

$$\Rightarrow \frac{BC}{CA} = \frac{CA}{CD}$$

$$\Rightarrow BC \times CD = CA^2$$

Hence,  $CA^2 = CB \times CD$ .




Fig. 7.30



**Example 9.** Two vertical poles of heights  $a$  metres and  $b$  metres are  $p$  metres apart. Prove that the height of the point of intersection of the lines joining the top of each pole to the foot of the other is  $\frac{ab}{a+b}$  metres.

**Solution :** Let  $AB$  and  $CD$  be the two vertical poles of heights  $a$  metres and  $b$  metres where  $A, C$  are the feet of the poles so that,  $AC = p$  metres.

Let  $AD$  and  $CB$  intersect at  $O$  (Fig 7.31). Draw  $OE \perp AC$ .

Let  $AE = x$  metres,  $CE = y$  metres. Also, let  $OE = h$  metres.

Then,  $p = AC = AE + EC = x + y$

$$\Rightarrow p = x + y \text{ ---(1)}$$

In  $\Delta^s ABC$  and  $EOC$

$$\angle BAC = \angle OEC = 90^\circ$$

$$\angle ACB = \angle ECO \quad (\text{same angle})$$

$\therefore$  by AA similarity,

$$\Delta ABC \sim \Delta EOC$$

$$\therefore \frac{AB}{EO} = \frac{AC}{EC}$$

$$\Rightarrow \frac{a}{h} = \frac{p}{y}$$

$$\Rightarrow y = \frac{ph}{a} \text{ ---(2)}$$

In  $\Delta^s ACD$  and  $AEO$ ,

$$\angle ACD = \angle AEO = 90^\circ$$

$$\angle DAC = \angle OAE \quad (\text{same angle})$$

$\therefore$  by AA similarity,

$$\Delta ACD \sim \Delta AEO$$

$$\therefore \frac{CD}{EO} = \frac{AC}{AE}$$

$$\Rightarrow \frac{b}{h} = \frac{p}{x}$$




Fig. 7.31

$$\Rightarrow x = \frac{ph}{b} \quad \text{---(3)}$$

From (1), (2), (3) we have

$$p = \frac{ph}{b} + \frac{ph}{a}$$

$$\Rightarrow 1 = h \left( \frac{1}{b} + \frac{1}{a} \right)$$

$$\Rightarrow 1 = h \left( \frac{a+b}{ab} \right)$$

$$\Rightarrow h = \frac{ab}{a+b} \text{ metres.}$$

Hence, the required height is  $\frac{ab}{a+b}$ .

**Example 10.** BP and EQ are medians of the triangles ABC and DEF respectively. If

$$\frac{BC}{EF} = \frac{AC}{DF} = \frac{BP}{EQ}, \text{ prove that } \Delta ABC \sim \Delta DEF.$$

**Solution :** **Given :** medians BP and EQ of  $\Delta^s ABC$  and  $DEF$  respectively.

$$\text{Also, } \frac{BC}{EF} = \frac{AC}{DF} = \frac{BP}{EQ}.$$

**To prove :**  $\Delta ABC \sim \Delta DEF$ .

**Proof :** Since BP is the median of  $\Delta ABC$ ,

$$2AP = 2PC = AC.$$

Similarly,  $2DQ = 2QF = DF$ .

It is given that

$$\frac{BC}{EF} = \frac{AC}{DF} = \frac{BP}{EQ}$$

$$\Rightarrow \frac{BC}{EF} = \frac{2 \times PC}{2 \times QF} = \frac{BP}{EQ}$$


$$\Rightarrow \frac{BC}{EF} = \frac{PC}{QF} = \frac{BP}{EQ}$$

$\therefore$  by SSS similarity,

$$\Delta BPC \sim \Delta EQF$$

$$\therefore \angle BCP = \angle EQF$$

$$\Rightarrow \angle BCA = \angle EFD$$



**Fig. 7.32**

$$\text{But, } \frac{BC}{EF} = \frac{AC}{DF}$$

$$\therefore \text{ by SAS similarity}$$

$$\triangle ABC \cong \triangle DEF.$$

**Example 11.** ABCD is a trapezium in which  $AB \parallel DC$ . If the diagonals AC and BD intersect at P and  $\triangle APD \sim \triangle BPC$ , prove that  $AD = BC$ .

**Solution :** **Given :** Trapezium ABCD in which  $AB \parallel DC$  and  $\triangle APD \sim \triangle BPC$  where P is the point of intersection of the diagonals AC and BD.

**To prove :**  $AD = BC$

**Proof :** Since  $\triangle APD \sim \triangle BPC$ ,

$$\frac{AP}{BP} = \frac{DP}{CP} = \frac{AD}{BC} \quad \text{-----(1)}$$

Again, it is given that  $AB \parallel DC$

$$\therefore \angle CDP = \angle ABP \quad (\text{alternate angles})$$

$$\angle DCP = \angle BAP \quad (\text{alternate angles})$$

$\therefore$  by AA similarity,

$$\triangle ABP \sim \triangle CDP$$

$$\therefore \frac{AP}{CP} = \frac{BP}{DP}$$

$$\Rightarrow \frac{AP}{BP} = \frac{CP}{DP}$$


$$\Rightarrow \frac{DP}{CP} = \frac{CP}{DP} \quad [\text{using (1)}]$$

$$\Rightarrow DP^2 = CP^2$$

$$\Rightarrow DP = CP$$

$$\therefore \text{From (1), } \frac{AP}{BP} = \frac{AD}{BC} = \frac{DP}{CP} = 1$$

$$\therefore AD = BC.$$



**Fig. 7.33**

**Example 12.** ABC is right triangle right angled at A and AD is drawn perpendicular to BC meeting it at D. Prove that  $AB^2 = BD \times BC$ .

**Solution :** **Given :** A right  $\triangle ABC$  right angled at A and  $AD \perp BC$ .

**To prove :**  $AB^2 = BD \times BC$

**Proof :** Since AD is the altitude of the rt  $\triangle ABC$ ,  
right angled A,


$$\triangle ABC \sim \triangle DBA$$

$$\therefore \frac{AB}{DB} = \frac{BC}{BA} = \frac{AC}{DA}$$

$$\Rightarrow \frac{AB}{DB} = \frac{BC}{AB}$$

$$\Rightarrow AB^2 = DB \times BC$$

$$\therefore AB^2 = BD \times BC$$



*Fig. 7.34*

### EXERCISE 7.4

- If two triangles are similar, prove that the corresponding
  - medians are proportional
  - altitudes are proportional
- ABCD is a parallelogram and E is the mid-point of AB. If F is the point of intersection of  $\overline{DE}$  and  $\overline{BC}$ , prove that  $BC = BF$ .
- ABC is an isosceles triangle in which  $AB = AC$  and D is a point on AC such that  $BC^2 = AC \cdot CD$ . Prove that  $BC = BD$ .
- If in a triangle ABC,  $AD \perp BC$  and  $AD^2 = BD \cdot DC$ , prove that  $\angle BAC = 90^\circ$ .
- Find the height of a vertical tower which casts a shadow of length 36m at the time when the shadow of a vertical post of length 5m is 3m. (Ans. 60m)
- Prove that the line segments joining the mid-points of the sides of a triangle form four triangles, each of which is similar to the original triangle.
- In  $\triangle ABC$ , DE is parallel to base BC with D on AB and E on AC. If  $\frac{AD}{DB} = \frac{2}{3}$ , find  $\frac{BC}{DE}$ . (Ans.  $\frac{5}{2}$ )
- In a  $\triangle ABC$ , P and Q are points on AB and AC respectively such that  $PQ \parallel BC$ . Prove that median AD bisects PQ.
- ABC is an isosceles triangle in which  $AB = AC$ . AD is drawn perpendicular to BC. From a point E on CB produced, EF is drawn perpendicular to AC. Prove that  $\triangle ADC \sim \triangle EFC$ .

10. **ABC is a right triangle right angled at B and D is the foot of the perpendicular drawn from B on AC. If  $DM \perp BC$  and  $DN \perp AB$  where M, N lie on BC, AB respectively, prove that**  
 (i)  $DM^2 = DN \times MC$  (ii)  $DN^2 = DM \times AN$ .
11. Through the vertex D of parallelogram ABCD, a line is drawn to intersect the sides BA produced and BC produced at E and F respectively. Prove that  $\frac{AD}{AE} = \frac{BF}{BE} = \frac{CF}{CD}$
12. The perimeter of two similar triangles ABC and PQR are respectively, 72 cm and 48 cm. If PQ = 20 cm, find AB. (Ans: 30cm)
13. Two right triangles ABC and DBC are drawn on the same hypotenuse BC and on the same side of BC. If AC and BD intersect at P, prove that  $AP \times PC = BP \times PD$ .
14. If one angle of a triangle is equal to one angle of another triangle and the bisectors of these equal angles divide the opposite sides in the same ratio, prove that the triangles are similar.
15. If two triangles are equiangular, prove that the ratio of the corresponding sides is same as the ratio of the corresponding altitudes.
16. If two triangles are equiangular, prove that the ratio of the corresponding sides is same as the ratio of the corresponding angle bisector segments.

### 7.6 Areas of Similar Triangles

In this section, we will discuss a theorem concerning the ratio of areas of similar triangles.

**Theorem 7.8** The ratio of the areas of two similar triangles is equal to the ratio of the squares of their corresponding sides.

**Given :**  $\triangle ABC \sim \triangle DEF$

**To prove :**  $\frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2}$

**Construction :** Draw  $AM \perp BC$  and  $DN \perp EF$  (Fig. 7.34).

$$\begin{aligned} \text{Proof : } \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} &= \frac{\frac{1}{2} BC \times AM}{\frac{1}{2} EF \times DN} \\ &= \frac{BC}{EF} \times \frac{AM}{DN} \quad \text{-----(1)} \end{aligned}$$

In  $\Delta^s$  ABM and DEN,  
 $\angle ABM = \angle DEN$  [ $\because \Delta ABC \sim \Delta DEF$ ],  
 $\therefore \angle B = \angle E$  ]

$\angle AMB = \angle DNE = 90^\circ$

$\therefore$  by AA similarity  
 $\Delta ABM \sim \Delta DEN$

$$\therefore \frac{AB}{DE} = \frac{AM}{DN}$$

But  $\frac{AB}{DE} = \frac{BC}{EF}$  [ $\because \Delta ABC \sim \Delta DEF$ ]

$$\therefore \frac{AM}{DN} = \frac{BC}{EF}$$

Then, from (1)

$$\frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{BC}{EF} \times \frac{BC}{EF} = \frac{BC^2}{EF^2}$$

Again,  $\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$

$$\Rightarrow \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2}$$

Hence,  $\frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2}$  .




Fig. 7.34

**Example 13.** If two triangles are similar and their areas are in the ratio 4:9 ; prove that their perimeters are in ratio 2:3.

**Solution :** Let ABC and DEF be two triangles such that

$$\Delta ABC \sim \Delta DEF \text{ and } \frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{4}{9}$$

$$\text{But } \frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2}$$

$$\Rightarrow \frac{4}{9} = \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2}$$

$$\Rightarrow \frac{2}{3} = \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = \frac{AB+BC+AC}{DE+EF+DF} \text{ [taking square roots]}$$

$$\Rightarrow \frac{AB+BC+CA}{DE+EF+FD} = \frac{2}{3}$$

$$\text{Hence, } \frac{\text{perimeter of } \triangle ABC}{\text{perimeter of } \triangle DEF} = \frac{2}{3}.$$

### EXERCISE 7.5

1. If the areas of two similar triangles are equal, prove that they are congruent.
2. If D, E, F are respectively the mid-points of the sides BC, CA, AB of a  $\triangle ABC$ , prove that  $\frac{\text{ar}(\triangle DEF)}{\text{ar}(\triangle ABC)} = \frac{1}{4}$
3. Prove that the areas of two similar triangles are in the ratio of the square of the corresponding altitudes.
4. Prove that the areas of two similar triangles are in the ratio of the square of the corresponding medians.
5. Prove that the area of the equilateral triangle described on the side of a square is half the area of the equilateral triangle described on its diagonal.

### 7.7 Pythagoras Theorem

In this section, we shall prove an important Theorem known as Pythagoras Theorem (also known as Baudhayan Theorem) using the concept of similarity of triangles.

#### Theorem 7.9 (Pythagoras Theorem)

**In a right triangle the square of the hypotenuse is equal to the sum of squares of the other two sides.**

**Given :** A right  $\triangle ABC$  right angled at A (Fig 7.35).

**To prove :**  $BC^2 = AB^2 + AC^2$

**Construction :** Draw  $AD \perp BC$  (Fig 7.35)

**Proof :** In  $\triangle ABC$  and  $\triangle DBA$ ,

$$\angle BAC = \angle BDA = 90^\circ$$

$$\angle ABC = \angle DBA \quad (\text{same angle})$$

$\therefore$  by AA similarity,

$$\triangle ABC \sim \triangle DBA$$




Fig . 7.35

$$\begin{aligned} \therefore \frac{AB}{DB} &= \frac{BC}{BA} \\ \Rightarrow AB^2 &= BC \cdot BD \quad \dots\dots(1) \end{aligned}$$

Similarly, from  $\Delta^s ABC$  and  $DAC$ , it can be shown that

$$AC^2 = BC \cdot DC \quad \dots\dots(2)$$

From (1) and (2), we have

$$\begin{aligned} AB^2 + AC^2 &= BC(BD + DC) \\ &= BC \times BC \\ &= BC^2 \end{aligned}$$

Hence,  $BC^2 = AB^2 + AC^2$ .

The converse of the above theorem is also true and is proved below.

**Theorem 7.10 (Converse of Pythagoras Theorem)**

**In a triangle, if the square of one side is equal to the sum of the squares of the remaining two, the angle opposite to the first side is a right angle.**

**Given :** A  $\Delta ABC$  in which  $BC^2 = AB^2 + AC^2$  (Fig.7.36).

**To prove :**  $\angle A = 90^\circ$

**Construction :** Draw a right angle  $QPR$  such that  $PQ = AB$  and  $PR = AC$ . Join  $QR$ .

**Proof :** In  $\Delta PQR$ , by Pythagoras theorem,

$$\begin{aligned} QR^2 &= PQ^2 + PR^2 \\ &= AB^2 + AC^2 \quad (\text{by construction}) \end{aligned}$$

But,  $AB^2 + AC^2 = BC^2$  (given)

$$\therefore QR^2 = BC^2$$

$$\Rightarrow QR = BC$$

Now, in  $\Delta^s ABC$  and  $PQR$

$AB = PQ$ ,  $AC = PR$  and  $BC = QR$

$\therefore$  by SSS congruence,


$\Delta ABC \cong \Delta PQR$

$\therefore \angle BAC = \angle QPR$

But  $\angle QPR = 90^\circ$  (by construction)

$\therefore \angle BAC = 90^\circ$

i.e.  $\angle A = 90^\circ$ .



**Fig. 7.36**



**Example 14.** Prove that the sum of squares on the four sides of a rhombus is equal to the sum of squares on the diagonals.

**Solution :** **Given :** A rhombus ABCD

$$\begin{aligned}\text{To prove : } & AB^2 + BC^2 + CD^2 + DA^2 \\ & = AC^2 + BD^2\end{aligned}$$

**Proof :** We know that the diagonals of rhombus bisect each other at  $90^\circ$ . Let O be the point of intersection of AC and BD (Fig 7.37).

$$\text{Then, } BO = DO = \frac{1}{2} BD$$

$$\text{and } CO = AO = \frac{1}{2} AC.$$

From the rt.  $\triangle AOB$ , by Pythagoras theorem,


$$AB^2 = BO^2 + AO^2$$

$$\text{Similarly, } BC^2 = BO^2 + CO^2$$

$$CD^2 = CO^2 + DO^2$$

$$\text{and } DA^2 = DO^2 + AO^2$$

$$\begin{aligned}\therefore AB^2 + BC^2 + CD^2 + DA^2 &= 2[AO^2 + BO^2 + CO^2 + DO^2] \\ &= 2[(AO^2 + CO^2) + (BO^2 + DO^2)] \\ &= 2\left[2 \cdot \frac{1}{4} AC^2 + 2 \cdot \frac{1}{4} BD^2\right] \\ &= AC^2 + BD^2\end{aligned}$$



**Example 15.** D and E are points on the sides CA and CB respectively of a triangle ABC right angled at C. Prove that  $AE^2 + BD^2 = AB^2 + DE^2$ .

**Solution :** **Given :** Points D and E on CA and CB respectively of a  $\triangle ABC$  in which  $\angle C = 90^\circ$ .

$$\text{To prove : } AE^2 + BD^2 = AB^2 + DE^2$$

**Construction :** Join AE, BD, DE (Fig. 7.38)

**Proof :** From the rt.  $\triangle ACE$ , by Pythagoras theorem

$$AE^2 = AC^2 + CE^2 \quad \dots(1)$$

From the rt.  $\triangle ABC$ ,

$$BD^2 = BC^2 + CD^2 \quad \text{---(2)}$$

From the rt.  $\triangle ACB$ ,

$$AB^2 = AC^2 + BC^2 \quad \text{---(3)}$$

From the rt.  $\triangle DCE$ ,

$$DE^2 = CD^2 + CE^2 \quad \text{---(4)}$$

From (1) and (2),

$$\begin{aligned} AE^2 + BD^2 &= (AC^2 + CE^2) + (BC^2 + CD^2) \\ &= (AC^2 + BC^2) + (CD^2 + CE^2) \\ &= AB^2 + DE^2 \quad \text{(using (3) \& (4))} \end{aligned}$$

$$\therefore AE^2 + BD^2 = AB^2 + DE^2.$$




Fig. 7.38

**Example 16.** Two poles of height 9 m and 14 m stand on a horizontal ground. If the distance between their feet is 12 m, find the distance between their tops.

**Solution :** Let AB and CD be the poles of respective heights 9m and 14 m where the ends B and D are on the ground (Fig 7.39).

Draw  $AE \perp CD$ .

Then  $DE = AB = 9\text{m}$

$$\begin{aligned} \therefore CE &= CD - DE \\ &= 14 - 9 \\ &= 5 \text{ m} \end{aligned}$$

From the rt  $\triangle AEC$ , by Pythagoras theorem,

$$\begin{aligned} AC^2 &= AE^2 + CE^2 \\ &= BD^2 + CE^2 \\ &= 12^2 + 5^2 \\ &= 169 \end{aligned}$$

$$\Rightarrow AC = 13 \text{ m}$$

$\therefore$  the distance between their tops is 13 m.




Fig. 7.39

## EXERCISE 7.6

1. Sides of triangles are given below. Determine which of them are right triangles.
  - (i) 3 cm, 4cm, 5cm
  - (ii) 5cm, 12cm, 13cm
  - (iii) 4cm, 5cm, 7cm
  - (iv) 8cm, 11cm, 15cm
  - (v) 9cm, 40cm, 41cm
2. A man goes 15m due west and then 8m due north, Find his distance from the starting point.
3. The length of a side of a rhombus is 5 cm and the length of one of its diagonal is 6cm. Find the length of the other diagonal.
4. A ladder 13 m long reaches a window which is 12 m above the ground on one side of a street keeping its foot at the same point, the ladder is turned to other side of the street and it just reaches a window 5m high. Find the width of the street.
5. A ladder reaches 1m below the top of a vertical wall when its foot is at a distance of 6m from the wall. When the foot is shifted 2m nearer the wall, the ladder just reaches the top of the wall; find the height of the wall.
6. Find the length of the altitude and area of an equilateral triangle having 'a' as the length of a side.
7. If D, E, F are the mid-points of the sides BC, CA, AB of a right  $\triangle ABC$  (rt.  $\angle$  at A) respectively, prove that  $3(AD^2 + BE^2 + CF^2) = 4(AB^2 + BC^2 + CA^2)$
8. In a right triangle ABC right angled at C, if p is the length of the perpendicular segment drawn from C upon AB, then prove that
  - (i)  $ab = pc$
  - (ii)  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2}$ , where a = BC, b = CA and c = AB.
9. In an equilateral triangle ABC the side BC is trisected at D. Prove that  $9AD^2 = 7AB^2$ .
10. If A be the area of a right triangle and b one of the sides containing the right angle, prove that the length of the altitude on the hypotenuse is  $\frac{2Ab}{\sqrt{b^4 + 4A^2}}$ .
11. Prove that the equilateral triangles described on the two sides of a right angled triangle are together equal to the equilateral triangle on the hypotenuse in terms of their area.
12. If O is any point in the interior of a rectangle ABCD, prove that  $OA^2 + OC^2 = OB^2 + OD^2$ .

**ANSWER**

- |    |         |          |          |         |   |        |
|----|---------|----------|----------|---------|---|--------|
| 1. | (i) Yes | (ii) Yes | (iii) No | (iv) No | (v) Yes                                       | 2. 17m |
| 3. | 8cm     | 4. 17m   | 5. 10.5m | 6.      | $\frac{\sqrt{3}}{2}a, \frac{\sqrt{3}}{4}a^2.$ |        |

**SUMMARY**

**In this chapter, you have studied the following points:**


- Two figures having the same shape and not necessarily the same size are called similar figures.
- All congruent figures are always similar but the similar figures need not be congruent.
- Two polygons having the same number of sides are similar if (i) the corresponding angles are equal and (ii), the corresponding sides are proportional.
- If a line is drawn parallel to one side of a triangle to intersect the other two in distinct points, the other two sides are divided in the same ratio.
- If a line divides two sides of a triangle in the same ratio, the line is parallel to the third side.
- The internal bisector of an angle of a triangle divides the opposite side in the ratio of the other two sides.
- If in two triangles, the corresponding angles are equal, their corresponding sides are equal and the two triangles are similar.
- If the corresponding sides of two triangles are proportional their corresponding angles are equal and the two triangles are similar.
- If one angle of a triangle is equal to one angle of another triangle and the sides including these angles are proportional, the two triangles are similar.
- If a perpendicular is drawn from the vertex of the right angle of a right triangle to the hypotenuse, the triangles on each side of the perpendicular are similar to the whole triangle and to each other.
- The ratio of the areas of two similar triangles is equal to the ratio of the squares of their corresponding sides.
- In a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.
- In a triangle, if the square of one side is equal to the sum of squares of the remaining two, the angle opposite to the first side is a right angle.

\*\*\*\*\*

### 8.1 Introduction

In the previous classes, you have studied that a circle is a closed figure consisting of all those points which are at a constant distance (radius) from a fixed point (centre) in the plane. You have also studied various related terms like chord, diameter, circumference, arc, segment, sector etc.

Let us now examine the different possible situations when a circle and a line say  $l$  are given in a plane. Indeed, there are three possibilities (in terms of the number of common points between the circle and the line) as given in Fig. 8.1 below:



**Fig. 8.1**

In Fig. 8.1 (i), the circle and the line have no common point i.e. the line does not intersect the circle.

In Fig. 8.1 (ii), the circle and the line have two common points P and Q i.e. the line intersects the circle at two distinct points P and Q. The line  $l$  in this case is called a **secant** of the circle.

In Fig. 8.1 (iii), there is only one point P common to the circle and the line i.e. the line intersects the circle at only one point P. The line  $l$  in this case, is called a **tangent** to the circle.

In this chapter, we shall study about tangents to a circle, their existence and also study some of their properties.

circle at the point of contact. Also, note that all points of the tangent except the point of contact are exterior points of the circle.

Let us now explore some interesting properties of a tangent. To start with, draw a circle with centre  $O$  and draw a tangent  $AB$  to the circle at a point  $P$  (Fig. 8.3). Join the radius  $OP$ . Now, measure the angle between the radius  $OP$  and the tangent  $AB$ . You will find that  $OP$  is perpendicular to  $AB$ . This property of tangent is proved in the following theorem.

**Theorem 8.1** The tangent at any point of a circle is perpendicular to the radius through the point of contact.

**Given :** A tangent  $AB$  to a circle whose centre is  $O$ , the point of contact being  $P$ . (Fig. 8.4)

**To Prove :**  $OP \perp AB$

**Construction :** Let us take a point  $Q$  on  $AB$  other than  $P$  and join  $OQ$ .

**Proof:** Since all points on the tangent other than the point of contact are outside the circle,  $Q$  lies in the exterior of the circle.

$\therefore OQ > OP$




Fig. 8.4

Since this is true for every point on the tangent except the point  $P$ ,  $OP$  is the shortest of all the segments that can be drawn from the centre  $O$  to the points of the tangent  $AB$ .

And, we know that the shortest segment that can be drawn from a point to a line is the perpendicular segment from the point to the line.


Hence,  $OP \perp AB$

### 8.3 Tangents to a Circle through a point

Let us take a circle and a point  $P$ . The following three cases will arise.

**Case I (Fig. 8.5 (i)) :** If  $P$  lies inside the circle, then any line through  $P$  intersects the circle at two points. So, it is not possible to draw any tangent to the circle through  $P$ .

Thus, there is no tangent to a circle passing through a point inside the circle.



(i)

$OA = OB$  (radii of the same circle)  
 and  $OP = OP$  (common)  
 $\therefore \triangle OPA \cong \triangle OPB$  (RHS congruence)  
 Hence,  $PA = PB$  (corresponding sides)

**Remark :** The above theorem can also be proved by using the Pythagoras theorem and is left to the students as an exercise.

**Example 1.** The length of a tangent to a circle from a point P which is at a distance of 10 cm from the centre of the circle is 8 cm. Find the radius of the circle.

**Solution :** Let O be the centre of the circle and PA be a tangent segment drawn from P to the circle (Fig. 8.7).

It is given that  $OP = 10$  cm and  $PA = 8$  cm.


We are to find the radius of the circle. Join OA.

Then,  $\angle OAP = 90^\circ$  (Theorem 8.1)

Now, in the right  $\triangle OAP$ ,  
we have by Pythagoras Theorem,

$$\begin{aligned}
 OP^2 &= PA^2 + OA^2 \\
 \Rightarrow OA^2 &= OP^2 - PA^2 \\
 \Rightarrow OA &= \sqrt{OP^2 - PA^2} \\
 &= \sqrt{10^2 - 8^2} = 6
 \end{aligned}$$

Therefore, the radius of the circle is 6 cm.



**Fig. 8.7**


**Example 2.** PA and PB are tangent segments drawn from an external point P to a circle with centre O. If  $\angle AOB = 120^\circ$ , find  $\angle APB$ .

**Solution :** PA and PB are tangent segments drawn from an external point P to a circle with centre O and it is given that  $\angle AOB = 120^\circ$  (Fig. 8.8).

We are to find  $\angle APB$ .

Here, we have

$\angle OAP = \angle OBP = 90^\circ$  (Theorem 8.1)



**Fig. 8.8**

**Proof :** AB is a tangent to the smaller circle at P and OP is its radius.

$$\therefore OP \perp AB \text{ (Theorem 8.1)}$$

$$\Rightarrow \angle OPA = \angle OPB = 90^\circ$$

Now in the two right  $\Delta^s$  AOP and BOP,


$$OA = OB \text{ (radii)}$$

$$OP = OP \text{ (common)}$$

$\therefore$  By RHS congruence theorem,

$$\Delta AOP \cong \Delta BOP$$

$\therefore AP = BP$  (corresponding sides)



**Fig. 8.10**

**Example 5.** A circle touches the side BC of a  $\Delta ABC$  at P and the sides AB and AC produced, at Q and R respectively. Prove that AQ is half the perimeter of  $\Delta ABC$ .

**Given :** A circle touching the side BC of a  $\Delta ABC$  at P and AB and AC produced, at Q and R respectively (Fig. 8.11).

**To prove :**  $AQ = \frac{1}{2}(AB+BC+AC)$

**Proof :** BP and BQ are tangent segments drawn from the external point B to the circle.

$$\therefore BQ = BP \text{ (Theorem 8.2)--- (i)}$$

Similarly,

$$CP = CR \text{ ---(ii)}$$

$$\text{Also, } AQ = AR \text{ ---(iii)}$$

$$\text{Now, } 2AQ = AQ + AR$$

$$= AQ + AR \quad [\text{Using (iii)}]$$


$$= (AB + BQ) + (AC + CR)$$

$$= (AB + BP) + (AC + CP) \quad [\text{Using (i) \& (ii)}]$$

$$= AB + (BP + CP) + AC$$

$$= AB + BC + AC$$

$$\Rightarrow AQ = \frac{1}{2}(AB + BC + AC)$$



**Fig. 8.11**



**ANSWER**

1. 12 cm      2.  $40^\circ$       5. 16 cm

**SUMMARY**

**In this chapter, you have studied the following points :**

1. A line may intersect a circle at no point, two distinct points or only one point.
2. A line which intersects a circle at two points is called a secant of the circle.
3. A tangent to a circle is a line that intersects the circle at only one point. The point is called the point of contact and the tangent is said to touch the circle at the point of contact.
4. The tangent to a circle is a special case of the secant, when the two ends of the corresponding chord coincide.
5. There is one and only one tangent to a circle at any point on it.
6. The tangent at any point of a circle is perpendicular to the radius through the point of contact.
7. The number of tangents that can be drawn to a circle through a given point is
  - (i) zero if the point lies inside the circle
  - (ii) one if the point lies on the circle
  - (iii) two if the point lies outside the circle.
8. The lengths of tangents drawn from an external point to a circle are equal.

\*\*\*\*\*

### 9.1 Introduction

In the previous class, you have learnt about the need and importance of geometrical construction. You have also learnt construction of triangle under certain given data and construction of circumcircle and incircle of a given triangle. Here in this chapter we shall deal with some more construction problems. You are reminded once again that as far as practicable, only two geometrical instruments, namely a ruler and a compass will be used in geometrical construction. The analysis part of a construction is given only to reveal the clues leading to the construction process. It is not a part of the construction and will be omitted when the steps of construction are self evident.

### 9.2 Division of a line segment in a given ratio

There are two methods for dividing a given line segment in a given ratio, one based on the Basic Proportionality theorem and the other on the property of similar triangles. We deal with both the methods by taking an example. Suppose, a given line segment AB is to be divided in a given ratio, say 2:3.

#### First method :

**Given :** A line segment AB

**Required :** To divide AB in the ratio 2:3


**Analysis:** [Let C be the point of division. If a triangle  $ABP_5$  is formed with AB as a side and  $CP_2$  is drawn parallel to  $BP_5$  meeting  $AP_5$  at  $P_2$  then  $AP_2:P_2P_5=AC:CB=2:3$ .

So,  $P_2P_5$  will be of 3 units when  $AP_2$  is of 2 units and as such C can be located]

#### Steps of Construction :

- (i) Draw a ray AX inclined to AB at a certain angle preferably an acute angle.
- (ii) Taking a suitable length as unit, mark five (2+3) points  $P_1, P_2, P_3, P_4, P_5$ , on  $\overline{AX}$  such that  $AP_1=P_1P_2=P_2P_3=P_3P_4=P_4P_5=1$  unit
- (iii) Join  $P_5B$  and through  $P_2$  draw  $P_2C$  parallel to  $P_5B$  meeting AB at C.

Then C is the point on AB such that  $AC:CB=2:3$




**Proof:** In the triangle  $ABP_5$ , since  $CP_2$  is parallel to  $BP_5$  therefore  
 $AC:CB=AP_2:P_2P_5$  (Basic Proportional Theorem)  
 But  $AP_2:P_2P_5=2:3$  (by construction)  
 Hence  $AC:CB=2:3$

**Second Method:**

**Given :** A line segment AB

**Required :** To divide AB in the ratio 2:3

**Analysis:** [If similar triangles  $ACP_2$  and  $BCQ_3$  are drawn so that the point of division C lies on the segment  $P_2Q_3$ , then  $AP_2$  is parallel to  $BQ_3$  and  $AP_2:BQ_3=AC:BC=2:3$ . Hence  $BQ_3$  is of 3 unit when  $AP_2$  is of 2 units and as such C can be located]



**Steps of construction :**

- (i) Draw a ray  $AX$  inclined to  $AB$  at a certain angle.
- (ii) Draw a ray  $BY$  parallel to  $AX$  so that  $\angle ABY = \angle BAX$ .
- (iii) Mark Points  $P_1, P_2$  on  $AX$  and  $Q_1, Q_2, Q_3$  on  $BY$  such that  $AP_1 = P_1P_2 = BQ_1 = Q_1Q_2 = Q_2Q_3 (=1 \text{ unit})$
- (iv) Join  $P_2Q_3$  intersecting  $AB$  at  $C$ .

Then  $AC:CB = 2:3$

**Proof :** In the triangle  $ACP_2$  and  $BCQ_3$ ,

$$\angle P_2AC = \angle Q_3BC \quad (\because AP_2 \parallel BQ_3)$$

$$\text{and } \angle ACP_2 = \angle BCQ_3 \quad (\text{vertically opposite angles})$$

By AA similarity theorem

$$\Delta ACP_2 \sim \Delta BCQ_3$$

$$\begin{aligned} \therefore AC:CB &= AC:BC \\ &= AP_2:BQ_3 \\ &= 2:3 \end{aligned}$$

### 9.3 Construction of a triangle similar to a given triangle as per given scale factor

The construction can be illustrated by means of examples. Here, 'scale factor' means the ratio of the sides of the triangle to be constructed to the corresponding sides of the given triangle. Two cases arise according as the scale factor is less than or greater than unity. We deal with both the cases separately in the following examples.

**Example 1.** Construct a triangle similar to given triangle ABC, with its sides equal to  $\frac{3}{5}$  (scale factor <1) of the corresponding sides of the triangle ABC.


**Solution:**

**Given :** A triangle ABC

**Required :** To construct a triangle similar to the triangle ABC with its sides equal to  $\frac{3}{5}$  of the corresponding sides of the triangle ABC.

**Steps of construction :**

- (i) Draw any ray BX inclined to BC at a certain angle on the opposite sides of A.
  - (ii) Mark 5 points  $P_1, P_2, P_3, P_4, P_5$ , on BX such that  $BP_1 = P_1P_2 = P_2P_3 = P_3P_4 = P_4P_5$
  - (iii) Join  $P_5C$  and draw  $P_3C'$  parallel to  $P_5C$ , meeting BC at  $C'$ .
  - (iv) Draw  $C'A'$  parallel to CA, meeting AB at  $A'$ .
- Then  $\triangle A'BC'$  is the required triangle.



**Fig. 9.3**

**Proof :** By construction

$$\frac{BC'}{C'C} = \frac{BP_3}{P_3P_5} = \frac{3}{2}$$

$$\therefore \frac{BC}{BC'} = \frac{BC' + C'C}{B'C} = 1 + \frac{C'C}{B'C} = 1 + \frac{2}{3} = \frac{5}{3}$$

$$\therefore \frac{BC'}{BC} = \frac{3}{5}$$

Also, as  $C'A' \parallel CA$        $\triangle A'BC' \sim \triangle ABC$

$$\therefore \frac{A'B}{AB} = \frac{A'C}{AC} = \frac{BC'}{BC} = \frac{3}{5}$$

**Example 2.** Construct a triangle similar to a given triangle ABC with its sides equal to  $\frac{5}{3}$  (scale factor >1) of the corresponding sides of the triangle ABC.


**Solution :**

**Given:** A triangle ABC

**Required :** To construct a triangle similar to the triangle ABC with its sides equal to  $\frac{5}{3}$  of the corresponding sides of the triangle ABC.

**Steps of Construction :**

- (i) Draw any ray BX inclined to BC at a certain angle, on the opposite side of A.
- (ii) Mark 5 points  $P_1, P_2, P_3, P_4, P_5$ , on BX such that  
 $BP_1 = P_1P_2 = P_2P_3 = P_3P_4 = P_4P_5$
- (iii) Join  $P_3C$  and draw  $P_5C'$  parallel to  $P_3C$  and meeting BC produced at  $C'$
- (iv) Draw  $C'A'$  parallel to CA, meeting BA produced at  $A'$



**Fig. 9.4**

Then  $\Delta A'BC'$  is the required triangle.

**Proof :** Since  $AC \parallel A'C'$

$$\therefore \angle BAC = \angle BA'C'$$

$$\text{and } \angle ACB = \angle A'C'B$$

By AA similarity  $\Delta ABC \sim \Delta A'BC'$

$$\therefore \frac{AB}{A'B} = \frac{AC}{A'C'} = \frac{BC}{BC'}$$

$$\text{But } \frac{BC}{BC'} = \frac{BP_3}{BP_5} = \frac{3}{5}$$

$$\text{or } \frac{BC'}{BC} = \frac{5}{3}$$

$$\text{Hence, } \frac{A'B}{AB} = \frac{A'C'}{AC} = \frac{BC'}{BC} = \frac{5}{3}$$

**Note :** In the constructions of examples 1 and 2, you could take any ray through any vertex of the given triangle inclined at a certain angle to any of the sides through the vertex and proceed similarly.

## EXERCISE 9.1

1. Draw any line segment and divide it in the ratio  
(i) 3:7 (ii) 4:7 (iii) 3:8 (iv) 4:9
2. Draw a line segment of 8.7 cm and divide it in the ratio 5:9. Measure the two parts correct to a millimetre. (Answer : 3.1cm and 5.6 cm)
3. Construct a right triangle with sides 3cm, 4cm and 5cm. And then construct a triangle similar to the right triangle, the scale factor being  $\frac{7}{4}$ .
4. Draw a triangle with sides 5cm, 7cm and 8cm. Construct a triangle with sides  $\frac{4}{5}$  of the corresponding sides of first triangle.
5. Draw a triangle ABC in which AB=6cm, BC=7cm and  $\angle B = 45^\circ$ . Then construct a triangle whose sides are  $\frac{5}{8}$  of the corresponding sides of the triangle ABC.
6. Construct a triangle ABC in which BC=5cm,  $\angle B = 60^\circ$  and AB+AC=8cm. Then construct a triangle whose sides are  $\frac{5}{4}$  of the corresponding sides of the triangle ABC.
7. Construct an isosceles triangle of given base and altitude. Then construct another isosceles triangle whose sides are  $\frac{3}{2}$  times the corresponding sides of the first triangle.

## 9.4 Construction of a tangent (a pair of tangents) to a circle from an external point

**Given:** A circle with centre O and an external point P.


**Required:** To construct a tangent (a pair of tangents) from P to the circle.

**Analysis :** [If PA is a tangent touching the circle at A, then  $OA \perp PA$  and hence the  $\triangle POA$  is right angled at A. So, the circle described on the segment PO as a diameter, passes through A. As such, A can be located]

**Steps of construction:**

- (i) Join PO.
- (ii) Bisect PO at M
- (iii) Draw a circle with centre M and radius MP (or MO) cutting the given circle at A and A'
- (iv) Draw rays PA and PA'.

They are the pair of tangents from P to the circle.



**Fig . 9.5**

**Proof :** The angles  $\text{PAO}$  and  $\text{PA'O}$  are angles in semicircle. Therefore each of them is a right angle. Thus,  $\text{PA} \perp \text{OA}$  and also  $\text{PA}' \perp \text{OA}'$ . Hence  $\text{PA}$  and  $\text{PA}'$  are tangents to the circle from the external point  $\text{P}$ .

**Note:** The circle having centre at  $\text{M}$  and radius equal to  $\text{MP}$  intersects the given circle at two points. Therefore, only two tangents can be drawn from the external point  $\text{P}$  to the given circle.

### EXERCISE 9.2

1. Construct a pair of tangents from an external point which is at a distance of 5.5 cm from the centre of a circle whose radius is 2.5 cm.
2. Draw a circle of radius 3 cm. Construct a pair of tangents to the circle from a point 5 cm away from the centre. Measure the lengths of the tangents. Verify your measurement by geometrical reasoning.
3. Draw a circle of radius 3.5 cm. Construct a pair of tangents to the circle, inclined to each other at  $60^\circ$ . Find the distance of the point of intersection of the two tangents from the centre of the circle (Answer : 7cm)
4. Draw a circle with a given line segment  $\text{AB}$  as a diameter. Then construct a tangent to the circle from a given external point  $\text{P}$ .

### SUMMARY

**In this chapter you have studied the following constructions :**

1. Division of a line segment in a given ratio.
2. Construction of a triangle similar to a given triangle as per given scale factor.
3. Construction of a pair of tangents from an external point to a circle.

\*\*\*\*\*

### 10.1 Introduction


You have already known what the term “Trigonometry” means. It is the study of the relationships between the sides and angles of a triangle. Astronomers used trigonometry to calculate distance from the Earth to the planets and stars. It is also used in geography and in navigation and in many technologically advanced methods in Engineering.

In this chapter, we shall study some ratios of the sides of a right triangle called trigonometric ratios of an acute angle of the right triangle. We will restrict our discussion to trigonometric ratio of acute angles only, though it can be extended to those of other angles also. Here we shall establish some relationships between the trigonometric ratios and also we shall calculate numerical values of trigonometric ratio of some standard angles.

We shall use “an angle and its measure’ and “a side and its measure” synonymously.


### 10.2 Trigonometric Ratios of an Acute Angle

Let us take a right triangle ABC, right angled at B, as shown in Fig. 10.1. AC is the **hypotenuse** of the right triangle. We shall try to give names to the other two sides in view of their positions with respect to angle A. Note the position of the side BC with respect to  $\angle A$ . It faces  $\angle A$ . We call it the side opposite to  $\angle A$  or simply the “opposite side” with reference to  $\angle A$ . Similarly, the side AB is called the side adjacent to  $\angle A$  or simply “adjacent side”.



Further, let us note that relative to the  $\angle C$ , AB becomes the opposite side, BC the adjacent side while the hypotenuse AC remains the same (Fig. 10.2).

We have studied the concept of ‘ratio’ in earlier classes. We now define certain ratios involving sides of a right triangle and call them ‘trigonometric ratios’.





The trigonometric ratios of  $\angle A$  in the right triangle ABC (Fig 10.3) are defined as follows :

$$\text{sine of } \angle A = \frac{\text{opposite side}}{\text{hypotenuse}} = \frac{BC}{AC}$$


$$\text{cosine of } \angle A = \frac{\text{adjacent side}}{\text{hypotenuse}} = \frac{AB}{AC}$$

$$\text{tangent of } \angle A = \frac{\text{opposite side}}{\text{adjacent side}} = \frac{BC}{AB}$$

$$\text{cotangent of } \angle A = \frac{\text{adjacent side}}{\text{opposite side}} = \frac{AB}{BC}$$

$$\text{secant of } \angle A = \frac{\text{hypotenuse}}{\text{adjacent side}} = \frac{AC}{AB}$$

$$\text{cosecant of } \angle A = \frac{\text{hypotenuse}}{\text{opposite side}} = \frac{AC}{BC}$$



The ratio defined above are abbreviated as  $\sin A$ ,  $\cos A$ ,  $\tan A$ ,  $\cot A$ ,  $\sec A$  and  $\text{cosec } A$  respectively. You must have seen from the above definitions that the trigonometric ratios of an acute angle in a right triangle express the relationships between the angle and the ratios of its sides.

A point of caution is that  $\sin A$  is used as an abbreviation of 'sine of  $\angle A$ '.  $\sin A$  is not the product of 'sin' and 'A'. 'sin' left alone from A has no meaning. Similarly  $\tan A$  is not the product of 'tan' and 'A'. Similar interpretations follow for other trigonometric ratios.

Now, try to define trigonometric ratios of acute angle C in the right triangle ABC (Fig 10.3).


**Note:** Since the hypotenuse is the longest side of a right triangle, the value of  $\sin A$  or  $\cos A$  is always less than 1 for an acute angle A.

### 10.3 Uniqueness of Trigonometric Ratios

In this unit let us investigate the uniqueness of trigonometric ratios.

Consider a right  $\triangle ABC$ , right angled at B.

Take a point P on the hypotenuse AC (P may be within AC or AC produced as shown in Fig 10.4), and draw PM perpendicular to AB. Will the trigonometric ratios of  $\angle A$  in the right triangle PAM differ from those of  $\angle A$  in the right triangle ABC?



To answer this, first let us note the relationship between these two triangles. Are these triangles similar? Yes, using the appropriate similarity criterion (vide chapter 5), you will see that they are similar. Therefore, by the property of similar triangles, the corresponding sides of the triangles are proportional.

Thus, we have

$$\frac{AM}{AB} = \frac{AP}{AC} = \frac{MP}{BC}$$

$$\Rightarrow \frac{MP}{AP} = \frac{BC}{AC} = \sin A$$

Similarly,  $\Rightarrow \frac{AM}{AP} = \frac{AB}{AC} = \cos A$


$$\Rightarrow \frac{MP}{AM} = \frac{BC}{AB} = \tan A \text{ and so on.}$$

We thus see that the trigonometric ratios of  $\angle A$  in  $\Delta PAM$  do not differ from those of  $\angle A$  in  $\Delta CAB$ .

From this observation, we can conclude that the values of trigonometric ratios of an angle do not vary with the size of the right triangle considered. In short, the trigonometric ratios of an angle are uniquely defined.

#### 10.4 Motivation of Ratios of $0^\circ$ and $90^\circ$

Let us see what happens to the trigonometric ratios of angle  $A$  if it is made smaller and smaller in the right triangle  $ABC$  (Fig. 10.5), till it becomes zero.



**Fig. 10.5**

Keeping AB fixed, as  $\angle A$  gets smaller and smaller, BC becomes shorter and shorter and finally when  $\angle A$  measures  $0^\circ$ , C coincides with B i.e.  $BC = 0$  and  $AB = AC$ .

This helps to find the values of  $\sin A$  and  $\cos A$  when  $A = 0^\circ$ . In fact, we have

$$\sin 0^\circ = \frac{BC}{AC} = \frac{0}{AC} = 0$$

$$\cos 0^\circ = \frac{AB}{AC} = \frac{AC}{AC} = 1$$


$$\tan 0^\circ = \frac{BC}{AB} = \frac{0}{AB} = 0$$

$$\cot 0^\circ = \frac{AB}{BC} = \frac{AB}{0}, \text{ which is not defined. (why?).}$$

$$\sec 0^\circ = \frac{AC}{AB} = \frac{AC}{AC} = 1$$

$$\text{and } \operatorname{cosec} 0^\circ = \frac{AC}{BC} = \frac{AC}{0}, \text{ which is again not defined.}$$

Next let us see what happens to the trigonometric ratios of  $\angle A$  when it is made larger and larger till it measures  $90^\circ$  (Fig 10.6). Keeping BC fixed as  $\angle A$  gets larger and larger, AB gets shorter and shorter and finally when  $\angle A$  measures  $90^\circ$ , A coincides with B.



**Fig. 10.6**

Thus  $AB = 0$ ,  $AC = BC$ . As such,

$$\sin 90^\circ = \frac{BC}{AC} = \frac{AC}{AC} = 1, \quad \cos 90^\circ = \frac{AB}{AC} = \frac{0}{AC} = 0.$$

Could you find the values of the other trigonometric ratios of  $90^\circ$ ?

### 10.5 Trigonometric Ratios of Some Standard Angles

In this section, we will find the values of trigonometric ratios of angles  $30^\circ$ ,  $45^\circ$  and  $60^\circ$ .

In the previous section you have already known the values of trigonometric ratios of  $0^\circ$  and  $90^\circ$  whenever defined.

**(a) Trigonometric Ratios of  $45^\circ$**

In the right  $\triangle ABC$ , right angled at B, if one angle is  $45^\circ$ , then the other angle is also  $45^\circ$ , i.e. if  $\angle A = 45^\circ$ , then

$$\angle C = 45^\circ \quad (\text{fig 10.7})$$

So,  $AB = BC$  (why)


Suppose,  $AB = BC = k$

Now  $AC^2 = AB^2 + BC^2$  (Pythagoras relation)

$$= k^2 + k^2$$

$$= 2k^2$$

$$\therefore AC = \sqrt{2}k$$



**Fig. 10.7**

Then, using definitions of trigonometric ratios, we have,

$$\sin 45^\circ = \frac{BC}{AC} = \frac{k}{\sqrt{2}k} = \frac{1}{\sqrt{2}}$$

$$\cos 45^\circ = \frac{AB}{AC} = \frac{k}{\sqrt{2}k} = \frac{1}{\sqrt{2}}$$

$$\tan 45^\circ = \frac{BC}{AB} = \frac{k}{k} = 1$$

and  $\cot 45^\circ = \frac{AB}{BC} = \frac{k}{k} = 1$

$$\sec 45^\circ = \frac{AC}{AB} = \sqrt{2},$$

$$\operatorname{cosec} 45^\circ = \frac{AC}{BC} = \sqrt{2}.$$

**(b) Trigonometric Ratios of  $30^\circ$  and  $60^\circ$**

To calculate the values of trigonometric ratios of  $30^\circ$  and  $60^\circ$ , take an equilateral triangle ABC. (Fig 10.8). In an equilateral triangle each angle is  $60^\circ$ . Therefore,

$$\angle A = \angle B = \angle C = 60^\circ.$$

From A draw AD perpendicular to BC meeting it at D.

Now,  $\triangle ABD \cong \triangle ACD$  (why)

$$\therefore BD = DC$$

and  $\angle BAD = \angle CAD$

Consequently,  $BD = DC = \frac{1}{2} BC$

and  $\angle BAD = \angle CAD = \frac{1}{2} \angle A = \frac{1}{2} \times 60^\circ = 30^\circ$

Observe that  $\triangle ABD$  is a right triangle, right angled at D with  $\angle BAD = 30^\circ$  and  $\angle ABD = 60^\circ$  (Fig. 10.8)

Now, let us find the length of sides of  $\triangle ABD$ ,

Suppose  $AB = 2k$ ,

then  $BD = \frac{1}{2} BC = \frac{1}{2} AB = k$

and  $AD^2 = AB^2 - BD^2 = (2k)^2 - k^2 = 3k^2$

$\therefore AD = \sqrt{3}k$

Then, we have,

$$\sin 30^\circ = \frac{BD}{AB} = \frac{k}{2k} = \frac{1}{2}$$

$$\cos 30^\circ = \frac{AD}{AB} = \frac{\sqrt{3}k}{2k} = \frac{\sqrt{3}}{2}$$

$$\tan 30^\circ = \frac{BD}{AD} = \frac{k}{\sqrt{3}k} = \frac{1}{\sqrt{3}}$$

$$\text{Also, cosec } 30^\circ = \frac{AB}{BD} = \frac{2k}{k} = 2$$

$$\sec 30^\circ = \frac{AB}{AD} = \frac{2k}{\sqrt{3}k} = \frac{2}{\sqrt{3}}$$


$$\cot 30^\circ = \frac{AD}{BD} = \frac{\sqrt{3}k}{k} = \sqrt{3}$$

Similarly, from definitions,

$$\sin 60^\circ = \frac{AD}{AB} = \frac{\sqrt{3}k}{2k} = \frac{\sqrt{3}}{2},$$

$$\cos 60^\circ = \frac{BD}{AB} = \frac{k}{2k} = \frac{1}{2}$$

$$\tan 60^\circ = \frac{AD}{BD} = \frac{\sqrt{3}k}{k} = \sqrt{3}$$



**Fig. 10.8**

$$\cot 60^\circ = \frac{BD}{AD} = \frac{k}{\sqrt{3}k} = \frac{1}{\sqrt{3}}$$

$$\sec 60^\circ = \frac{AB}{BD} = \frac{2k}{k} = 2$$

$$\operatorname{cosec} 60^\circ = \frac{AB}{AD} = \frac{2k}{\sqrt{3}k} = \frac{2}{\sqrt{3}}$$

We shall now display the values of all trigonometric ratios of  $0^\circ, 30^\circ, 45^\circ, 60^\circ, 90^\circ$  in a table for ready reference as follows :

$\angle A$	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$
$\sin A$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos A$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan A$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	not defined
$\cot A$	not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0
$\sec A$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	not defined
$\operatorname{cosec} A$	not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1

The use of the values of trigonometric ratios of these angles will be illustrated in the following examples.


**Example 1.** In a right triangle ABC, right angled at C, AB = 10 cm and  $\angle BAC = 30^\circ$  (Fig. 10.9). Find the length of sides BC and AC.

**Solution :** To find the length of the side BC, we will choose the trigonometric ratio involving BC and the given side AB. Since BC is opposite to  $\angle A$  and AB is hypotenuse, we have,

$$\frac{BC}{AB} = \sin A$$

$$\text{i.e. } \frac{BC}{10} = \sin 30^\circ = \frac{1}{2}$$

$$\therefore BC = \frac{10}{2} = 5 \text{ cm}$$



To find the length of AC, we have similarly,

$$\frac{AC}{AB} = \cos A$$

$$\text{i.e. } \frac{AC}{10} = \cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\therefore AC = 10 \times \frac{\sqrt{3}}{2} = 5\sqrt{3} \text{ cm.}$$

**Note :** To find AC, alternatively we could have used the Pythagoras theorem. Thus,

$$AC = \sqrt{AB^2 - BC^2} = \sqrt{10^2 - 5^2} = \sqrt{75}$$

$$= \sqrt{25 \times 3} = 5\sqrt{3} .$$

**Example 2.** If  $\sin(A+B) = \frac{\sqrt{3}}{2}$ ,  $\cos(A-B) = \frac{\sqrt{3}}{2}$ ,  $0^\circ < A+B \leq 90^\circ$ ,  $A > B$ , find A and B.

**Solution :** Given,  $\sin(A+B) = \frac{\sqrt{3}}{2} = \sin 60^\circ$   
 $\therefore A+B = 60^\circ \quad \dots(1)$

And  $\cos(A-B) = \frac{\sqrt{3}}{2} = \cos 30^\circ$   
 $\therefore A-B = 30^\circ \quad \dots(2)$

Solving (1) and (2), we get,

$$A = 45^\circ \text{ and } B = 15^\circ .$$

### EXERCISE 10.1

1. Find value of the following.

(i)  $\sin 30^\circ \cos 45^\circ + \cos 30^\circ \sin 45^\circ$       (ii)  $\frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ}$

(iii)  $\frac{\sin 30^\circ + \tan 45^\circ - \operatorname{cosec} 60^\circ}{\sec 30^\circ + \cos 60^\circ + \cot 45^\circ}$       (iv)  $\frac{5 \cos^2 60^\circ + 4 \sec^2 30^\circ - \tan^2 45^\circ}{\sin^2 30^\circ + \cos^2 30^\circ}$

2. In the right  $\triangle PQR$ , right angled at Q,  $PQ = 4\text{cm}$  and  $\angle PRQ = 30^\circ$ . Find the length of sides PR and QR.

3. If  $\tan(A+B) = \sqrt{3}$  and  $\tan(A-B) = \frac{1}{\sqrt{3}}$ ;  $0^\circ < A+B < 90^\circ$ ,  $A > B$ , find A and B.

### ANSWER

1. (i)  $\frac{1+\sqrt{3}}{2\sqrt{2}}$  (ii)  $\sqrt{3}$  (iii)  $\frac{3\sqrt{3}-4}{4+3\sqrt{3}}$  (iv)  $\frac{67}{12}$   
 2. PR = 8cm, QR =  $4\sqrt{3}$  cm. 3. A =  $45^\circ$ , B =  $15^\circ$ .

### 10.6 Relationships between the Trigonometric Ratios

Let us establish certain relationships that exist between the trigonometric ratios of an angle. Remember the definitions of  $\sin A$  and  $\operatorname{cosec} A$ . In fig 10.10, we have

$$\sin A = \frac{\text{opposite side of } \angle A}{\text{hypotenuse}} = \frac{BC}{AC}$$

$$\operatorname{cosec} A = \frac{\text{hypotenuse}}{\text{opposite sides of } \angle A} = \frac{AC}{BC}$$


So, what do you see ?

$$\operatorname{cosec} A = \frac{AC}{BC} = \frac{1}{\frac{BC}{AC}} = \frac{1}{\sin A} \quad (1)$$

Similarly,

$$\sec A = \frac{AC}{AB} = \frac{1}{\frac{AB}{AC}} = \frac{1}{\cos A} \quad (2)$$

$$\cot A = \frac{AB}{BC} = \frac{1}{\frac{BC}{AB}} = \frac{1}{\tan A} \quad (3)$$



The above three relations (1), (2) and (3) are known as **reciprocal relations** of the trigonometric ratios.

Further, observe that  $\tan A = \frac{BC}{AB}$

$$= \frac{\frac{BC}{AC}}{\frac{AB}{AC}}$$



$$= \frac{\sin A}{\cos A} \dots\dots\dots(4)$$

$$\text{and, } \cot A = \frac{\cos A}{\sin A} \dots\dots\dots(5)$$

These two relations (4) and (5) are known as **quotient relations**.

Now, refer to Fig 10.10. In the  $\triangle ABC$ , right angled at B,

$$\text{we have } AB^2 + BC^2 = AC^2$$

$$\text{Hence, } \frac{AB^2}{AC^2} + \frac{BC^2}{AC^2} = \frac{AC^2}{AC^2}$$

$$\Rightarrow \left(\frac{AB}{AC}\right)^2 + \left(\frac{BC}{AC}\right)^2 = 1$$

$$\Rightarrow (\cos A)^2 + (\sin A)^2 = 1$$

$$\Rightarrow \cos^2 A + \sin^2 A = 1 \dots\dots\dots(6)$$

This relation holds for every A such that  $0 \leq A \leq 90^\circ$ .

$$\text{Again, } AB^2 + BC^2 = AC^2$$

$$\Rightarrow \frac{AB^2}{AB^2} + \frac{BC^2}{AB^2} = \frac{AC^2}{AB^2}$$

$$\Rightarrow \left(\frac{AB}{AB}\right)^2 + \left(\frac{BC}{AB}\right)^2 = \left(\frac{AC}{AB}\right)^2$$

$$1 + \tan^2 A = \sec^2 A \dots\dots\dots(7)$$

Is this equation true for  $A = 0$ ? Yes, it is. What about  $A = 90^\circ$ ? We have seen that  $\tan A$  and  $\sec A$  are not defined when  $A = 90^\circ$ . So (7) is true for all A such that  $0^\circ \leq A < 90^\circ$ .

Again,

$$\frac{AB^2}{BC^2} + \frac{BC^2}{BC^2} = \frac{AC^2}{BC^2}$$

$$\Rightarrow \left(\frac{AB}{BC}\right)^2 + \left(\frac{BC}{BC}\right)^2 = \left(\frac{AC}{BC}\right)^2$$

$$\Rightarrow \cot^2 A + 1 = \operatorname{cosec}^2 A \dots\dots\dots(8)$$

Remember that  $\cot A$  and  $\operatorname{cosec} A$  are not defined for  $A = 0^\circ$ . Therefore (8) is true for all A such that  $0^\circ < A \leq 90^\circ$ .

The above three relations (6), (7), and (8) are known as **Pythagorean relations of trigonometric ratios**.

From (6), (7) and (8) we see that

$$\begin{aligned}\cos^2 A &= 1 - \sin^2 A, & \sin^2 A &= 1 - \cos^2 A, \\ \sec^2 A - \tan^2 A &= 1, & \tan^2 A &= \sec^2 A - 1. \\ \operatorname{cosec}^2 A - \cot^2 A &= 1, & \cot^2 A &= \operatorname{cosec}^2 A - 1\end{aligned}$$

**Note :** For the sake of convenience , we write  $\cos^2 A$ ,  $\sin^2 A$  etc. in place of  $(\cos A)^2$ ,  $(\sin A)^2$  etc. But  $\cot A = (\tan A)^{-1} \neq \tan^{-1} A$  (it is called tangent inverse A).

The symbol  $\tan^{-1} A$  has a different meaning, which will be discussed in higher classes. Similar conventions hold for other trigonometric ratios as well.

### 10.7 Trigonometric Identities

Let us recall that an equation is called an identity if it is true for all values of the variables involved. Thus, an equation involving trigonometric ratios of an angle ( this being a variable) is called a trigonometric identity if it is true for all admissible values of the angle involved.

All the trigonometric relations established in the previous section are standard trigonometric identities.

Further consider the following relation:

$$\cos^2 A - \cos A = \cos A(\cos A - 1).$$

This is an identity because it is true for all values of A;

whereas  $\cos^2 A - \cos A = 0$  is not an identity, since it is satisfied only for some particular values of A (in fact for  $A = 0^\circ$  and  $A = 90^\circ$  only).

Using the standard identities we can express each trigonometric ratio in terms of a given ratio. Of course, from definition of trigonometric ratios we can do the same. The process is illustrated in the following examples.

**Example 3.** Given  $\sin A = \frac{3}{4}$ , find the other trigonometric ratios of A.

**Solution :** **First method:**

we know that  $\cos^2 A + \sin^2 A = 1$

$$\therefore \cos^2 A + \left(\frac{3}{4}\right)^2 = 1$$

$$\begin{aligned}\Rightarrow \cos^2 A &= 1 - \left(\frac{3}{4}\right)^2 \\ &= 1 - \frac{9}{16} \\ &= \frac{16-9}{16} \\ &= \frac{7}{16} \\ \therefore \cos A &= \frac{\sqrt{7}}{4} \quad (\text{neglecting negative value})\end{aligned}$$

$$\text{Then, } \tan A = \frac{\sin A}{\cos A} = \frac{\frac{3}{4}}{\frac{\sqrt{7}}{4}} = \frac{3}{\sqrt{7}}$$

$$\cot A = \frac{1}{\tan A} = \frac{\sqrt{7}}{3}$$

$$\sec A = \frac{1}{\cos A} = \frac{4}{\sqrt{7}}$$

$$\text{and } \operatorname{cosec} A = \frac{1}{\sin A} = \frac{4}{3}.$$

**Second method:** Draw a right  $\triangle ABC$  right angled at B (Fig. 10.11)

$$\text{we know that } \sin A = \frac{BC}{AC} = \frac{3}{4}.$$

Therefore, if  $BC = 3k$ , then  $AC = 4k$ , where  $k$  is a positive number.

We also know that  $AC^2 = AB^2 + BC^2$


$$\therefore (4k)^2 = AB^2 + (3k)^2$$

$$\Rightarrow 16k^2 = AB^2 + 9k^2$$

$$\Rightarrow AB^2 = 16k^2 - 9k^2$$

$$= 7k^2$$

$$\therefore AB = \sqrt{7}k.$$



**Fig. 10.11**

Now, by definitions,

$$\cos A = \frac{AB}{AC} = \frac{\sqrt{7}k}{4k} = \frac{\sqrt{7}}{4}$$

$$\tan A = \frac{BC}{AB} = \frac{3k}{\sqrt{7}k} = \frac{3}{\sqrt{7}},$$

$$\cot A = \frac{AB}{BC} = \frac{\sqrt{7}k}{3k} = \frac{\sqrt{7}}{3},$$

$$\sec A = \frac{AC}{AB} = \frac{4k}{\sqrt{7}k} = \frac{4}{\sqrt{7}}$$

$$\operatorname{cosec} A = \frac{AC}{BC} = \frac{4k}{3k} = \frac{4}{3}.$$

**Example 4.** In a right triangle ABC, right angled at C, if  $\tan A = \frac{1}{\sqrt{3}}$ , find the value of  $\cos^2 A - \sin^2 A$ .

**Solution :** In  $\triangle ABC$ ,  $\tan A = \frac{1}{\sqrt{3}}$  (Fig. 10.12)

$$\Rightarrow \frac{BC}{AC} = \frac{1}{\sqrt{3}}.$$

$$\sqrt{3}BC = AC \dots\dots\dots(1)$$

Let  $BC = k$ , then  $AC = \sqrt{3}k$

Now,  $AB^2 = AC^2 + BC^2$

$$= (\sqrt{3}k)^2 + k^2$$

$$= 3k^2 + k^2$$


$$= 4k^2$$

$$\therefore AB = \sqrt{4k^2} = 2k$$

$$\text{Then } \cos A = \frac{AC}{AB} = \frac{\sqrt{3}k}{2k} = \frac{\sqrt{3}}{2}$$

$$\text{and } \sin A = \frac{BC}{AB} = \frac{k}{2k} = \frac{1}{2}$$

$$\therefore \cos^2 A - \sin^2 A = \left(\frac{\sqrt{3}}{2}\right)^2 - \left(\frac{1}{2}\right)^2 = \frac{3}{4} - \frac{1}{4} = \frac{2}{4} = \frac{1}{2}.$$



**Fig. 10.12**

**Example 5.** In a right triangle DEF, right angled at E, EF = 24 cm. DE = 7 cm. Find sin F and cos F.

**Solution :** In  $\triangle DEF$ , we have,

$$\begin{aligned} DF^2 &= DE^2 + EF^2 \\ &= 7^2 + 24^2 \\ &= 49 + 576 \\ &= 625 \end{aligned}$$

$$\therefore DF = \sqrt{625} = 25 \text{ cm}$$

$$\text{Then, } \sin F = \frac{DE}{DF} = \frac{7}{25}$$

$$\cos F = \frac{EF}{DF} = \frac{24}{25}$$




Fig. 10.13

**Example 6.** Express tan A, sec A and sin A in terms of cot A.

**Solution :** We know that  $\cot A = \frac{1}{\tan A}$

$$\therefore \tan A = \frac{1}{\cot A}$$

Next,

$$\begin{aligned} \sec^2 A &= 1 + \tan^2 A = 1 + \left(\frac{1}{\cot A}\right)^2 \\ &= 1 + \frac{1}{\cot^2 A} = \frac{\cot^2 A + 1}{\cot^2 A} \end{aligned}$$

$$\therefore \sec A = \pm \sqrt{\frac{\cot^2 A + 1}{\cot^2 A}}$$

This gives,  $\sec A = \frac{\sqrt{\cot^2 A + 1}}{\cot A}$  (why)

Further,  $\operatorname{cosec}^2 A = 1 + \cot^2 A$

$$\Rightarrow \frac{1}{\sin^2 A} = 1 + \cot^2 A$$

$$\Rightarrow \sin^2 A = \frac{1}{1 + \cot^2 A}$$

$$\therefore \sin A = \frac{1}{\sqrt{1 + \cot^2 A}}$$

**Example 7.** Determine whether the following is an identity :

$$\tan^2 \theta + \cot^2 \theta = 2$$

**Solution :**  $\tan^2 \theta + \cot^2 \theta = 2$

$$\Rightarrow \tan^2 \theta + \frac{1}{\tan^2 \theta} = 2$$

$$\Rightarrow \tan^4 \theta + 1 = 2 \tan^2 \theta$$

$$\Rightarrow \tan^4 \theta + 1 - 2 \tan^2 \theta = 0$$

$$\Rightarrow (\tan^2 \theta - 1)^2 = 0$$

$$\Rightarrow \tan^2 \theta - 1 = 0$$

$$\Rightarrow \tan^2 \theta = 1$$

This relation cannot hold for all admissible values of  $\theta$ . For instance the relation does not hold when  $\theta = 60^\circ$ . Hence it is not an identity

**Second method :** If it can be shown that the equality does not hold for some particular value of  $\theta$ , then the relation is not an identity.

Taking  $\theta = 30^\circ$ , we have,

$$\text{L.H.S} = \tan^2 30^\circ + \cot^2 30^\circ$$

$$= \left( \frac{1}{\sqrt{3}} \right)^2 + (\sqrt{3})^2$$

$$= \frac{1}{3} + 3 = \frac{1+9}{3}$$

$$= \frac{10}{3} \neq 2 \quad (\text{R.H. S})$$

$\therefore$  the relation is not an identity

**Example 8.** Prove the following identities

$$(i) \frac{1}{\operatorname{cosec} \theta - \cot \theta} - \frac{1}{\sin \theta} = \frac{1}{\sin \theta} - \frac{1}{\operatorname{cosec} \theta + \cot \theta}$$

$$(ii) \sin^6 \theta + \cos^6 \theta = 1 - 3 \sin^2 \theta \cos^2 \theta$$

$$(iii) \frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} = \sec \theta \operatorname{cosec} \theta + 1$$

**Solution :** (i)  $\frac{1}{\operatorname{cosec}\theta - \cot\theta} - \frac{1}{\sin\theta} = \frac{1}{\sin\theta} - \frac{1}{\operatorname{cosec}\theta + \cot\theta}$

i.e.  $\frac{1}{\operatorname{cosec}\theta - \cot\theta} + \frac{1}{\operatorname{cosec}\theta + \cot\theta} = \frac{1}{\sin\theta} + \frac{1}{\sin\theta}$

i.e.  $\frac{1}{\operatorname{cosec}\theta - \cot\theta} + \frac{1}{\operatorname{cosec}\theta + \cot\theta} = \frac{2}{\sin\theta} \dots\dots\dots(1)$

Now, L.H.S. of (1) =  $\frac{1}{\operatorname{cosec}\theta - \cot\theta} + \frac{1}{\operatorname{cosec}\theta + \cot\theta}$   
 $= \frac{\operatorname{cosec}\theta + \cot\theta + \operatorname{cosec}\theta - \cot\theta}{\operatorname{cosec}^2\theta - \cot^2\theta}$   
 $= \frac{2\operatorname{cosec}\theta}{1} \quad (\because \operatorname{cosec}^2\theta = 1 + \cot^2\theta)$   
 $= \frac{2}{\sin\theta} = \text{R.H.S of (1)}$

This establishes the given identity.

**Alternative method :**

L.H.S. =  $\frac{\operatorname{cosec}\theta + \cot\theta}{(\operatorname{cosec}\theta - \cot\theta)(\operatorname{cosec}\theta + \cot\theta)} - \operatorname{cosec}\theta$   
 $= \frac{\operatorname{cosec}\theta + \cot\theta}{\operatorname{cosec}^2\theta - \cot^2\theta} - \operatorname{cosec}\theta$   
 $= \operatorname{cosec}\theta + \cot\theta - \operatorname{cosec}\theta$   
 $= \operatorname{cosec}\theta + \frac{(\cot\theta - \operatorname{cosec}\theta)(\cot\theta + \operatorname{cosec}\theta)}{(\cot\theta + \operatorname{cosec}\theta)}$   
 $= \operatorname{cosec}\theta + \frac{\cot^2\theta - \operatorname{cosec}^2\theta}{\cot\theta + \operatorname{cosec}\theta}$   
 $= \operatorname{cosec}\theta - \frac{\operatorname{cosec}^2\theta - \cot^2\theta}{\operatorname{cosec}\theta - \cot\theta}$   
 $= \operatorname{cosec}\theta - \frac{1}{\operatorname{cosec}\theta + \cot\theta}$   
 $= \frac{1}{\sin\theta} - \frac{1}{\operatorname{cosec}\theta + \cot\theta} = \text{R.H.S}$

$$\begin{aligned}
 \text{(ii) } \sin^6 \theta + \cos^6 \theta &= (\sin^2 \theta)^3 + (\cos^2 \theta)^3 \\
 &= (\sin^2 \theta + \cos^2 \theta)^3 - 3 \sin^2 \theta \cos^2 \theta (\sin^2 \theta + \cos^2 \theta) \\
 &= 1^3 - 3 \sin^2 \theta \cos^2 \theta \\
 &= 1 - 3 \sin^2 \theta \cos^2 \theta.
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) L.H.S.} &= \frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} \\
 &= \frac{\frac{\sin \theta}{\cos \theta}}{1 - \frac{\cos \theta}{\sin \theta}} + \frac{\frac{\cos \theta}{\sin \theta}}{1 - \frac{\sin \theta}{\cos \theta}} \\
 &= \frac{\frac{\sin \theta}{\cos \theta}}{\frac{\sin \theta - \cos \theta}{\sin \theta}} + \frac{\frac{\cos \theta}{\sin \theta}}{\frac{\cos \theta - \sin \theta}{\cos \theta}} \\
 &= \frac{\sin^2 \theta}{\cos \theta (\sin \theta - \cos \theta)} + \frac{\cos^2 \theta}{\sin \theta (\cos \theta - \sin \theta)} \\
 &= \frac{\sin^2 \theta}{\cos \theta (\sin \theta - \cos \theta)} - \frac{\cos^2 \theta}{\sin \theta (\sin \theta - \cos \theta)} \\
 &= \frac{\sin^3 \theta - \cos^3 \theta}{\cos \theta \sin \theta (\sin \theta - \cos \theta)} \\
 &= \frac{(\sin \theta - \cos \theta)(\sin^2 \theta + \cos \theta \sin \theta + \cos^2 \theta)}{\cos \theta \sin \theta (\sin \theta - \cos \theta)} \\
 &= \frac{(\sin^2 \theta + \cos^2 \theta + \sin \theta \cos \theta)}{\cos \theta \sin \theta} \\
 &= \frac{1 + \sin \theta \cos \theta}{\cos \theta \sin \theta} \\
 &= \frac{1}{\cos \theta \sin \theta} + 1 \\
 &= \sec \theta \operatorname{cosec} \theta + 1 \qquad \dots \dots \dots \text{(ii)} \\
 &= \text{R.H.S.}
 \end{aligned}$$



## EXERCISE 10.2

1. In a right  $\triangle ABC$ , right angled at C,  $AC = 3\text{cm}$ ,  $AB = 5\text{cm}$ .  
Find the value of (i)  $\sin A$ ,  $\cos A$  (ii)  $\sin B$ ,  $\cos B$
2. In a right  $\triangle DEF$ , right angled at E,  $DF = 13\text{ cm}$ ,  $DE = 12\text{ cm}$ , find  
(i)  $\tan D - \cot F$   
(ii)  $\tan D - \cot D$
3. If  $\cos A = \frac{3}{5}$ , calculate  $\sin A$  and  $\tan A$ .
4. If  $\tan A = \frac{3}{4}$ , find  $\cos A$  and  $\operatorname{cosec} A$ .
5. If  $\sec A = \frac{13}{12}$ , find  $\tan A$  and  $\sin A$ .
6. If  $\cot A = \frac{4}{3}$ , find the value of  $\cos^2 A - \sin^2 A$
7. If  $\tan \theta = \frac{a}{b}$ , show that  $\frac{a \sin \theta - b \cos \theta}{a \sin \theta + b \cos \theta} = \frac{a^2 - b^2}{a^2 + b^2}$ .
8. In a right  $\triangle PQR$ , right angled at Q,  $PR - PQ = 1\text{cm}$  and  $QR = 5\text{cm}$ . Find the values of  $\sin R$ ,  $\cos R$  and  $\tan R$ .
9. Express  $\cos \theta$ ,  $\tan \theta$  and  $\sec \theta$  in terms of  $\sin \theta$ .
10. Determine whether the following equations are identities :  
(i)  $\cot^2 \theta + \cos \theta = \sin^2 \theta$  (ii)  $\frac{\tan \theta + \sin \theta}{\tan \theta - \sin \theta} = \frac{\sec \theta + 1}{\sec \theta - 1}$   
(iii)  $\tan^2 \theta - \sin^2 \theta = \tan^2 \theta \sin^2 \theta$   
(iv)  $\tan \phi \operatorname{cosec} \phi = \sec \phi$
11. Prove the following identities :  
(i)  $\frac{1 + \cos \theta}{1 - \cos \theta} = (\operatorname{cosec} \theta + \cot \theta)^2$   
(ii)  $\frac{1 + \cos A}{\sin A} = \frac{\sin A}{1 - \cos A}$

$$(iii) \frac{\cos A}{1 + \sin A} + \frac{1 + \sin A}{\cos A} = 2 \sec A$$

$$(iv) \sec^2 \theta + \operatorname{cosec}^2 \theta = \sec^2 \theta \operatorname{cosec}^2 \theta$$

$$(v) \frac{1 + \sin \theta}{1 - \sin \theta} = (\sec \theta + \tan \theta)^2$$

$$(vi) \frac{1 + \cos \theta - \sin^2 \theta}{\sin \theta (1 + \cos \theta)} = \cot \theta$$

$$(vii) \frac{\cos \theta}{\operatorname{cosec} \theta + 1} + \frac{\cos \theta}{\operatorname{cosec} \theta - 1} = 2 \tan \theta$$

$$(viii) \cos^2 \theta - \sin^2 \theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

$$(ix) \sin^4 \theta - \cos^4 \theta = \sin^2 \theta - \cos^2 \theta$$

$$(x) \left(1 + \frac{1}{\tan^2 A}\right) \left(1 + \frac{1}{\cot^2 A}\right) = \frac{1}{\sin^2 A - \sin^4 A}$$

$$(xi) \tan^2 A + \cot^2 A + 2 = \sec^2 A \operatorname{cosec}^2 A$$

$$(xii) \frac{\sin^3 A + \cos^3 A}{\sin A + \cos A} = 1 - \sin A \cos A$$

$$(xiii) \frac{1 + \cos \theta + \sin \theta}{1 + \cos \theta - \sin \theta} = \frac{1 + \sin \theta}{\cos \theta}$$

$$(xiv) \frac{\cos A}{1 - \tan A} + \frac{\sin A}{1 - \cot A} = \sin A + \cos A$$

$$(xv) \sin A(1 + \tan A) + \cos A(1 + \cot A) = \sec A + \operatorname{cosec} A$$

$$(xvi) \sin A(\sec A + \operatorname{cosec} A) - \cos A(\sec A - \operatorname{cosec} A) = \sec A \operatorname{cosec} A .$$

### ANSWER

$$1. \quad (i) \sin A = \frac{4}{5}, \cos A = \frac{3}{5}, \quad (ii) \sin B = \frac{3}{5}, \cos B = \frac{4}{5}.$$

$$2. \quad (i) 0 \quad (ii) \frac{-119}{60}.$$

$$3. \quad \sin A = \frac{4}{5}, \quad \tan A = \frac{4}{3}$$

$$4. \quad \cos A = \frac{4}{5}, \quad \operatorname{cosec} A = \frac{5}{3}$$

$$5. \quad \tan A = \frac{5}{12}, \quad \sin A = \frac{5}{13}$$

$$6. \quad \frac{7}{25}$$

$$8. \quad \sin R = \frac{12}{13}, \quad \cos R = \frac{5}{13}, \quad \tan R = \frac{12}{5}$$

$$9. \quad \cos \theta = \sqrt{1 - \sin^2 \theta}, \quad \tan \theta = \frac{\sin \theta}{\sqrt{1 - \sin^2 \theta}}, \quad \sec \theta = \frac{1}{\sqrt{1 - \sin^2 \theta}}$$

### 10.8 Trigonometric Ratios of Complementary Angles

We know that in any right triangle, there are two acute angles. What is the relation between them? They are complementary i.e. their sum is  $90^\circ$ .

Fig. 10.14 shows a right triangle ABC, whose acute angles are  $\angle A$  and  $\angle C$ .


Hence,  $\angle A + \angle C = 90^\circ$  i.e. they are complementary.

We have,

$$\left. \begin{aligned} \sin A &= \frac{BC}{AC}, \quad \cos A = \frac{AB}{AC}, \quad \tan A = \frac{BC}{AB} \\ \operatorname{cosec} A &= \frac{AC}{BC}, \quad \sec A = \frac{AC}{AB}, \quad \cot A = \frac{AB}{BC} \end{aligned} \right\} \dots\dots(i)$$

Let us find the trigonometric ratios of  $C = 90^\circ - A$ .

First let us check what should be the side opposite to and adjacent to angle  $90^\circ - A$ . You will find that AB is the opposite side and BC, the adjacent side for the angle C i.e.  $90^\circ - A$ .



$$\therefore \left. \begin{aligned} \sin(90^\circ - A) &= \frac{AB}{AC}, \cos(90^\circ - A) = \frac{BC}{AC}, \tan(90^\circ - A) = \frac{AB}{BC}, \\ \operatorname{cosec}(90^\circ - A) &= \frac{AC}{AB}, \sec(90^\circ - A) = \frac{AC}{BC}, \cot(90^\circ - A) = \frac{BC}{AB} \end{aligned} \right\} \dots(ii)$$

Let us now compare the ratios in (i) and (ii). We will find that

$$\sin(90^\circ - A) = \frac{AB}{AC} = \cos A \text{ and } \cos(90^\circ - A) = \frac{BC}{AC} = \sin A,$$

$$\tan(90^\circ - A) = \frac{AB}{BC} = \cot A \text{ and } \cot(90^\circ - A) = \frac{BC}{AB} = \tan A$$

$$\sec(90^\circ - A) = \frac{AC}{BC} = \operatorname{cosec} A, \text{ and } \operatorname{cosec}(90^\circ - A) = \frac{AC}{AB} = \sec A.$$

$$\text{Thus, } \sin(90^\circ - A) = \cos A, \quad \cos(90^\circ - A) = \sin A$$

$$\tan(90^\circ - A) = \cot A, \quad \cot(90^\circ - A) = \tan A$$

$$\sec(90^\circ - A) = \operatorname{cosec} A, \quad \operatorname{cosec}(90^\circ - A) = \sec A,$$

for all values of  $A$  lying between  $0^\circ$  and  $90^\circ$ .

It would be a good exercise to check whether these hold for  $A = 0^\circ$  and  $A = 90^\circ$ .

**Example 9.** Find the value of  $\frac{\sin 18^\circ}{\cos 72^\circ}$

**Solution :** We know that  $\cos A = \sin(90^\circ - A)$

$$\therefore \cos 72^\circ = \sin(90^\circ - 72^\circ) = \sin 18^\circ$$

$$\text{Hence, } \frac{\sin 18^\circ}{\cos 72^\circ} = \frac{\sin 18^\circ}{\sin 18^\circ} = 1.$$

**Example 10.** If  $\tan 2A = \cot(A - 18^\circ)$ , where  $2A$  is an acute angle, find the value of  $A$ .

**Solution :** We know that

$$\tan 2A = \cot(90^\circ - 2A)$$

Hence, the given relation can be written as

$$\cot(90^\circ - 2A) = \cot(A - 18^\circ)$$

$$\therefore 90^\circ - 2A = A - 18^\circ. \text{ [Since } 90^\circ - 2A \text{ and } A - 18^\circ \text{ are acute angles.]}$$

$$\begin{aligned} \text{Then, } 3A &= 90^\circ + 18^\circ \\ &= 108^\circ \\ \therefore A &= \frac{108^\circ}{3} = 36^\circ. \end{aligned}$$

**Example 11.** Express  $\sin 67^\circ + \cos 65^\circ$  in terms of trigonometric ratios of angles between  $0^\circ$  and  $45^\circ$ .

**Solution :**

$$\begin{aligned} \sin 67^\circ + \cos 65^\circ &= \cos(90^\circ - 67^\circ) + \sin(90^\circ - 65^\circ) \\ &= \cos 23^\circ + \sin 25^\circ. \end{aligned}$$

**Example 12.** In the triangle ABC and DEF,  $\angle A = 50^\circ$ ,  $\angle B = 90^\circ$  and  $AC = 4\text{cm}$ ,  $\angle D = 40^\circ$ ,  $\angle E = 90^\circ$  and  $DE = 4\text{cm}$ .

Prove that  $\sqrt{\frac{\text{area of } \triangle ABC}{\text{area of } \triangle DEF}} = \sin 50^\circ$ .

**Solution :** In the Fig. 10.15. (i),  $\frac{AB}{AC} = \cos 50^\circ$

$$\begin{aligned} \Rightarrow AB &= AC \cos 50^\circ \\ &= 4 \cos 50^\circ \text{ cm.} \end{aligned}$$


and,  $\frac{BC}{AC} = \sin 50^\circ$

$$\begin{aligned} \Rightarrow BC &= AC \sin 50^\circ \\ &= 4 \sin 50^\circ \text{ cm} \end{aligned}$$

$$\begin{aligned} \therefore \text{Area of } \triangle ABC &= \frac{1}{2} AB \cdot BC \\ &= \frac{1}{2} \times 4 \cos 50^\circ \times 4 \sin 50^\circ \text{ cm}^2 \\ &= 8 \cos 50^\circ \sin 50^\circ \text{ cm}^2 \end{aligned}$$

In Fig 10.15 (ii),  $\frac{EF}{DE} = \tan 40^\circ$

$$\begin{aligned} \Rightarrow EF &= DE \tan 40^\circ \text{ cm.} \\ &= 4 \tan 40^\circ \text{ cm.} \end{aligned}$$



**Fig. 10.15**

$$\begin{aligned}\therefore \text{Area of } \triangle DEF &= \frac{1}{2} DE \cdot EF \\ &= \frac{1}{2} 4.4 \tan 40^\circ \text{ cm}^2 \\ &= 8 \tan 40^\circ \text{ cm}^2\end{aligned}$$

$$\begin{aligned}\text{Now, } \frac{\text{area of } \triangle ABC}{\text{area of } \triangle DEF} &= \frac{8 \cos 50^\circ \sin 50^\circ}{8 \tan 40^\circ} \\ &= \frac{\cos 50^\circ \sin 50^\circ}{\frac{\sin 40^\circ}{\cos 40^\circ}} \\ &= \frac{\cos 40^\circ \cos 50^\circ \sin 50^\circ}{\sin 40^\circ} \\ &= \frac{\sin(90^\circ - 40^\circ) \sin(90^\circ - 50^\circ) \sin 50^\circ}{\sin 40^\circ} \\ &= \frac{\sin 50^\circ \sin 40^\circ \sin 50^\circ}{\sin 40^\circ} \\ &= \sin^2 50^\circ\end{aligned}$$

$$\therefore \sqrt{\frac{\text{area of } \triangle ABC}{\text{area of } \triangle DEF}} = \sin 50^\circ$$

### EXERCISE 10.3

1. Find the value of

$$(i) \frac{\tan 49^\circ}{\cot 41^\circ} \quad (ii) \frac{\cos 70^\circ}{\sin 20^\circ} \quad (iii) \frac{\tan 26^\circ}{\cot 64^\circ}$$

$$(iv) \frac{\operatorname{cosec} 35^\circ}{\sec 55^\circ} \quad (v) \cos 37^\circ - \sin 53^\circ$$

$$(vi) \operatorname{cosec} 47^\circ - \sec 43^\circ$$

2. Show that

$$(i) \sec 53^\circ \sin 53^\circ \tan 37^\circ = 1$$

$$(ii) \tan 48^\circ \tan 23^\circ \tan 42^\circ \tan 67^\circ = 1$$

$$(iii) \sin 40^\circ \sec 40^\circ \sec 50^\circ = \operatorname{cosec} 50^\circ$$

$$(iv) \cos 38^\circ \cos 52^\circ - \sin 38^\circ \sin 52^\circ = 0$$

3. If  $\tan A = \cot B$ , show that  $A + B = 90^\circ$ .
4. If  $\sec 4A = \operatorname{cosec}(A - 20^\circ)$ , where  $4A$  is an acute angle, find the value of  $A$ .
5. Find the value of  $\theta$ , where all the angles are acute or equal to  $90^\circ$  :
  - (i)  $\operatorname{cosec} 4\theta = \sec 6\theta$
  - (ii)  $\tan(2\theta + 35^\circ) = \cot 3\theta$
  - (iii)  $\sin(\theta + 55^\circ) = \cos(7\theta - 5^\circ)$
6. Express  $\cot 85^\circ + \cos 70^\circ$  in terms of trigonometric ratios of angles between  $0^\circ$  and  $45^\circ$ .
7. If  $A, B, C$  are angles of a triangle  $ABC$ , show that  $\sin\left(\frac{B+C}{2}\right) = \cos\frac{A}{2}$ .
8. Compare the areas of the right triangles  $ABC$  and  $DEF$  in which  $\angle A = 40^\circ$ ,  $\angle B = 90^\circ$  and  $AC = 5\text{cm}$ ;  $\angle D = 50^\circ$ ,  $\angle E = 90^\circ$  and  $DE = 5\text{cm}$ .

### ANSWER


1. (i) 1 (ii) 1 (iii) 1 (iv) 1 (v) 0 (vi) 0
4.  $A = 22^\circ$ .
5. (i)  $\theta = 9^\circ$  (ii)  $\theta = 11^\circ$  (iii)  $\theta = 5^\circ$ .
6.  $\tan 5^\circ + \sin 20^\circ$
8.  $\frac{\operatorname{ar}(\triangle ABC)}{\operatorname{ar}(\triangle DEF)} = \sin^2 40^\circ$  (or  $\cos^2 50^\circ$ )

### 10.9 HEIGHTS AND DISTANCES

In this section, we shall see how trigonometry is applied in finding the heights and distances of various objects without actually measuring them. While dealing with such problems we will come across two terms known as the **angle of elevation** and the **angle of depression**.


**Angle of Elevation :**

Suppose you are looking up at an object B, say a Kite, (Fig 10.6) A being the position of your eye. Then AB, the line drawn from eye to the object viewed, is the line of sight. It makes an angle BAC with the horizontal line AC through your eye. This  $\angle BAC$  is called the angle of elevation of B from A. Thus the angle of elevation of a point observed is the angle formed by the line of sight with the horizontal, when the point being observed is above the horizontal through the eye.

**Angle of Depression :**

Suppose you are sitting on a balcony of a house and are looking down at a football B on the ground. (Fig 10.17), A being the position of your eye. AB is the line of sight and it makes an angle BAC with AC, the horizontal through A.

This  $\angle BAC$  is called the angle of depression of B from A. Thus, the angle of depression of a point observed is the angle formed by the line of sight with the horizontal when the point observed is below the horizontal through the eye.



After knowing these terms, we illustrate the process of using trigonometry in finding heights and distances with the help of some examples.

**Example 13.** A pole stands vertically on the ground. From a point on the ground, 15m away from the foot of the pole, the angle of elevation of the top of the pole is  $60^\circ$ . Find the height of the pole.



**Solution :** First let us draw a simple representative diagram of the problem.

Let BC represent the pole and let A be the point of observation.


Then, AC is the line of sight.

Hence  $\angle BAC = 60^\circ$ , and  $\angle ABC = 90^\circ$  (why?) and  $AB = 15\text{m}$ . Think of a trigonometric ratio involving BC and the known side AB.

$$\text{Clearly, } \frac{BC}{AB} = \tan A = \tan 60^\circ = \sqrt{3}$$

$$\therefore BC = AB\sqrt{3} = 15\sqrt{3}.$$

i.e. height of the pole =  $15\sqrt{3}$  m.



**Fig. 10.18**

**Example 14.** Find the angle of elevation of the top of a tower of height  $100\sqrt{3}$  m from a point on the ground, 100m away from its foot.

**Solution :** Let BC be the tower and A be the point of observation.

Then,  $BC = 100\sqrt{3}$ ,  
 $\angle BAC = \theta$ , the angle of elevation  
 and  $AB = 100$  m.

In right triangle ABC,


$$\frac{BC}{AB} = \tan \theta$$

$$\Rightarrow \frac{100\sqrt{3}}{100} = \tan \theta$$

$$\text{i.e. } \tan \theta = \sqrt{3}$$

$$\Rightarrow \theta = 60^\circ$$

$\therefore$  the required angle of elevation is  $60^\circ$ .



**Fig. 10.19**

**Example 15.** Find the height of a tree, if its shadow is 10m long when the altitude of the sun is  $30^\circ$ .


**Solution :** Let BC represent the tree and AB its shadow. In the right  $\triangle ABC$  (right angled at B) we have  $\angle A = 30^\circ$  (Fig. 10.20) and  $AB = 10\text{m}$ .

$$\text{Now, } \frac{BC}{AB} = \tan 30^\circ$$

$$\Rightarrow BC = AB \frac{1}{\sqrt{3}}$$

$$\Rightarrow BC = \frac{10\sqrt{3}}{3} \text{ m}$$

$$\therefore \text{the height of the tree is } \frac{10\sqrt{3}}{3} \text{ m.}$$



**Example 16.** The shadow of a tower, standing on level ground, is found to be 45m longer when the sun's altitude is  $30^\circ$  than when it was at  $60^\circ$ . Find the height of the tower.

**Solution :** Let AB be the tower and BC, BD be the shadows when the altitude of the sun are  $60^\circ$  and  $30^\circ$  respectively,  
Then  $\angle BCA = 60^\circ$  and  $\angle BDA = 30^\circ$  and  $CD = 45$   
Let  $BC = x$  m.

$$\text{In } \triangle ABC, \tan C = \frac{AB}{BC}$$

$$\begin{aligned} \text{i.e. } AB &= BC \tan 60^\circ \\ &= BC \sqrt{3} \\ &= \sqrt{3}x \quad \dots(i) \end{aligned}$$

$$\text{In } \triangle ABD, \tan D = \frac{AB}{BD}$$

$$\Rightarrow \tan 30^\circ = \frac{AB}{BC+CD}$$


$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AB}{x+45}$$

$$\begin{aligned} \Rightarrow x+45 &= \sqrt{3}AB \\ &= \sqrt{3}x \cdot \sqrt{3}x \quad \text{by (i)} \\ &= 3x \end{aligned}$$

$$\Rightarrow 45 = 2x.$$

$$\therefore x = \frac{45}{2} = 22.5 \text{ m}$$

$$\therefore AB = \text{height of the tower} = \sqrt{3}x = \sqrt{3} \times 22.5 \text{ m}$$



**Example 17.** The angles of depression of the top and the bottom of a 7m tall tree from the top of a tower are  $45^\circ$  and  $60^\circ$  respectively. Find the height of the tower.

**Solution :** Let AB represent the tower and CD the tree .  
Draw DE parallel to CB.  
Then

$$\angle XAD = 45^\circ$$

$$\text{and } \angle XAC = 60^\circ \quad (\text{Fig 10.22})$$

$$\therefore \angle ADE = 45^\circ \text{ and } \angle ACB = 60^\circ$$

$$\text{In right } \triangle ABC, \quad \frac{AB}{BC} = \tan 60^\circ$$

$$\Rightarrow AE + EB = BC \cdot \sqrt{3}$$

$$\Rightarrow h + DC = BC \cdot \sqrt{3}$$

$$(\because EB = DC \text{ and if } AE = h)$$

$$\Rightarrow h + 7 = \sqrt{3}BC \dots (1)$$

$$\text{In right } \triangle ADE, \quad \frac{AE}{ED} = \tan 45^\circ = 1$$

$$\Rightarrow AE = ED$$

$$\Rightarrow h = BC \dots (2)$$

Using (2) in (1)

$$h + 7 = \sqrt{3}h$$

$$\Rightarrow \sqrt{3}h - h = 7$$

$$\Rightarrow (\sqrt{3} - 1)h = 7$$


$$\therefore h = \frac{7}{\sqrt{3} - 1} = \frac{7(\sqrt{3} + 1)}{2}$$

$$\therefore AB = \text{height of the tower} = AE + EB = AE + DC$$

$$= h + 7$$

$$= \left[ \frac{7}{2}(\sqrt{3} + 1) + 7 \right]$$

$$= \frac{7}{2}(3 + \sqrt{3})\text{m}$$



**Fig. 10.22**

**EXERCISE 10.4**

1. In a right  $\triangle ABC$ ,  $\angle B = 90^\circ$ .
  - (i) Find BC, if  $AC = 15\text{m}$ ,  $\angle CAB = 30^\circ$
  - (ii) Find AB, if  $AC = 40\text{m}$ ,  $\angle CAB = 45^\circ$
  - (iii) Find BC if  $AC = 30\text{m}$ ,  $\angle CAB = 60^\circ$
2. Inclination of a ladder leaning against a wall is  $30^\circ$  and the foot of the ladder is 2m away from the wall. Find the length of ladder.
3. The string of a kite is 100m long and it makes an angle of  $60^\circ$  with the horizontal. Find the height of the kite, assuming that there is no slack in the string.
4. A ladder 10 m long is leaning against a wall. Its foot is 5 m away from the wall. Find the angle it makes with the wall.
5. A vertical post 9m high casts a shadow  $3\sqrt{3}$  m long. Find the altitude of the Sun.
6. From the top of a light house the angles of depression of two ships, one directly behind the other, are  $45^\circ$  and  $30^\circ$  respectively. If the ships are 200m apart, find the height of the light house.
7. The angle of elevation of the top of a tower from a point A on the ground, is  $30^\circ$ . On moving a distance of 20 m. towards the foot of the tower to a point B, the angle of elevation increases to  $60^\circ$ . Find the height of the tower and the distance of the tower from the point A.
8. Two men, on opposite sides of a temple 75m high, observe the angles of elevation of the top of the temple to be  $30^\circ$  and  $60^\circ$  respectively. Find the distance between the two men.
9. A pole 5m high is fixed on the top of a tower. The angle of elevation of the top of the pole observed from a point A on the ground is  $60^\circ$  and the angle of depression of the point A from the top of the tower is  $45^\circ$ . Find the height of the tower.
10. From the top of a tower 50 m high the angles of depression of the top and the bottom of a pole are observed to be  $45^\circ$  and  $60^\circ$  respectively. Find the height of the pole, if the pole and the tower stand on the same horizontal plane.
11. From the top of a tree 9 m. high, the angle of elevation of the top of a building and the angle of depression of the foot of the building are respectively  $45^\circ$  and  $60^\circ$ . What is the height of the building?
12. A vertical stick 10cm. long casts a shadow 8 cm. long. At the same time, a tree casts a shadow 3 m long. Find the height of the tree.

13. A pole is partly broken by the wind and its top touches the ground at an angle of  $30^\circ$  and at a distance of 8m. from the foot of the pole. Find the whole length of the pole.
14. From a light house the angles of depression of two ships on opposite sides of the light house are observed to be  $30^\circ$  and  $45^\circ$ . If the height of the light house be 100 metres, find the distance between the ships if the line joining them passes through the foot of the light house.
15. The angle of elevation of a bird from the eye of a man on the bank of a pond is  $30^\circ$  and the angle of depression of its reflection in the pond is  $60^\circ$ . Find the height of the bird above the pond if the distance of the eye from the foot of the man is 1.5 metres.
16. A bridge over a river makes an angle of  $45^\circ$  with the river bank. If the length of the bridge over the river is 150m. what is the width of the river?
17. From a point on the ground 40 m. away from the foot of a tower, the angle of elevation of the top of the tower is  $30^\circ$  and the angle of elevation of the top of a water tank (on the top of the tower) is  $45^\circ$ . Find the (i) height of the tower and (ii) depth of the tank.
18. A straight highway leads to the foot of a tower of height 50m. From the top of the tower, the angles of depression of two cars standing on the highway are  $30^\circ$  and  $60^\circ$ . What is the distance between the two cars and how far is each car from the tower ?
19. As observed from the top of a light house 100 m. above the water level, the angles of depression of two ships are  $30^\circ$  and  $45^\circ$ . If one ship is to the north and the other to the east of the light house, find the distance between the two ships.
20. The angles of elevation of the top of a tower from two points at distances  $a$  and  $b$  from the base and in the same straight line with it are complementary. Prove that height of the tower is  $\sqrt{ab}$ .
21. A tower subtends an angle  $\alpha$  at a point on the same level as the foot of the tower and at a second point  $h$  metres above the first, the depression of the foot of the tower is  $\beta$ . Show that the height of the tower is  $h \tan \alpha \cot \beta$
22. A vertical tower stands on a horizontal plane and is surmounted by a vertical flagstaff of height  $h$ . At a point on the plane, the angle of elevation of the bottom of the flagstaff is  $\alpha$  and that of the top of the flagstaff is  $\beta$ . Prove that the height of the tower is  $\frac{h \tan \alpha}{\tan \beta - \tan \alpha}$ .

## ANSWER

1. (i)  $BC = \frac{15}{2}$  cm      (ii)  $AB = 20\sqrt{2}$  cm      (iii)  $BC = 15\sqrt{3}$  cm
2.  $\frac{4\sqrt{3}}{3}$  m.
3.  $50\sqrt{3}$
4.  $30^\circ$
5.  $60^\circ$
6. 273.2m
7. height of tower = 17.3m  
distance of tower from the point A = 30m.
8.  $100\sqrt{3}$  m.
9. 6.83 m.
10. 21.13 m.
11.  $(9+3\sqrt{3})$  m.
12. 3.75m.
13.  $8\sqrt{3}$  m.      14.  $100(1+\sqrt{3})$  m.      15. 3 metres      16.  $75\sqrt{2}$  m.
17.  $\frac{40\sqrt{3}}{3}$  m.,  $\frac{40(3-\sqrt{3})}{3}$  m.      18.  $\frac{100\sqrt{3}}{3}$  m.;  $50\sqrt{3}$  m. and  $\frac{50\sqrt{3}}{3}$  m.
19. 200 m.

## SUMMARY

In this chapter, you have learnt the following points:

1. In a right  $\triangle ABC$ , right angled at B,

$$\sin A = \frac{\text{side opposite to } \angle A}{\text{hypotenuse}}, \quad \cos A = \frac{\text{side adjacent to } \angle A}{\text{hypotenuse}}$$

$$\tan A = \frac{\text{side opposite to } \angle A}{\text{side adjacent to } \angle A}, \quad \cot A = \frac{\text{side adjacent to } \angle A}{\text{side opposite to } \angle A}$$

$$\sec A = \frac{\text{hypotenuse}}{\text{side adjacent to } \angle A}, \quad \operatorname{cosec} A = \frac{\text{hypotenuse}}{\text{side opposite to } \angle A}$$

2. The values of trigonometric ratios for angles  $0^\circ$ ,  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$ , and  $90^\circ$ .

3.  $\operatorname{cosec} A = \frac{1}{\sin A}$ ,  $\sec A = \frac{1}{\cos A}$ ,  $\cot A = \frac{1}{\tan A}$ ,  $\tan A = \frac{\sin A}{\cos A}$ ,  $\cot A = \frac{\cos A}{\sin A}$ ,

$$\sin^2 A + \cos^2 A = 1, \quad \sec^2 A = 1 + \tan^2 A, \quad \operatorname{cosec}^2 A = 1 + \cot^2 A$$

4. If one of the trigonometric ratios of an acute angle is known, the remaining trigonometric ratios of the angle can be easily determined.

5. The values of  $\sin A$  or  $\cos A$  never exceeds 1, whereas the value of  $\sec A$  or  $\operatorname{cosec} A$  is always greater than or equal to 1.

6.  $\sin(90^\circ - A) = \cos A$ ,  $\cos(90^\circ - A) = \sin A$   
 $\tan(90^\circ - A) = \cot A$ ,  $\cot(90^\circ - A) = \tan A$   
 $\sec(90^\circ - A) = \operatorname{cosec} A$ ,  $\operatorname{cosec}(90^\circ - A) = \sec A$ .

\*\*\*\*\*

### 11.1 Introduction

In class IX, you have seen that to locate the position of a point on a plane, a pair of mutually perpendicular lines known as the coordinate axes are required. Usually, one of the axes is taken along a horizontal line and the other along a vertical line. The horizontal line is known as the  $x$ -axis and the vertical line is known as the  $y$ -axis. The point of intersection of the  $x$ -axis and the  $y$ -axis is known as the origin. The distance of a point from the  $y$ -axis is called its  $x$ -coordinate or abscissa and the distance of a point from the  $x$ -axis is called its  $y$ -coordinate or ordinate. You have seen that the coordinates of a point on the plane are of the form  $(x, y)$  where  $x$  is the abscissa and  $y$  is the ordinate. The coordinates of a point on the  $x$ -axis are of the form  $(x, 0)$  and those of a point on the  $y$ -axis are of the form  $(0, y)$ . You have also learnt about plotting of points in a plane when their coordinates are given. Further, you have learnt that the distance between two points  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  is  $\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$  and in particular, the distance of the point  $R(x, y)$  from the origin is  $\sqrt{x^2 + y^2}$ .

In this chapter, you will see how the co-ordinates of the point, which divides the line segment joining two given points in a given ratio, can be found. Next, the area of a triangle will be found in terms of the coordinates of its vertices. Henceforth, we shall write,  $(x_1, y_1) = (x_2, y_2)$  whenever the points, whose coordinates are  $(x_1, y_1)$  and  $(x_2, y_2)$ , are coincident.

### 11.2 SECTION FORMULA

Let  $P$  and  $Q$  be two points in a plane say, the plane of the paper (Fig 11.1). Let us take a ratio say, 2:1, at which the line  $PQ$  is to be divided. Let  $R$  be the point on  $PQ$  such that  $PR = 2RQ$  (Fig 11.1). Then  $PR:RQ = 2:1$ . Therefore, the point  $R$  divides  $PQ$  in the ratio 2:1 and observe that  $R$  lies on  $PQ$ .





Fig. 11.1



Again, let  $R'$  be the point on  $PQ$  produced (Fig 11.2) such that  $PQ = QR'$ . Then  $PR' = PQ + QR' = 2QR'$  ie.  $PR' : R'Q = 2:1$



Therefore, the point  $R'$  also divides  $PQ$  in the same ratio 2:1 and observe that  $R'$  lies on  $PQ$  produced. We have seen that  $R$  and  $R'$  divide  $PQ$  in the same ratio 2:1. We have also observed that  $R$  lies on  $PQ$  whereas  $R'$  lies on  $PQ$  produced. Here  $R$  is said to divide  $PQ$  internally and  $R'$  is said to divide  $PQ$  externally in the ratio 2:1. In other words,  $R$  is the point of internal division of  $PQ$  in the ratio 2:1 and  $R'$  is the point of external division of  $PQ$  in the same ratio 2:1.

Thus, we have seen that, for a given ratio, there are two points of division of a line joining two points: one, the point of internal division and the other, the point of external division. Here, we shall consider the case of internal division only.

In this section, we will find a formula, known as section formula, for finding the coordinates of the point  $R$  in terms of the coordinates of  $P$  and  $Q$  when  $R$  divides  $PQ$  internally in the ratio  $m:n$ .

Let  $XOX$ ,  $YOY$  be the coordinate axes so that  $O$  is the origin. Let  $(x_1, y_1)$  and  $(x_2, y_2)$  be the coordinates of  $P$  and  $Q$  respectively. Let  $(x, y)$  be the coordinates of the point  $R$  which divides  $PQ$  internally in the ratio  $m:n$  (Fig 11.3).

Draw  $PA$ ,  $QB$  and  $RC$  perpendicular to the  $x$ -axis and draw  $PD$ ,  $RE$  perpendicular to  $RC$ ,  $QB$  respectively as in Fig 11.3.

Then, in  $\triangle PDR$ ,

$$PD = AC = OC - OA = x - x_1$$

$$RD = CR - CD = y - y_1$$

In  $\triangle REQ$

$$RE = CB = OB - OC = x_2 - x$$

$$QE = BQ - BE = BQ - CR = y_2 - y$$

In  $\triangle PDR$  and  $REQ$ ,

$$\angle PDR = \angle REQ = 90^\circ \text{ (why ?)}$$

$$\angle RPD = \angle QRE \text{ (corresponding angles)}$$

∴ by AA similarity

$$\triangle PDR \sim \triangle REQ$$

$$\therefore \frac{PR}{RQ} = \frac{PD}{RE} = \frac{RD}{QE}$$

$$\text{i.e. } \frac{m}{n} = \frac{x - x_1}{x_2 - x} = \frac{y - y_1}{y_2 - y}$$

$$\text{Considering } \frac{m}{n} = \frac{x - x_1}{x_2 - x},$$


$$\text{we get } x = \frac{mx_2 + nx_1}{m + n}$$

$$\text{Again, considering } \frac{m}{n} = \frac{y - y_1}{y_2 - y}, \text{ we get } y = \frac{my_2 + ny_1}{m + n}$$

$$\therefore (x, y) = \left( \frac{mx_2 + nx_1}{m + n}, \frac{my_2 + ny_1}{m + n} \right)$$

Thus, the coordinates of the point R which divides the line joining  $P(x_1, y_1)$  and  $Q(x_2, y_2)$

in the ratio  $m:n$  are  $\left( \frac{mx_2 + nx_1}{m + n}, \frac{my_2 + ny_1}{m + n} \right)$ . It is known as the section formula.



**Note:** To find the abscissa of the point of division,  $m$  multiplies to  $x_2$  and  $n$  to  $x_1$  i.e. cross wise. Similar is the case for finding the ordinate also.

**Deduction:** The mid-point of a line segment is the point which divides the line segment in the ratio 1:1. Thus, the coordinates of the mid-point of the line segment joining  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  are

$$\left( \frac{1 \cdot x_2 + 1 \cdot x_1}{1 + 1}, \frac{1 \cdot y_2 + 1 \cdot y_1}{1 + 1} \right) \text{ i.e. } \left( \frac{x_2 + x_1}{2}, \frac{y_2 + y_1}{2} \right)$$

**Example 1.** Find the coordinates of the point which divides the line segment joining the points  $(-1, 3)$  and  $(4, -7)$  internally in the ratio 3:4.

**Solution :** Let  $P(x, y)$  be the required point. Then, by Section formula,

$$x = \frac{3 \cdot 4 + 4 \cdot (-1)}{3 + 4} = \frac{8}{7}, \quad y = \frac{3 \cdot (-7) + 4 \cdot 3}{3 + 4} = \frac{-9}{7}$$

$\therefore$  the required point is  $\left(\frac{8}{7}, \frac{-9}{7}\right)$

**Example 2.** Find the points of trisection of the line segment joining the points A(3, -2) and B(-3, -4).

**Solution:** Let P and Q be the points of trisection of AB

Then, P divides AB in the ratio 1:2

$$\therefore \text{The co-ordinates of P are } \left(\frac{1 \times (-3) + 2 \times 3}{1 + 2}, \frac{1 \times (-4) + 2 \times (-2)}{1 + 2}\right)$$



i.e.  $\left(1, \frac{-8}{3}\right)$

Again, Q divides AB in the ratio 2:1

$$\therefore \text{the coordinates of Q are } \left(\frac{2 \times (-3) + 1 \times 3}{2 + 1}, \frac{2 \times (-4) + 1 \times (-2)}{2 + 1}\right)$$

i.e.  $\left(-1, \frac{-10}{3}\right)$

$\therefore$  the required points of trisection of AB are  $\left(1, \frac{-8}{3}\right)$  and  $\left(-1, \frac{-10}{3}\right)$

**Example 3.** In what ratio, is the line joining (-3, -1) and (-8, -9) divided at the point  $\left(-5, \frac{-21}{5}\right)$  ?

**Solution :** Let the required ratio be m:n

$$\text{Then, } \left(-5, \frac{-21}{5}\right) = \left(\frac{m \cdot (-8) + n \cdot (-3)}{m + n}, \frac{m \cdot (-9) + n \cdot (-1)}{m + n}\right)$$

i.e.  $-5 = \frac{-8m - 3n}{m + n}, \quad \frac{-21}{5} = \frac{-9m - n}{m + n}$

From the 1st eq<sup>n</sup>,

$$-5(m + n) = -8m - 3n$$

i.e.  $8m - 5m = 5n - 3n$

i.e.  $3m = 2n$

i.e.  $\frac{m}{n} = \frac{2}{3}$

i.e.  $m:n = 2:3$

∴ The required ratio is 2:3

**Note:** The ratio  $m:n$  could have been solved from the 2nd eqn  $-\frac{21}{5} = \frac{-9m - n}{m + n}$ , as well.

**Example 4.** Find the ratio in which the line segment joining A(2, - 4) and B(-3,6) is divided by the x-axis. Find the point of division also.

**Solution :** Let  $m:n$  be the required ratio. Then, the coordinates of the point dividing AB in

the ratio  $m:n$  are  $\left(\frac{-3m + 2n}{m + n}, \frac{6m - 4n}{m + n}\right)$ . But this point lies on the x-axis and so, the ordinate is zero,

i.e.  $\frac{6m - 4n}{m + n} = 0$

i.e.  $6m - 4n = 0$

i.e.  $6m = 4n$

i.e.  $\frac{m}{n} = \frac{2}{3}$

i.e.  $m:n = 2:3$


∴ The required ratio is 2:3

Again, the point of division is  $\left(\frac{-3m + 2n}{m + n}, \frac{6m - 4n}{m + n}\right)$

i.e.  $\left(\frac{-3\frac{m}{n} + 2}{\frac{m}{n} + 1}, 0\right)$  i.e.  $\left(\frac{-3 \times \frac{2}{3} + 2}{\frac{2}{3} + 1}, 0\right)$  i.e. (0,0)

**Example 5.** The line segment joining A(-2,9) and B(6,3) is a diameter of a circle with centre C. Find the coordinates of C.

**Solution :**



Since the centre C is the mid-point of the diameter AB, the coordinates of C are  $\left(\frac{-2+6}{2}, \frac{9+3}{2}\right)$  i.e. (2,6).

**Example 6.** Find the coordinates of the centroid of the triangle with vertices  $A(x_1, y_1)$ ,  $B(x_2, y_2)$  and  $C(x_3, y_3)$ .


**Solution :** Let D be the mid point of the side AB.

Then the co-ordinates of D are  $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$ . Join CD and let G be the centroid of  $\triangle ABC$ . Then G divides CD internally in the ratio 2:1.

$\therefore$  the coordinates of G are

$$\left(\frac{2 \cdot \frac{x_1+x_2}{2} + 1 \cdot x_3}{2+1}, \frac{2 \cdot \frac{y_1+y_2}{2} + 1 \cdot y_3}{2+1}\right)$$

$$\text{i.e. } \left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$$



$$\therefore \text{ the centroid } \triangle ABC \text{ is } \left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$$

**Example 7.** Show that the points A (3,1) , B(0, -2) , C(1,1) and D (4,4) are the vertices of a parallelogram ABCD.

**Solution:** Mid-points of AC


$$= \left(\frac{3+1}{2}, \frac{1+1}{2}\right) = (2,1)$$

Again, mid-point of BD

$$= \left(\frac{0+4}{2}, \frac{-2+4}{2}\right) = (2,1)$$

$\therefore$  the mid-point of AC coincides with the mid-point of BD i.e., the diagonals of ABCD bisect each other.

$\therefore$  ABCD is a parallelogram.



**Example 8.** If three cosecutive vertices of a par<sup>m</sup> are A(1,-2), B(3,6) and C(5,10), find its fourth vertex.

**Solution :** Let D (x,y) be the fourth vertex of the par<sup>m</sup>.  
Then, the diagonals AC and BD bisect each other.

∴ mid-point of AC = mid-point of BD

$$\text{i.e. } \left( \frac{1+5}{2}, \frac{-2+10}{2} \right) = \left( \frac{3+x}{2}, \frac{6+y}{2} \right)$$

$$\text{i.e. } (3,4) = \left( \frac{3+x}{2}, \frac{6+y}{2} \right)$$

$$\text{i.e. } 3 = \frac{3+x}{2}, 4 = \frac{6+y}{2}$$

$$\text{i.e. } x = 3, y = 2.$$

∴ The fourth vertex is (3,2)

### EXERCISE 11.1

1. Find the coordinates of the point which divides the line segment joining the points (6,3) and (-4,5) in the ratio 3:2 internally.
2. Find the coordinates of the point which divides the line joining (1,-2) and (4,7) internally in the ratio 1:2.
3. Find the coordinates of the mid-point of the line joining the points (4,7) and (6,9).
4. Find the co-ordinates of the point which divides internally the line joining the points (p,q) and (q,p) in the ratio p-q : p+q.
5. Let A(-1,5) and B(6,-2) be two given points. Find the coordinates of the point P such that  $AP = \frac{3}{7} AB$  and P lies on AB.
6. The mid-point of the line segment joining A(2a,4) and B(-2,3b) is M(1,2a+1). Find the values of a and b.
7. AB is a diameter of a circle with centre C(-1,6). If the coordinates of A are (-7,3), find the coordinates of B.
8. Find the coordinates of the point where the diagonals of the parallelogram ABCD formed by joining the points A(-2,-1), B(1,0), C(4,3) and D(1,2) meet.
9. Find the ratio in which the line segment joining (-2,-3) and (5,6) is divided by x-axis. Also, find the coordinates of the point of division.
10. In what ratio is the line segment joining the points (-2,-3) and (3,7) divided by the y-axis? Also, find the co-ordinates of the point of division.

11. If a vertex of a triangle be  $(1,1)$  and the middle points of the sides through it be  $(-2,3)$  and  $(5,2)$ , find the other vertices.
12. Show that the points  $(3,-2)$ ,  $(4,0)$ ,  $(6,-3)$  and  $(5,-5)$  are the vertices of a parallelogram taken in order.
13. Show that the four points  $(1,2)$ ,  $(3,0)$ ,  $(7,4)$  and  $(5,6)$  taken in order are the angular points of a rectangle.
14. Determine the ratio in which the point  $P(10,m)$  divides the join of  $A(5,2)$  and  $B(17,14)$ . Also, find the value of  $m$ .
15. Find the centroid of  $\triangle ABC$  whose vertices are  $A(-1,0)$ ,  $B(5,-2)$  and  $C(8,2)$ .
16. If the points  $A(a,-11)$ ,  $B(5,b)$ ,  $C(2,15)$  and  $D(1,1)$  taken in order are the vertices of a parallelogram  $ABCD$ , find the values of  $a$  and  $b$ .
17. Three vertices of a rectangle are the points  $(3,4)$ ,  $(-1,2)$  and  $(2,-4)$ ; what are the coordinates of the fourth vertex.

### ANSWER


- |                                       |                    |                     |  |
|---------------------------------------|--------------------|---------------------|--|
| 1. $\left(0, \frac{21}{5}\right)$     | 2. $(2,1)$         | 3. $(5,8)$          | 4. $\left(\frac{p^2 - q^2 + 2pq}{2p}, \frac{p^2 + q^2}{2p}\right)$ |
| 5. $(2,2)$                            | 6. $a=2, b=2$      | 7. $(5,9)$          | 8. $(1,1)$   |
| 9. $1:2; \left(\frac{1}{3}, 0\right)$ | 10. $2:3; (0,1)$   | 11. $(-5,5), (9,3)$ | 14. $5:7; m = 7$   |
| 15. $(4,0)$                           | 16. $a = 4, b = 3$ | 17. $(6, -2)$       |  |

### 11.3 AREA OF A TRIANGLE

In earlier classes, the area of a triangle have been computed by using the formula

$$\text{Area of a triangle} = \frac{1}{2} \times \text{Base} \times \text{Altitude}$$

In class IX, you have used Heron's formula to find the area of a triangle when the length of its sides are given. When the coordinates of the vertices of a triangle are given, you can find the lengths of the three sides of the triangle by using distance



formula and hence the area of the triangle by using the Heron's formula. But, when the lengths of the sides are irrational numbers, this process becomes tedious. So, it is preferred to compute the area in terms of the coordinates of the vertices of the triangle as follows:

Let  $A(x_1, y_1)$ ,  $B(x_2, y_2)$  and  $C(x_3, y_3)$  be the vertices of a  $\triangle ABC$  (fig 11.4).

Draw  $AL$ ,  $BM$  and  $CN$  perpendicular to the  $x$ -axis as in the adjoining figure. From the figure

$$\begin{aligned} \text{Area of } \triangle ABC &= \text{Area of trapezium } ABML + \text{Area of trapezium } ALNC \\ &\quad - \text{Area of trapezium } BMNC \end{aligned}$$

But, we know that area of trapezium

$$= \frac{1}{2} \times \text{sum of parallel sides} \times \text{perp. distance between them}$$

$\therefore$  Area of  $\triangle ABC$

$$\begin{aligned} &= \frac{1}{2} (AL+BM)ML + \frac{1}{2} (AL+CN)LN - \frac{1}{2} (BM+CN)MN \\ &= \frac{1}{2} (y_1 + y_2)(x_1 - x_2) + \frac{1}{2} (y_1 + y_3)(x_3 - x_1) - \frac{1}{2} (y_2 + y_3)(x_3 - x_2) \\ &= \frac{1}{2} [x_1 \{(y_1 + y_2) - (y_1 + y_3)\} + x_2 \{(y_2 + y_3) - (y_1 + y_2)\} + x_3 \{(y_1 + y_3) - (y_2 + y_3)\}] \\ &= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \end{aligned}$$

$$\therefore \text{Area of a triangle} = \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

**Sign of Area :** If the three points are taken in the anticlockwise sense, then the area calculated of the  $\triangle ABC$  will be positive, while if the points are taken in clockwise sense, then the area calculated will be negative. In case, the area calculated is negative, the negative sign will be ignored i.e. the absolute value of the area is taken only because negative area is meaningless.

**Note: (1)** If the area of a  $\triangle ABC$  is zero, then the three vertices  $A$ ,  $B$ ,  $C$  are collinear (ie, lie on the same line) and conversely.

**(2)** To find the area of a polygon, we divide it into triangles having no common area and take the numerical value of the area of each of the triangles and add.



**Example 1.** Find the area of a triangle whose vertices are (6,3),(-3,4) and (4,-2).

**Solution :** Let  $A(x_1, y_1) = (6, 3)$ ,  $B(x_2, y_2) = (-3, 4)$  and  $C(x_3, y_3) = (4, -2)$

Then,

$$\begin{aligned} \text{area of } \triangle ABC &= \frac{1}{2}[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \\ &= \frac{1}{2}[6(4 - 2) + (-3)(-2 - 3) + 4(3 - 4)] \\ &= \frac{1}{2}[36 + 15 - 4] \\ &= \frac{1}{2} \times 47 \\ &= \frac{47}{2} \end{aligned}$$

$\therefore$  the required area of the triangle is  $\frac{47}{2}$  sq. units.

**Example 2.** Find the area of the triangle formed by the points (a, c + a), (a,c) and (-a, c-a).

**Solution :** Let  $A(x_1, y_1) = (a, c + a)$ ,  $B(x_2, y_2) = (a, c)$  and  $C(x_3, y_3) = (-a, c - a)$ .

$$\begin{aligned} \text{Then, area of } \triangle ABC &= \frac{1}{2}[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \\ &= \frac{1}{2}[a(c - c + a) + (c - a - c - a) + (-a)(c + a - c)] \\ &= \frac{1}{2}[a^2 + a(-2a) - a(a)] \\ &= \frac{1}{2}[a^2 - 3a^2] \\ &= -a^2 \end{aligned}$$

Neglecting the -ve sign, area of  $\triangle ABC = a^2$  sq. units.

**Example 3.** Show that the points (-5, 1), (5,5) and (10,7) are collinear.

**Solution :** Let  $A(x_1, y_1) = (-5, 1)$ ,  $B(x_2, y_2) = (5, 5)$  and  $C(x_3, y_3) = (10, 7)$

$$\text{Then, area of } \triangle ABC = \frac{1}{2}[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

$$\begin{aligned}
 &= \frac{1}{2}[(-5)(5-7)+5(7-1)+10(1-5)] \\
 &= \frac{1}{2}[10+30-40] \\
 &= \frac{1}{2} \times 0 \\
 &= 0
 \end{aligned}$$

∴ the points A, B, C i.e. the given points are collinear.

**Example 4.** Find the value of p for which the points (-5,1), (1,p) and (4,-2) are collinear.

**Solution :** Let A(x<sub>1</sub>, y<sub>1</sub>) = (-5,1), B(x<sub>2</sub>, y<sub>2</sub>) = (1, p) and C(x<sub>3</sub>, y<sub>3</sub>) = (4,-2).

Then, since A, B,C are collinear, area of ΔABC = 0

$$\therefore \frac{1}{2}[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] = 0$$

$$\text{i.e. } (-5)(p + 2) + 1(-2 - 1) + 4(1 - p) = 0$$

$$\text{i.e. } -5p - 10 - 3 + 4 - 4 - 4 = 0$$

$$\text{i.e. } 9p = -9$$

Hence, p = -1.

**Example 5.** Find the area of the quadrilateral whose vertices are (1,1),(3,4), (5,-2) and (4,-7) taken in order.

**Solution:** Let A(1,1), B(3,4), C(5,-2), D(4,-7) be the vertices.

Then, area of quadrilateral ABCD

= area of ΔABC + area of ΔACD


$$= \frac{1}{2}|1(4+2)+3(-2-1)+5(1-4)|$$

$$+ \frac{1}{2}|1(-2+7)+5(-7-1)+4(1+2)|$$

$$= \frac{1}{2}|6-9-15| + \frac{1}{2}|5-40+12|$$

$$= \frac{1}{2}|-18| + \frac{1}{2}|-23|$$

$$= \left(\frac{1}{2} \times 18\right) + \left(\frac{1}{2} \times 23\right)$$



$$= \frac{41}{2}$$

$\therefore$  area of the quadrilateral is  $\frac{41}{2}$  sq. units.

### EXERCISE 11.2

1. Find the area of a triangle whose vertices are

(i)  $(-5, -3), (-4, -6), (2, -1)$

(ii)  $\left(at_1, \frac{a}{t_1}\right), \left(at_2, \frac{a}{t_2}\right), \left(at_3, \frac{a}{t_3}\right)$

(iii)  $(5, 2), (4, 7), (7, -4)$

2. Find the area of a quadrilateral whose vertices are  $(-4, -2), (-3, -5), (3, -2)$  and  $(2, 3)$  taken in order.

3. Show that the following points are collinear.

(i)  $(-1, 1), (5, 7), (8, 10)$

(ii)  $(3a, 0), (0, 3b), (a, 2b)$

(iii)  $(a, b+c), (b, c+a), (c, a+b)$

4. In each of the following, find the value of 'x' for which the points are collinear.

(i)  $(8, 1), (x, -4), (2, -5)$

(ii)  $\left(-\frac{1}{2}, 3\right), (-5, 6), (-8, x)$

(iii)  $(x, 5), (2, -5), (4, -9)$

5. If the points  $(x, y), (-5, 7)$  and  $(-4, 5)$  are collinear, then show that  $2x + y + 3 = 0$ .

6. Prove that the points  $(a, 0), (0, b)$  and  $(1, 1)$  are collinear, if  $\frac{1}{a} + \frac{1}{b} = 1$ .

7. If the points  $(x_1, y_1), (x_2, y_2), (x_3, y_3)$  lie on the same line, prove that

$$\frac{y_2 - y_3}{x_2 x_3} + \frac{y_3 - y_1}{x_3 x_1} + \frac{y_1 - y_2}{x_1 x_2} = 0$$

8. If a, b, c are distinct, prove that the points  $(a, a^2)$ ,  $(b, b^2)$ ,  $(c, c^2)$  can never be collinear.
9. Find the area of the quadrilateral whose vertices are (1,1), (7, -3), (12, 2) and (7, 21) taken in order.
10. Find the area of the rhombus whose vertices, taken in order, are (3,0), (4,5), (-1,4), (-2, -1).

**ANSWER**

1. (i)  $\frac{23}{2}$  sq. units (ii)  $\frac{a^2}{2t_1 t_2 t_3} (t_1 - t_2)(t_2 - t_3)(t_3 - t_1)$  (iii) 2 sq. units
2. 28 sq. units    4. (i) 3 (ii) 8 (iii) -3    9. 132 sq. units
10. 24 sq. units.


**SUMMARY**

In this chapter, the following points have been studied :

1. The co-ordinates of the point R which divides the line joining  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  internally in the ratio  $m : n$  are  $\left( \frac{mx_2 + nx_1}{m + n}, \frac{my_2 + ny_1}{m + n} \right)$
2. The mid-point of the line segment joining  $(x_1, y_1)$  and  $(x_2, y_2)$  is  $\left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$
3. Area of a triangle with vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$  is  $\frac{1}{2} |x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)|$

\*\*\*\*\*

©BOSEM Not to be republished



In the subsequent sections, we shall discuss the methods of finding surface area and volume of such combinations.

## 12.2 Perimeter and Area of a Circle

In class (VIII), you have learnt about the perimeter and the area of a circle. Recall that the perimeter of a circle is the measure of its circumference and the circumference of a circle bears a constant ratio to its diameter. This constant ratio is denoted by the Greek letter  $\pi$  (read as pi). In other words,

$$\frac{\text{Circumference}}{\text{Diameter}} = \pi$$

$$\begin{aligned} \therefore \text{Circumference} &= \pi \times \text{Diameter} \\ &= \pi \times d, \text{ where 'd' denotes diameter} \\ &= \pi \times 2r, \text{ where 'r' denotes radius} \\ &= 2\pi r \end{aligned}$$

The value of  $\pi$  is generally taken as  $\frac{22}{7}$  or 3.14. Both these are approximate values. Further, you may recall that the area of a circle is  $\pi r^2$ , where 'r' is the radius of the circle. You have also verified this fact in class (viii), by cutting a circle into a number of sectors and rearranging them [see Fig. 12.3]. Here the shape in Fig. 12.3(ii) approaches a rectangle of length  $\frac{1}{2} \times 2\pi r$  and breadth  $r$ , giving that the area of the circle

$$= \frac{1}{2} \times 2\pi r \times r = \pi r^2.$$

$$\begin{aligned}\Rightarrow R &= \frac{616}{2\pi} \\ &= \frac{616 \times 7}{2 \times 22} \\ &= 98 \text{ m}\end{aligned}$$

Also, the width of the track = 7 m

$$\begin{aligned}\therefore \text{the inner radius of the track, } r &= (98-7) \\ &= 91 \text{ m}\end{aligned}$$

$$\begin{aligned}\text{and the area of the track} &= \pi(R^2 - r^2) \\ &= \frac{22}{7}(98+91)(98-91) \\ &= \frac{22}{7} \times 189 \times 7 \\ &= 4158 \text{ m}^2\end{aligned}$$

$\therefore$  the required area of the track is 4158 m<sup>2</sup>

### EXERCISE 12.1

$\left( \text{Unless otherwise stated, use } \pi = \frac{22}{7} \right)$

1. Find the area of a circle whose radius is 14 cm.
2. Find the area of a quadrant of a circle whose circumference is 44 cm.
3. The circumference of a circle exceeds the diameter by 21 cm. Find the area of the circle.
4. The area of a circle is 1386 cm<sup>2</sup>; find its circumference.
5. The sum of the radii of two circles is 35 cm and the difference of their areas is 770 cm<sup>2</sup>. Find their circumferences.
6. The diameter of each wheel of a vehicle is 42 cm. Each wheel makes 20 revolutions in 12 seconds. Find the speed of the vehicle in metre per minute.
7. Find the radius of a circle whose area is equal to the difference of the areas of two circles of radii 6 cm and 10 cm respectively.
8. The difference between the perimeter and the radius of a circle is 37 cm. Find the area of the circle.

**Recall :**

The line segment joining any two points of a circle is called a **Chord**. In Fig. 12.5,  $\overline{AB}$  is a chord of the circle.

A part of a circle between two distinct points on it, is called an **arc** of the circle. In fig. 12.5,  $\widehat{AXB}$  is a **minor** arc and  $\widehat{BYA}$  is a **major** arc.

The part (or portion) of the circular region enclosed by two radii (say)  $OA$ ,  $OB$  and the corresponding arc  $AB$  of a circle is called a sector of the circle. [Fig. 12.6] Here, for the sector  $OAXB$ ,  $\angle AOB$  is called sectorial angle. In Fig. 12.6, the shaded region  $OAXB$  is a minor sector and the unshaded region  $OAYB$  is the corresponding major sector.

The part (or portion) of the circular region enclosed by a chord and the corresponding arc of a circle is called a segment of a circle. In Fig. 12.7, the shaded region  $AXB$  is the minor segment and the unshaded region  $AYB$  is the major segment.

Let us now find the area of a sector. In Fig. 12.6,  $OAXB$  is a sector of a circle with centre  $O$  and radius  $r$ .

Let  $\angle AOB = \theta$  (in degrees).

Here, you recall that the area of a circle is  $\pi r^2$ , where  $r$  is the radius of the circle.

We may consider the whole circular region as a sector of sectorial angle  $360^\circ$ .

When the sectorial angle is  $360^\circ$ , the area of the sector is  $\pi r^2$

$\therefore$  when the sectorial angle is  $1^\circ$ , the area of the sector is  $\frac{\pi r^2}{360}$



$\therefore$  when the sectorial angle is  $\theta^\circ$ , the area of the sector is  $\frac{\pi r^2 \theta}{360}$

Thus, the area of a sector of sectorial angle  $\theta$ , is  $\frac{\pi r^2 \theta}{360}$ .

From Fig. 12.6 we may observe that

area of the major sector  $OAYB$  = area of circle – area of the minor sector  $OAXB$

$$\begin{aligned} &= \pi r^2 - \frac{\pi r^2 \theta}{360} \\ &= \left( \frac{360 - \theta}{360} \right) \pi r^2 \end{aligned}$$





$$= \frac{r^2}{2} \left( \frac{\theta\pi}{180} - \sin\theta \right)$$

And area of the major segment OAYB = area of the circle - area of the minor segment

$$\begin{aligned} &= \pi r^2 - \frac{r^2}{2} \left[ \frac{\theta\pi}{180} - \sin\theta \right] \\ &= r^2 \left[ \pi \left( 1 - \frac{\theta}{360} \right) + \frac{\sin\theta}{2} \right] \\ &= r^2 \left[ \frac{\pi(360 - \theta)}{360} + \frac{\sin\theta}{2} \right] \end{aligned}$$

Henceforth 'sector' and 'segment' will mean the 'minor sector' and the 'minor segment' respectively, unless otherwise stated.

**Example 3.** If a chord of a circle of radius 12 cm subtends an angle of  $60^\circ$  at the centre, find the area of the corresponding (i) minor sector (ii) major sector and (iii) major segment (take  $\pi = 3.14$  and  $\sqrt{3} = 1.73$ ).

**Solution :** Here,

$$\theta = 60^\circ, r = 12 \text{ cm}$$


Then

$$\begin{aligned} \text{(i) the area of the minor sector} &= \frac{\pi r^2 \theta}{360} \\ &= 3.14 \times 12^2 \times \frac{60}{360} \\ &= 3.14 \times 12 \times 2 \\ &= 75.36 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{(ii) the area of the corresponding major sector} &= \pi r^2 - \text{area of the minor sector} \\ &= 3.14 \times 12 \times 12 - 75.36 \\ &= 452.16 - 75.36 \\ &= 376.8 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{(iii) we know that the area of the major segment} &= \pi r^2 - \text{area of the minor segment} \dots(1) \end{aligned}$$

[ For obtaining the area of the minor segment refer Fig 12.9.]



**Example 5.** A chord of a circle of radius 4.2 cm subtends  $120^\circ$  at the centre. Find the area of the corresponding minor segment of the circle.

(take  $\pi = \frac{22}{7}$  and  $\sqrt{3} = 1.73$ )

**Solution :** We have

$$r = 4.2 \text{ cm}$$

$$\theta = 120^\circ$$

For finding the minor segment refer Fig. 12.10. In  $\triangle OAB$ , draw  $OM \perp AB$  and  $OA=OB$ .

$\therefore$  by RHS congruence,  $\triangle AMO \cong \triangle BMO$ .

So, M is the mid-point of AB

and  $\angle AOM = \angle BOM$

$$\begin{aligned} &= \frac{1}{2} \times 120^\circ \\ &= 60^\circ \end{aligned}$$

From  $\triangle OMA$ , we have

$$\frac{OM}{OA} = \cos 60^\circ = \frac{1}{2}$$

$$\Rightarrow OM = \frac{1}{2} \times OA$$

$$\text{And } \frac{AM}{OA} = \sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\therefore AM = \frac{\sqrt{3}}{2} \times OA$$


$$\therefore \text{ area of the } \triangle OAB = \frac{1}{2} \times AB \times OM$$

$$= AM \times OM$$

$$= \frac{\sqrt{3}}{2} \times OA \times \frac{1}{2} \times OA$$

$$= \frac{\sqrt{3}}{4} \times (4.2)^2$$

$$= 7.63 \text{ cm}^2$$



**Fig. 12.10**

10. A chord of a circle of radius 12 cm subtends a right angle at the centre. Find the area of the corresponding (i) major sector and (ii) minor segment (use  $\pi = 3.14$ ).
11. A chord of a circle of radius 18 cm subtends an angle of  $120^\circ$  at the centre. Find the area of the corresponding (i) minor sector and (ii) major segment (use  $\pi=3.14$  and  $\sqrt{3}=1.73$ ).
12. A chord of a circle of radius 6 cm subtends an angle of  $60^\circ$  at the centre. Find the area of the corresponding (i) minor segment and (ii) major segment.(take  $\pi=3.14$  and  $\sqrt{3} = 1.73$  ) .
13. The radius of a circle is 5 cm. A chord of length  $5\sqrt{2}$  cm is drawn in the circle. Find the area of the major segment (take  $\pi=3.14$ ).
14. A chord of a circle of diameter 12 cm makes an angle of  $30^\circ$  at the centre. Find the area of the minor segment of the circle.
15. A chord of a circle subtends an angle of  $60^\circ$  at the centre . If the length of the chord is 12 cm, find the area of the two segments into which the chord divides the circle (take  $\pi=3.14$  and  $\sqrt{3} = 1.73$ )
16. A grass field is in the form of an equilateral triangle of sides 48 m. A cow is tethered at a vertex of the field. If the cow can reach upto 21 m from the vertex, find  
 (i) the area of the part of the field in which the cow can graze.  
 (ii) how much area remains ungrazed.  
 (iii) the increase in grazing area if the length of the rope be increased to 35 m (use  $\pi=3.14$  ).

### ANSWER

1.  $26 \text{ cm}^2$     2.  $60^\circ$     3.  $308 \text{ cm}^2$     4.  $44 \text{ cm}$   
 5.  $90^\circ, 24.64 \text{ cm}^2$     6.  $18.48 \text{ cm}^2$     7. (i)  $51.33 \text{ cm}^2$     (ii)  $77 \text{ cm}^2$   
 (iii)  $102.67 \text{ cm}^2$  (iv)  $154 \text{ cm}^2$     (v)  $205.33 \text{ cm}^2$     8.  $10.47 \text{ cm}^2$   
 9.(i)  $9.9$     (ii)  $31.19 \text{ cm}^2$  (iii)  $22.5 \text{ cm}$     (iv)  $11.34 \text{ cm}^2, 113.40 \text{ cm}^2$   
 10. (i)  $339.12 \text{ cm}^2$ ,    (ii)  $41.04 \text{ cm}^2$  11.  $339.12 \text{ cm}^2, 818.37 \text{ cm}^2$   
 12. (i)  $3.27 \text{ cm}^2$     (ii)  $109.77 \text{ cm}^2$     13.  $71.38 \text{ cm}^2$   
 14.  $0.43 \text{ cm}^2$     15.  $13.08 \text{ cm}^2, 439.08 \text{ cm}^2$     16. (i)  $230.79 \text{ m}^2$   
 (ii)  $765.69 \text{ m}^2$     (iii)  $410.29 \text{ m}^2$

∴ length of the radius of each circle = 7 cm.

$$\begin{aligned} \text{Now, area of the sector AXZ} &= \frac{60}{360} \pi \cdot 7^2 \\ &= \frac{49\pi}{6} \text{ cm}^2 \end{aligned}$$

$$\text{Similarly, area of the sector BYX} = \frac{49\pi}{6} \text{ cm}^2$$

$$\text{and area of the sector CZY} = \frac{49\pi}{6} \text{ cm}^2$$

$$\begin{aligned} \therefore \text{the total area of the three sectors} &= 3 \cdot \frac{49\pi}{6} \\ &= \frac{49\pi}{2} \text{ cm}^2 \end{aligned}$$


∴ the required area regions

$$= \text{Area of } \triangle ABC - \text{total area of the three sectors}$$

$$= 49\sqrt{3} - \frac{49\pi}{2}$$

$$= 49 \left( \sqrt{3} - \frac{\pi}{2} \right) \text{ cm}^2$$

**Example 8.** In Fig 12.13, AB and CD are two diameters of a circle (with centre O) perpendicular to each other and OA is the diameter of the smaller circle. If OA = 6 cm, find the total area of the shaded regions.



**Fig. 12.13**

**Solution :** Here OA = OB = OC = OD = 6 cm

Now, area of the shaded region I

$$= \text{area of the quadrant AOC} - \text{semi circle of smaller circle}$$

$$= \frac{1}{4} \times \pi \times 6^2 - \frac{1}{2} \times \pi \times 3^2$$

$$= 9\pi - \frac{9\pi}{2}$$

$$= \frac{9}{2} \pi \text{ cm}^2$$

Similarly, area of the shaded region II =  $\frac{9}{2} \pi \text{ cm}^2$ .

9. BC is the arc of a quadrant of a circle of radius 8cm. A semi - circle is described on the chord BC on the side opposite to the centre of the quadrant. Find the area enclosed between arc BC and the semi circle.
10. In a 400 m track the length of each of the straight portions is 90 m and the diameter of each of the inner semi-circular ends is 70 m. If the uniform width of the track is 7m, find the cost of turfing the track at the rate of ₹ 250 per square metre.
11. Two flower beds are in the form of segments (minor ) of the circle circumscribing a square field of side 42 m, with two opposite sides of square as the corresponding chords. Find the cost of planting flowers at the rate of ₹ 200 per square metre.
12. On a rectangular window of dimension, 84 cm × 56 cm, twentyfour nonoverlapping circular designs each of radius 7 cm are made at the rate of ₹ 25 per 20 cm<sup>2</sup>. Find the cost of making the designs. Also find the area of the remaining portion.
13. In a round table of diameter 112 cm, there is a portion in the form of a regular hexagon inscribed in it. Designs are made on the remaining portion of the table top at the rate of ₹ 8 per 15 cm<sup>2</sup>. Find the cost of designing (use  $\sqrt{3} = 1.73$ ).

14. Find the area and perimeter of the shaded region in the given Fig.12.14, having given that OA = OB = 28cm.




Fig. 12.14

15. In the given Fig. 12.15, AC = CB = 18cm. Find the area of the shaded region using the fact that

$$OP = \frac{1}{6} AB.$$




Fig. 12.15

16. In the given fig. 12.16, O is the centre and AOB is the diameter of the circle. Find the area of the shaded region.




Fig. 12.16

the total surface area of the cylinder

$$= \text{curved surface area} + \text{area of two circular ends}$$

$$= 2\pi rh + 2\pi r^2$$

$$= 2\pi r(h + r)$$

and the volume = area of the base  $\times$  height

$$= \pi r^2 \times h$$

$$= \pi r^2 h$$

If  $l$ ,  $r$  and  $h$  are respectively the slant height, radius of the base and height of a right circular cone, then  $l^2 = r^2 + h^2$ . Further, the total surface area

$$= \text{curved surface area} + \text{area of base}$$

$$= \pi rl + \pi r^2$$

$$= \pi r(l + r)$$

and the volume =  $\frac{1}{3} \pi r^2 h$

If  $r$  be the radius of sphere, then its surface area is  $4\pi r^2$  and its volume is  $\frac{4}{3} \pi r^3$ .

If  $r$  is the radius of a hemisphere, then its total surface area

$$= \text{curved surface area} + \text{area of circular face}$$

$$= 2\pi r^2 + \pi r^2$$

$$= 3\pi r^2$$

and volume =  $\frac{2}{3} \pi r^3$

Henceforth, cylinder and cone will mean right circular cylinder and right circular cone respectively, unless otherwise stated.

**Example 9.** A vessel is in the form of a hollow cylinder surmounted on a hemispherical bowl of the same radius. The diameter of the hemisphere is 12cm and the total height of the vessel is 14 cm. Find

- (i) the total surface area of the vessel
- (ii) the volume of the vessel
- (iii) the cost of painting the outer surface of the vessel at the rate of ₹ 5 per square cm. (take  $\pi = 3.14$ ).

$$= \frac{1}{3} \times 3.14 \times 3^2 \cdot 4$$

$$= 37.68 \text{ cm}^3.$$

and surface area of the right circular cone  $= \pi r l$

$$= 3.14 \times 3.5$$

$$= 47.1 \text{ cm}^2.$$

Also, volume of the hemisphere  $= \frac{2}{3} \pi r^3$

$$= \frac{2}{3} \times 3.14 \times 3^3$$

$$= 56.52 \text{ cm}^3$$

and surface area of the hemisphere  $= 2\pi r^2$

$$= 2 \times 3.14 \times 3^2$$

$$= 56.52 \text{ cm}^2$$

(i) The required volume of the toy

$$= \text{volume of the cone} + \text{volume of the hemisphere}$$

$$= 37.68 + 56.52 = 94.2 \text{ cm}^3$$

(ii) The required surface area of the toy

$$= \text{surface area of the cone} + \text{surface area of the hemisphere}$$

$$= 47.1 + 56.52 = 103.62 \text{ cm}^2.$$

**Example 11.** A canvas-tent is in the form of a cylinder of diameter 16 m and height 5 m surmounted by a cone of equal base and height 6 m. Find the capacity of the tent and the cost of the canvas at Rs 150 per square metre [use  $\pi = 3.14$ ].

**Solution :**

Here,

radius of the base of the cone ( $r$ )  
 $=$  radius of the cylinder ( $r$ )

$$= \frac{16}{2} = 8 \text{ m.}$$

and height ( $h$ )  $= 6$  m

$$\therefore \text{slant height } (l) = \sqrt{r^2 + h^2}$$

$$= \sqrt{8^2 + 6^2}$$

$$\begin{aligned}\therefore \text{curved surface area of the cylinder} &= 2\pi rh \\ &= 2 \cdot \frac{22}{7} \times 2.1 \times 4 \\ &= 52.8 \text{ m}^2.\end{aligned}$$

$$\begin{aligned}\text{and volume of the cylinder} &= \pi r^2 h \\ &= \frac{22}{7} \times 2.1 \times 4 \\ &= 55.44 \text{ m}^3.\end{aligned}$$

$$\begin{aligned}\text{and curved surface area of the hemispherical end} &= 2\pi r^2 \\ &= 2 \times \frac{22}{7} \times (2.1)^2 \\ &= 27.72 \text{ m}^2\end{aligned}$$

$$\begin{aligned}\text{and volume of the hemisphere} &= \frac{1}{2} \left( \frac{4}{3} \pi r^3 \right) \\ &= \frac{2}{3} \times \frac{22}{7} \times (2.1)^3 \\ &= 19.40 \text{ m}^3\end{aligned}$$

$$\begin{aligned}\therefore \text{the required total curved surface area} &= 52.8 + 27.72 \\ &= 80.52 \text{ m}^2.\end{aligned}$$

$$\begin{aligned}\text{and the required capacity of the boiler} &= 55.44 + 19.40 \\ &= 74.84 \text{ m}^3.\end{aligned}$$

#### EXERCISE 12.4

(Unless otherwise stated, use  $\pi = \frac{22}{7}$ )

1. A solid wooden toy is in the shape of a cone surmounted on a hemisphere of the same radius 2.1 cm. If the total height of the wooden toy is 8.1 cm, find its total surface area and volume.
2. A solid is in the form of a cylinder of diameter 7 cm with hemispherical ends. If the total length of the solid is 42 cm, find the total surface area and the volume of the solid.
3. An ice-cream is in the form of an inverted cone surmounted by a hemisphere of the same-radius. If the radius and the height of the cone are respectively 2.1 cm and 8 cm, find the volume of the ice-cream.



16. A godown is formed of a cuboid of dimensions  $63 \text{ m} \times 28 \text{ m} \times 5 \text{ m}$  covered by a half cylindrical roof. If the length and breadth of the cuboid are  $63 \text{ m}$  and  $28 \text{ m}$  respectively, find the volume of air inside the godown. Also find the cost of roofing at the rate of ₹ 500 per square metre.
17. A solid is in the form of a cylinder surmounted by a cone of the same radius. If the radius of the base and the height of the cone are ' $r$ ' cm and ' $h$ ' cm respectively and the total height of the solid is  $3h$ , prove that the volume of the solid is  $\frac{7}{3}\pi hr^2$ .

### ANSWERS

- |  |   |                                      |
|--|---|--------------------------------------|
| 1. $69.63 \text{ cm}^2$ , $47.12 \text{ cm}^3$   | 2. $924 \text{ cm}^2$ , $1527.16 \text{ cm}^3$    | 3. $56.36 \text{ cm}^3$              |
| 4. $24.75 \text{ cm}^3$                          | 5. 35   | 6. $2564.58 \text{ m}^3$ , ₹ 9,50,40 |
| 7. $5632 \text{ m}^3$ , ₹ 1,43,817.30            | 8. $678.86 \text{ m}^3$ , $310.39 \text{ m}^2$    |                                      |
| 9. 17.31 litres                                  | 10. $452.57 \text{ cm}^3$ , $320.57 \text{ cm}^2$ | 11. $8448 \text{ cm}^3$              |
| 12. $10560 \text{ cm}^3$                         | 13. 8.44 litres                                   | 14. $121.56 \text{ cm}^3$            |
| 15. $814.63 \text{ cm}^3$ , $475.2 \text{ cm}^2$ | 16. $28224 \text{ m}^3$ , ₹ 13,86,000             |                                      |

### 12.6 Conversion of solid from one shape to another

In this section, we shall discuss about the conversion of metallic solid into one or more solids of similar shape. For example, a solid metallic sphere is melted and recast into smaller spherical balls or recast into a wire; a solid metallic cone is melted and recast into the form of a solid cylinder etc. The calculation of surface areas and volumes in such cases will now be illustrated through examples.

We may note the following facts :

If  $r_1$  and  $r_2$  are respectively the inner and outer radii of a spherical shell (hollow sphere), then the volume of the material in the shell is  $\frac{4}{3}\pi(r_2^3 - r_1^3)$  cubic units

If  $r_1$  and  $r_2$  are respectively the internal and external radii of a cylindrical shell (hollow cylinder) of height  $h$ , then the volume of the material in the shell =  $\pi h(r_2^2 - r_1^2)$  cubic units.

**Example 13.** How many 5 cm solid cubes can be cut from a solid metallic cuboid measuring  $10 \text{ cm} \times 15 \text{ cm} \times 25 \text{ cm}$  ?

**Example 15.** Three solid metallic cones of the same height 18 cm and the same base radius 4 cm are melted together and recast into a solid sphere. Find the surface area of the sphere.

**Solution :** For the solid cone,  
 $r = 4$  cm,  $h = 18$  cm

$$\begin{aligned}\therefore \text{volume} &= \frac{1}{3} \pi r^2 h \\ &= \frac{1}{3} \pi \times 4^2 \times 18 \\ &= 96\pi \text{ cm}^3\end{aligned}$$

$$\begin{aligned}\therefore \text{volume of 3 such cones} &= 3 \times 96 \pi \\ &= 288 \pi \text{ cm}^3\end{aligned}$$

Let  $R$  be the radius of the sphere. Then its volume  $= \frac{4}{3} \pi R^3$

$$\text{By question, } \frac{4}{3} \pi R^3 = 288\pi$$

$$\begin{aligned}\Rightarrow R^3 &= \frac{3 \times 288}{4} \\ &= 216\end{aligned}$$

$$\therefore R = 6 \text{ cm}$$

$$\begin{aligned}\therefore \text{the surface area of the sphere} &= 4\pi r^2 \\ &= 4\pi \cdot 6^2 \\ &= 144\pi \text{ cm}^2\end{aligned}$$

**Example 16.** A solid metallic cylinder of height 24 cm and radius 3 cm is melted and recast into a cone of radius 6 cm. Find the height of the cone.

**Solution :** For the solid cylinder,

$$h = 24 \text{ cm, } r = 3 \text{ cm}$$

$$\begin{aligned}\therefore \text{volume of the cylinder} &= \pi r^2 h \\ &= \pi \cdot 3^2 \cdot 24 \\ &= 216\pi \text{ cm}^3\end{aligned}$$

For the solid cone,

$$r = 6 \text{ cm, } h = ?$$

**EXERCISE 12.5**

1. A solid cube is cut into two cuboids of equal volumes. Find the ratio of the total surface area of the given cube to that of one of the cuboids.
2. A sphere has the same surface area as the curved surface of a cone of height 16 cm and base radius 12 cm. Find the radius of the sphere.
3. A solid metallic cone is 81 cm high and radius of its base is 6 cm. If it is melted and recast into a solid sphere, find the curved surface area of the sphere.
4. A solid metallic sphere of diameter 28 cm is melted and recast into a number of cones, each of diameter 7 cm and height 4 cm. Find the number of cones so formed.
5. The internal and external radii of a hollow metallic sphere are 3 cm and 5 cm respectively. If the sphere is melted and recast into a solid cylinder of height  $2\frac{2}{3}$  cm, find the curved surface area of the cylinder.
6. A metallic cone of height 28 cm and radius of base 12 cm, is melted and recast into a cylinder of height  $9\frac{1}{3}$  cm. Find the curved surface area of the cylinder so formed.
7. 20 circular plates, each of radius 7 cm and thickness 1.5 cm are placed one above another to form a solid cylinder. Find the curved surface area and the volume of the cylinder so formed.
8. Find the volume of the largest sphere that can be carved out of a cube of side 4.2 cm. Also find the ratio of the volume of the cube to that of the sphere.
9. Find the volume of the largest cone that can be carved out of a cube of side 16.8 cm.
10. A solid cone of diameter 14 cm and height 8 cm is melted and recast into a hollow sphere. If the external diameter is 10 cm, find the internal diameter of the sphere.
11. A hemisphere of lead of radius 7 cm is cast into a cone of base radius 3.5 cm. Find the height of the cone.
12. A cylinder is cut lengthwise and flattened, thereby forming a rectangle of dimensions 66 cm  $\times$  28 cm. Find the curved surface area and volume of the original cylinder.
13. A sector of a circle of radius 6 cm has angle  $120^\circ$ . It is rolled up so that two bounding radii are joined together to form a cone. Find the volume of the cone.  
(Take  $\sqrt{2}=1.4$ )
14. A conical flask of radius  $x$  units and height  $2x$  units, is full of water. The water is poured into a cylindrical flask of radius  $\frac{2x}{3}$  units. Find the height of water in the cylindrical flask.


## 12.7 Frustum of a Right Circular Cone

In this section, you will learn to derive the formulas for finding the volume and the surface area of a frustum of a right circular cone. In our daily life, we use a number of solids of such form e.g. a bucket, a glass tumbler etc. The study of the volume and surface area of such type of solids will be very useful.

### Recall:

If a right circular cone is cut off by a plane parallel to the base, the portion of the cone between the plane and the base of the cone is called a **frustum** of the cone.

Let us now consider the following figures:-




**Fig. 12.18 (i)**

**Fig. 12.18 (ii)**

Of the two parts shown in Fig 12.18 (ii) the lower one is the frustum of the cone. It has two parallel flat circular bases namely base I and base II and a curved surface between the bases.

A section of the frustum by any plane containing its axis i.e. the line through the centres of the bases, is an isosceles trapezium ABCD in Fig 12.18(iii). The length of any of the pair of non parallel sides AD and BC is the slant height of the frustum.



**Fig. 12.18(iii)**

Also, the slant height of the frustum equals the difference between the slant heights of the cones OAB and OCD [Fig. 12.18(i)].

Also, the height of the frustum is equal to the difference between the heights of the cones OAB and OCD.

$$\begin{aligned}
 &= \frac{1}{3} \pi h \left( \frac{r_1^3 - r_2^3}{r_1 - r_2} \right) \\
 &= \frac{1}{3} \pi h \frac{(r_1 - r_2)(r_1^2 + r_1 r_2 + r_2^2)}{(r_1 - r_2)} \\
 &= \frac{1}{3} \pi h (r_1^2 + r_1 r_2 + r_2^2)
 \end{aligned}$$

Thus, the volume V of the frustum of cone is given by

$$V = \frac{1}{3} \pi h (r_1^2 + r_1 r_2 + r_2^2)$$

The volume may also be expressed in terms of the area of bases and the height.

If  $A_1$  and  $A_2$  ( $A_1 > A_2$ ) are the surface areas of the two circular bases, then

$$A_1 = \pi r_1^2$$

and  $A_2 = \pi r_2^2$ .

Thus, the volume of the frustum of cone is given by

$$\begin{aligned}
 V &= \frac{1}{3} h [\pi r_1^2 + \pi r_1 r_2 + \pi r_2^2] \\
 &= \frac{1}{3} h \left[ \pi r_1^2 + \pi r_2^2 + \sqrt{(\pi r_1^2)(\pi r_2^2)} \right] \\
 &= \frac{1}{3} h \left[ A_1 + A_2 + \sqrt{A_1 A_2} \right]
 \end{aligned}$$

Now from the right  $\triangle CAN$ , right  $\angle$  ed at N, we have

$$AN = r_1 - r_2$$

$$\text{and } AC^2 = l^2 = h^2 + (r_1 - r_2)^2$$

$$\therefore \text{slant height } l = \sqrt{h^2 + (r_1 - r_2)^2}$$

Let  $l_2$  be the slant height of the cone OCD

Since  $\triangle OMC \sim \triangle OLA$ , therefore

$$\frac{l_2}{l_1} = \frac{r_2}{r_1}$$

$$\Rightarrow \frac{l_1 - l}{l_1} = \frac{r_2}{r_1}$$

This gives,  $l_1 = \frac{l r_1}{r_1 - r_2}$  .....(3)

$$\begin{aligned}
 &= \frac{1}{3} \times 3.14 \times 8 \times 201 \\
 &= 1683.04 \text{ cm}^3
 \end{aligned}$$

$$\begin{aligned}
 \text{Slant height, } l &= \sqrt{h^2 + (r_1 - r_2)^2} \\
 &= \sqrt{8^2 + 6^2} \quad [\because r_1 - r_2 = 11 - 5 = 6] \\
 &= \sqrt{100} \\
 &= 10
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{ curved surface area} &= \pi l (r_1 + r_2) \\
 &= 3.14 \times 10 (11 + 5) \\
 &= 3.14 \times 10 \times 16 \\
 &= 502.4 \text{ cm}^2.
 \end{aligned}$$

**Example 19.** A bucket is in the form of a frustum of a cone whose top and bottom are of diameters 42 cm and 28 cm respectively. If the height of the bucket is 21 cm, find the capacity of the bucket in litres.

**Solution :** We have

$$r_1 = 21 \text{ cm, } r_2 = 14 \text{ cm.}$$

$$h = 21 \text{ cm.}$$

$$\begin{aligned}
 \therefore \text{ capacity of the bucket} &= \frac{1}{3} \pi h [r_1^2 + r_2^2 + r_1 r_2] \\
 &= \frac{1}{3} \times \frac{22}{7} \times 21 [21^2 + 14^2 + 21 \cdot 14] \\
 &= 22 [441 + 196 + 294] \\
 &= 22 \times 931 \\
 &= 20482 \text{ cm}^3 \\
 &= \frac{20482}{1000} \text{ litres} \\
 &= 20.48 \text{ litres.}
 \end{aligned}$$

**Example 20.** A container made up of a metal sheet is in the form of a frustum of a cone of height 20 cm with radii of its upper and lower ends as 40 cm and 25 cm respectively. Find the quantity of milk the container can hold . Also find the area of the metal sheet used for making the container. ( $\pi = 3.14$ ).

$$\therefore \frac{OM}{OL} = \frac{MD}{LB}$$

$$\Rightarrow \frac{h_2}{h_1} = \frac{r_2}{r_1} \dots\dots\dots(1)$$

Also,

$$\frac{\text{volume of the smaller cone}}{\text{volume of the given cone}} = \frac{1}{27}$$

$$\Rightarrow \frac{\frac{1}{3} \pi r_2^2 h_2}{\frac{1}{3} \pi r_1^2 h_1} = \frac{1}{27}$$

$$\Rightarrow \left(\frac{r_2}{r_1}\right)^2 \cdot \left(\frac{h_2}{h_1}\right)^2 = \frac{1}{27} \quad \text{[by (1)]}$$

$$\Rightarrow \left(\frac{h_2}{h_1}\right)^3 = \left(\frac{1}{3}\right)^3$$

$$\Rightarrow \frac{h_2}{h_1} = \frac{1}{3}$$

$$\Rightarrow h_2 = \frac{1}{3} \times h_1$$

$$= \frac{1}{3} \times 18 = 6$$

$$\therefore \text{the height of the resulting frustum} = h_1 - h_2 = 18 - 6 = 12 \text{ cm.}$$




Fig. 12.21

**EXERCISE 12.6**

(unless otherwise stated, use  $\pi = \frac{22}{7}$ )

1. If the radii of the circular ends of frustum of height 6 cm are 20 cm and 12 cm respectively, find the volume and the curved surface area of the frustum. (Take  $\pi = 3.14$ )
2. A bucket is in the form of a frustum of a cone. If the height of the bucket is 16 cm and the radii of the upper and lower ends are 18 cm and 6 cm respectively, find

13. A cone is cut into three parts by planes through the points of trisection of its altitude and parallel to the base. Prove that the volumes of the parts are in the ratio 1: 7:19.
14. A bucket is in the form of a frustum with a capacity of 45584 cubic cm. If the radii of the top and bottom of the bucket are 28 cm and 21cm respectively, find its height and surface area.

### ANSWER

1.  $4923.52 \text{ cm}^3, 1004.8 \text{ cm}^2$

2.  $24\text{cm}, 7844.57 \text{ cm}^2, 20 \text{ cm}, 1621.71\text{cm}^2$

3.  $65.76 \text{ litres}$

4.  $3072.3 \text{ cm}^3$

5.  $301.57 \text{ cm}^3.$

6.  $14589.14 \text{ cm}^2$

7. ₹  $122.80, 1177.5 \text{ cm}^2$

8.  $1: 7$

9.  $2:3$

10.  $7.5 \text{ cm}.$

12.  $8 \text{ cm}$

14.  $24 \text{ cm}, 5236 \text{ cm}^2$

### SUMMARY

In this chapter, you have studied the following points :

1. Circumference of a circle =  $2\pi r$
2. Area of a circle =  $\pi r^2$
3. Length of an arc of a sector of a circle with radius  $r$  and angle  $\theta$  (measure in degrees) is  $\frac{\theta}{360} \times 2\pi r$
4. Area of a sector of circle with radius  $r$  and angle  $\theta$  (measured in degrees) is  $\frac{\theta}{360} \times \pi r^2$ .
5. Area of segment of a circle = Area of the corresponding sector  
– Area of the corresponding triangle.



### 13.1 Introduction

In your previous classes, you have seen and studied Statistics as a subject dealing with collection, compilation, analysis and interpretation of data. We can also look upon it as the weighing of evidence, meaning thereby that standard statistical methods give quantified evidence to draw certain conclusions in respect of one or more characteristics of the data (population).

It has also been stated that statistics gives information which are of representative nature and do not pertain to information on particular individual or individuals.

For example, the per capita income of India has statistical importance to represent the income of an average Indian while the income of the richest person of the country has less importance to represent the income of a typical Indian.

The main parameters of a population that a statistician desires to know in order to draw conclusions on it are the measures of central tendency and location, deviation, skewness and kurtosis of the data. These are the five important indicators whose knowledge promote a systematic study of the population.

In fact they are the five most important parameters indicating the state of being of the population in respect of a characteristic or characteristics.


As an analogy, let us consider the case of a medical practitioner. A doctor generally wants to know the blood pressure reading, the number of heart beats per minute, the body mass index etc. of a person in order to study the existing conditions of some of the vital organs of the person. Actually, they are important indicators of the state of health of the person under investigation similar to the parameters of a population under study.

In elementary classes of statistics we shall devote our attention to the first parameter namely, the central tendency of the data and a little to the measures of location.

What we are to study in the following sections of this chapter are the methods and devices which are generally adopted to determine the measures of central tendency of the data and also their representation and interpretation in numerical and graphical forms.

Before we proceed to study the methods and devices for the estimation of certain parameters of a population, a brief discussion on the innate character of the subject Statistics is worth mentioning.

**In the case of other physical sciences:**



Now, you have seen both the routes and must have observed that one is the other way round of the other.

However, it is noteworthy to mark that empirical findings are the forerunners of almost all the conceptually established theories. History of scientific discoveries and inventions are replete with many anecdotes of keen observations leading to profound theories.

One such example is John Kepler's laws of planetary motion derived from purely observational data and the same being profitably used by Sir Isaac Newton leading to the discovery of his famous laws of gravitation.

### 13.2 Measures of Central Tendency

You have seen that one of the major objectives of the study of data, particularly numerical data is to find the measures of central tendency of the data. However, we should note that there is no unique name for the measure of central tendency. Depending on the nature of the data under study, we may take the mean or the median or the mode to represent the central tendency. In any case, a measure of central tendency is the value of the variate around which the other values are supposed to cluster.

We shall first describe the most commonly used measures of central tendency.

### 13.3 The Weighted Arithmetic Mean (AM)

For constructing a grouped frequency distribution, we make classes of the variate values into which all the observations are packed piecewise.

For example, when we consider the grouped frequency distribution of marks obtained by 250 students in an examination of 100 marks, suppose we write 10-20 in the column of classes and 35 in the corresponding column of frequencies against the class, we mean that the number of students securing marks 10 and above but less than 20 is 35.

Then the total number of persons =  $f_1 + f_2 + \dots + f_n = N$  and the total amount with all the persons =  $f_1x_1 + f_2x_2 + \dots + f_nx_n$ .

Hence, the average amount with a person denoted by  $\bar{x}$  is given by

$$\bar{x} = \frac{f_1x_1 + f_2x_2 + \dots + f_nx_n}{f_1 + f_2 + \dots + f_n}$$

$$\text{or, } \bar{x} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i} = \frac{1}{N} \sum_{i=1}^n f_i x_i$$

**Note 2.** The term weighted is used because of the analogy of the formula for  $\bar{x}$  and that of the distance of the centre of gravity of a system of weights  $w_1, w_2, \dots, w_n$  at respective distances of  $x_1, x_2, \dots, x_n$  from the origin along the same line. The distance  $\bar{x}$  of the centre of gravity of the system of weights from the origin is given by

$$\bar{x} = \frac{x_1w_1 + x_2w_2 + \dots + x_nw_n}{w_1 + w_2 + \dots + w_n}$$

The weights are comparable to the frequencies of the various classes.

For a grouped frequency distribution  $(f_i, x_i)$ ,  $i = 1, 2, \dots, n$ , two more means as measures of central tendency are used in statistics. They are the weighted Geometric Mean (GM) and the weighted Harmonic Mean (HM). They are respectively denoted by G and H

and are given by the formulae.  $G = (x_1^{f_1} \cdot x_2^{f_2} \cdot \dots \cdot x_n^{f_n})^{\frac{1}{f_1 + f_2 + \dots + f_n}}$

$$\text{or, } G = \left( \prod_{i=1}^n x_i^{f_i} \right)^{\frac{1}{N}}$$

$$\text{and } \frac{(f_1 + f_2 + \dots + f_n)}{H} = \frac{f_1}{x_1} + \frac{f_2}{x_2} + \dots + \frac{f_n}{x_n}$$

$$\text{i.e. } \frac{N}{H} = \sum_{i=1}^n \frac{f_i}{x_i}$$

$$\text{Or, } \frac{1}{H} = \frac{1}{N} \sum_{i=1}^n \frac{f_i}{x_i} \quad ; \quad N = \sum_{i=1}^n f_i \text{ in both the cases.}$$

**Solution :** The data is reproduced with the mid values of the classes

Classes	Mid value	Frequency	
	$x_i$	$f_i$	$f_i x_i$
0 – 10	5	55	275
10 – 20	15	57	855
20 – 30	25	80	2000
30 – 40	35	75	2625
40 – 50	45	62	2790
50 – 60	55	47	2585
60 – 70	65	25	1625
70 – 80	75	7	525
80 – 90	85	2	170
90–100	95	0	0

$$410=N \quad 13450=\sum f_i x_i$$

**(a) By Direct method**

The mean age  $\bar{x}$  is given by  $\bar{x} = \frac{1}{N} \sum_{i=1}^{10} f_i x_i$

$$\therefore \bar{x} = \frac{1}{410} \times 13450 = 32.8 \text{ years (nearly).}$$

Here ,  $\sum_{i=1}^{10} f_i = 410 = N$  and,

$$\sum_{i=1}^{10} f_i x_i = 13450.$$

**Note:** Since the frequency of the last class is 0, we can drop the class and take only 9 classes. This will not affect the value of  $\bar{x}$ . If 0 frequency occurs anywhere in the table, then also the class can be dropped without affecting the value of the mean. However, retaining it gives the information that there is no variate assuming that value or there is no observation in the corresponding class.

**(c) The step deviation or the change of origin and scale method**

In the solution of the problem in the above example by the assumed mean method if you examine the column of the deviation i.e.  $d_i$ , you will observe that these deviations have the greatest common factor 10, in this case. In fact 10 is the width of each of the classes.

Thus, when the deviations have common factors, the HCF i.e. the greatest common factor say,  $h$  can be easily found and we construct a new table using  $u_i = \frac{x_i - a}{h}$  as the new variate.

Thus taking  $a = 55$  and  $h = 10$  the new table is constructed as follows:

$x_i$	$u_i = \frac{x_i - a}{h} \left( = \frac{x_i - 55}{10} \right)$	$f_i$	$f_i u_i$
5	-5	55	-275
15	-4	57	-228
25	-3	80	-240
35	-2	75	-150
45	-1	62	-62
55	0	47	0
65	1	25	25
75	2	7	14
85	3	2	6
95	4	0	0

410

$$\begin{aligned} \sum f_i u_i &= -955 + 45 \\ &= -910 \end{aligned}$$

Now,  $u_i = \frac{x_i - a}{h}$

$$\Rightarrow hu_i = x_i - a$$

$$\Rightarrow f_i hu_i = f_i x_i - af_i$$

$$\Rightarrow \sum f_i hu_i = \sum f_i x_i - \sum af_i$$

$$\Rightarrow h \sum f_i u_i = \sum f_i x_i - a \sum f_i$$

$$\Rightarrow \frac{h}{N} \sum f_i u_i = \frac{1}{N} \sum f_i x_i - \frac{a}{N} \sum f_i$$

$$\Rightarrow h\bar{u} = \bar{x} - a$$

$$\Rightarrow \bar{x} = a + h\bar{u}, \bar{u} \text{ is the mean of the new variate } u \text{ whose values are } u_i \text{ (} i = 1, 2, \dots, n \text{)}$$

Here,  $N = 410$

$$\therefore \frac{N}{2} = 205.$$

From the cumulative frequency column we observe that 192 persons have their ages below 30 years. And 267 persons have their ages below 40. Thus, the age such that 205 persons are younger and another 205 are older than it must be between 30 and 40.

Thus, 30 – 40 is the median class.

$\therefore$  Median age =  $M = 30 +$  a fraction of the class width 10.

We assume that the frequency 75 of the median class is uniformly distributed over the width i.e. 10

of the class, so that the remaining portion  $\left(\frac{410}{2} - 192\right)$  must lie in a width of  $\frac{10}{75}(205 - 192)$

$$\begin{aligned} \therefore M &= 30 + \frac{10}{75} \times 13 \\ &= 31.7 \text{ years (nearly).} \end{aligned}$$

Thus, we use the formula

$$M = l + \frac{\frac{N}{2} - c}{f} \times h$$

where,  $l$  = lower limit of the median class

$N$  = size of the sample or population

$c$  = cumulative frequency of the class just before the median class

$f$  = frequency of the median class

$h$  = width of the median class

**Note : The given data can be modified in the following ways**

1.

Age in years	Number of persons
Below 100	410
' 90	410
' 80	408
' 70	401
' 60	376
' 50	329
' 40	267
' 30	192
' 20	112
' 10	55

$$\begin{aligned}
 \therefore \text{Mode} &= 20 + \frac{80 - 57}{2 \times 80 - 57 - 75} \times 10 \\
 &= 20 + \frac{23}{28} \times 10 \\
 &= 20 + 8.2 \\
 &= 28.2 \text{ years (nearly)}
 \end{aligned}$$

**Note : In an ideal case, the mean, median and mode of a distribution coincide at the same point.**

However, a rough test for consistency can be done by using Pearson's empirical formula viz.

Mean – Mode = 3 (Mean – Median) even though they do not coincide.

$$\begin{aligned}
 \text{In our case, Mean – Mode} &= 32.8 - 28.2 \\
 &= 4.6
 \end{aligned}$$

$$\begin{aligned}
 \text{and, } 3(\text{Mean – Median}) &= 3(32.8 - 31.7) \\
 &= 3.3
 \end{aligned}$$

Thus, the difference is not large.

### 13.7 Measures of Location or Partition values or Quantiles

For a grouped frequency distribution, certain values of the variate may give important information about the data. For example, we may be interested in the value of the variate such that it is more than one fourth of all the values of the variate in the data while is less than three fourths of all of them etc.

Such values are called the partition values or quantiles or measures of location because they are partitioning the whole data into parts.

Some of the important partition values are :

#### The First or Lower Quartile

This quartile denoted by  $Q_1$  is the value of the variate such that one fourth of all the values of the variate in the data have their values less than or equal to it while three fourths have their values greater than or equal to it and  $Q_1$  is given by

$$Q_1 = l + \frac{\frac{N}{4} - c}{f} \times h$$

**Example 2.** For the data of section 13.4, the first quartile  $Q_1$  and the third quartile  $Q_3$  are calculated as follows :

$$\text{Now, } \frac{N}{4} = \frac{410}{4} = 102.5$$

Here, 10 – 20 is the  $Q_1$  class

$$\begin{aligned} \therefore Q_1 &= 10 + \frac{102.5 - 55}{57} \times 10 \\ &= 10 + 8.33 \\ &= 18.3 \text{ (nearly)} \end{aligned}$$

$$\text{Again, } \frac{3}{4} N = \frac{3}{4} \times 410 = 307.5$$

Here, 40 – 50 is the  $Q_3$  class

$$\begin{aligned} \therefore Q_3 &= 40 + \frac{307.5 - 267}{62} \times 10 \\ &= 40 + 6.53 \\ &= 46.5 \text{ (nearly)}. \end{aligned}$$

In the same way,  $D_6$ , the 6<sup>th</sup> decile or the 60<sup>th</sup> percentile of the data is given by

$$D_6 = P_{60} = l + \frac{\frac{60}{100} N - c}{f} \times h, \text{ with usual notations}$$

$$\text{Now, } \frac{60}{100} N = \frac{6}{10} \times 410 = 246$$

30 – 40 is thus the  $P_{60}$  class

$$\begin{aligned} \therefore P_{60} &= 30 + \frac{\frac{60}{100} N - 192}{75} \times 10 \\ &= 30 + \frac{246 - 192}{75} \times 10 \\ &= 30 + 7.2 \\ &= 37.2 \text{ years} \end{aligned}$$

### 13.8 Applications of the measures of location

Although the measures of locations are simple as they look, they have many applications in taking important decisions objectively. Now a-days the results of public examinations, particularly competitive exams where a large number of students appear for limited number of seats, are declared with percentiles. These results enable the examining



Group of marks secured (classes)	No. of candidates (Frequency)	Cumulative frequency
0 - 10	5606	5606
10 - 20	8670	14276
20 - 30	10078	24354
30 - 40	25686	50040
40 - 50	25700	75740
50 - 60	35006	110746
60 - 70	24678	135424
70 - 80	16462	151886
80 - 90	4270	156156
90 - 100	1054	157210

Since the admission is to be given only to the uppermost 10%, we are to find the marks secured by the candidates such that 90% of the total number of candidates get marks below that mark and 10% get it and above it. Surely, it is  $P_{90}$ , i.e. the 90<sup>th</sup> percentile.

$$\text{Now, } N = 157210 \text{ and } \frac{90}{100}N = \frac{90}{100} \times 157210$$

$$= 141489$$

Here, 70-80 is the  $P_{90}$  class

$$\therefore \text{ From the formula, } P_{90} = l + \frac{\frac{90}{100}N - c}{f} \times h$$

we get,

$$P_{90} = 70 + \frac{141489 - 135424}{16462} \times 10$$

$$= 70 + \frac{60630}{16462}$$

$$= 74 \text{ (nearly).}$$


Thus, those candidates who secure 74 marks and above can be given admission.

Now, each and every student gets a unique mark between 0 and 100. Again, we can also find all the percentiles from  $P_1$  to  $P_{99}$ .  $P_0$  and  $P_{100}$  are surely 0 and 100 respectively in this case.

Thus a candidate getting any mark between 0 to 100 will have a percentile which indicates his/her relative position with respect to the total number of candidates.

To draw the more than ogive, we plot points with the lower limits of the classes as the abscissae and the more than cumulative frequency of the corresponding classes as the ordinates and join these points by free hand.

Thus, we plot the points (0,410), (10,355), (20,298), (30,218), (40,143), (50,81), (60,34), (70,9), (80,2), (90,0) and joining them by free hand we get the more than ogive.



The two ogives intersect at the point 'P' whose abscissa is more than 30. Thus  $M > 30$ .

**Note.1** Accurate values can be obtained by taking precise scale. We are showing only the principle.

**Note.2** You must have seen that the Median and the second Quartile are both measures of central tendency and measures of location.

**13.10 A general remark on the suitability of the different measures of central tendency**

We have seen that there are three measures of central tendency viz, the mean, the median and the mode.

Again, there are three means viz, the AM, the GM and the HM.

It is therefore quite natural to seek the criteria for the determination of the most suitable measure of central tendency for a particular case. For such situation, we examine the case in the following way.

Month	No. of T.V. sets sold
January	14
February	17
March	16
April	12
May	7
June	6
July	8
August	9
September	6
October	14
November	15
December	18

Find the average number of T.V. sets sold per month.

3. The following is the record of weight of children at the time of their birth as maintained in a maternity ward during a particular year--

Weight of children in kg.	No. of children
2.5	12
3.0	175
3.5	156
3.8	42
4.0	15
4.2	5

Find the average (mean) weight of a child at the time of birth.

4. Five coins were simultaneously tossed 1000 times, and at each toss the number of heads was observed. The number of tosses during which 0, 1, 2, 3, 4, 5 heads were obtained are shown in the table below. Find the mean number of heads per toss.

No of heads per toss	No. of tosses
0	38
1	144
2	342
3	287
4	164
5	25

Expenditure on water (in Rs.)	No. of families
30 – 40	12
40 – 50	18
50 – 60	20
60 – 70	15
70 – 80	12
80 – 90	11
90 – 100	6
100 – 110	4
110 – 120	2

**100**

Find the mean monthly expenditure of the families on water .

Also, determine the quartiles of the expenditure.

8. Find the median, lower and upper quartiles, 4th decile and 56<sup>th</sup> percentile of the following distribution of marks:

Marks	No. of students
0 – 4	10
4 – 8	12
8 – 12	18
12 – 14	7
14 – 18	5
18 – 20	8
20 – 25	4
25 and above	6

Daily savings (in ₹ )	No. of workers
1 – 2.99	6
3 – 4.99	53
5 – 6.99	85
7 – 8.99	56
9 – 10.99	21
11 – 12.99	16
13 – 14.99	4
15 – 16.99	4

[Hint : Convert into continuous classes]

### ANSWER

(1) 2.63      (2) 11.83      (3) 3.31 Kg      (4) 2.47

(5) 4.75      (6) Mean = 25.04, Median = 23.82

(7) Mean = ₹ 63.60,  $Q_1 = ₹ 47.22$ ,  $Q_3 = ₹ 78.33$

(8) Median = 10.89,  $Q_1 = 6.5$ ,  $Q_3 = 18.13$ ,  $D_4 = 9.33$ ,  $P_{56} = 11.82$

(9) 56.56

(10) 

Class	Frequency
20 - 40	28
60 - 80	24

(11) Median = 59.35,  $Q_1 = 31.04$ ,  $Q_3 = 68.52$ ,

Mode = 66.77

(12) Median = ₹ 6.49,  $Q_1 = ₹ 5.05$ ,  $Q_3 = ₹ 8.41$ ,

$D_8 = ₹ 8.85$ ,  $P_{56} = ₹ 6.84$ , Mode = ₹ 6.04

### 14.1 Introduction

We are familiar with the word probability. In fact, we have seen that it is the measure of chance and the probability that we are discussing is objective probability. We have also seen that the probability of the occurrence of an event is a real number lying between 0 and 1. In extreme cases, the end values may be assumed. In other words, for any event  $E$ , the probability  $P(E)$  of its occurrence is such that  $0 \leq P(E) \leq 1$ . You may note that the probability of an event may be an irrational number as you will come across later on (see Appendix II).

The definition of this measure of chance is given in three seemingly different forms and you are familiar with the one called the empirical or experimental definition. One of the other two is the classical or mathematical or a priori definition of probability due to Laplace and the last is the set theoretic or axiomatic or modern definition due to A. Kolmogorov.

We shall not discuss the last definition. This is discussed in higher classes with more sophisticated tools of modern Mathematics.

However, the value of the probability of an event  $E$  as calculated from these three seemingly different stand points converge to the same value as the number of trials becomes larger and larger.

In the following few sections, we shall discuss probability of the occurrence of an event from the classical or mathematical approach.

But before the actual discussion, we shall consider some terms associated with the probability of an event.

### 14.2 Random or non-deterministic experiment

An experiment, whose result cannot be uniquely predicted even if the previous results of the same experiment conducted under similar conditions are all known is called a random or more precisely a non-deterministic experiment.

For example, tossing a coin 100 times may give 51 heads and 49 tails. With this prior information, we cannot say the outcome of the 101<sup>th</sup> toss beforehand. In our daily social and personal lives, we come across a number of such examples and some of them are so important that the fate of some individuals or even a powerful government may depend on the outcome of such unpredictable phenomena.

Take the case of monsoon in India. How important is this phenomenon to the life of individuals and the government of the country! With all the records of the previous years, nobody can say precisely the shape of things to come in the coming years. Such is the nature of a non-deterministic experiment.

A non-deterministic experiment is also sometimes known as a trial. It is to be noted that one or more trials may constitute an experiment.

### 14.3 Sample space and event

The totality of all the possible outcomes of an experiment is called the sample space of the experiment. Any component of a sample space is an event.

For example, in tossing a coin once, the possible outcomes are a head or a tail to be denoted by H and T respectively. If we take these two outcomes together and write S to denote the collection of these two denoted by  $\{H, T\}$ , i.e.  $S = \{H, T\}$ , then S is the sample space of the experiment of tossing a coin once.

In this case, to get H or to get T are events and they are denoted by  $\{H\}$  and  $\{T\}$  respectively. Note that H means a head while  $\{H\}$  means the event of getting a head. The cases of getting none of them and getting any one of H and T are also events. In fact, for the experiment of tossing a coin, there are 4 events. Similarly, the sample space S of tossing two coins once or one coin twice is given by  $S = \{HH, HT, TH, TT\}$ .

You can verify that for this there are 16 events (How?).

Now what does  $\{HH\}$  represent? It is the event of getting head in both the tosses. Similarly,  $\{HH, HT, TH\}$  represents the event of getting at least one head in tossing two coins once or one coin twice.

### 14.4 Equally likely events

Events are said to be equally likely if there is no valid reason to say that one event has more chance to occur than the others.

For example, in tossing an unbiased coin, one of the faces will turn up but there is no valid reason to say that a particular face will surely turn up or has more chance to turn up.

### 14.5 Mutually exclusive events

Events are said to be mutually exclusive if the happening of one forecloses the happening of all the others.

For example, in tossing a fair die once, one of the six faces will turn up. But when a face, say, the one denoted by [2] or [· ·] turns up, any one of the remaining five faces namely [1],[3],[4],[5] and [6] or [·],[··][· ·],[· · ·],[· · ·] will not turn up.

#### 14.6 Independent events

Events are said to be independent if the occurrence of one has no effect on the occurrence of the other or others.

For example : If a coin is tossed twice, the event of getting H or T in the second toss is independent of the outcome of the first toss.

#### 14.7 Elementary events

An elementary event is one which cannot be further subdivided. This is analogous to an atom which is the smallest unit that is taking part in a chemical reaction.

Therefore, some writers use the term 'atom' of an event to represent an elementary event.

e.g. In the tossing of a coin once, the events {H} and {T} are elementary events.

On the other hand, in the experiment of tossing a coin twice, the event E of getting at least one head consists of the three mutually exclusive events viz {HH}, {HT} and {TH}. These constitute the event E. Thus, E is not an elementary event.

Further, any one of the three events is not an elementary event because, the event say {HT} is formed by two elementary events namely {H} in the 1st and {T} in the second toss. Similarly, {TT} is the event of getting T in both the tosses.

#### 14.8 Exhaustive set of events

A set of events is said to be exhaustive if all the possible outcomes are included.

For example, in tossing a fair die once, the six events {1},{2},{3},..., {6} constitute a set of exhaustive events.

#### 14.9 Favourable outcomes

For every experiment, out of the set of exhaustive outcomes, those entailing the occurrence of a particular event are called the favourable ones for the event.

For example, consider the following situation.

In an urn, there are 2 white and 4 black balls, all similar in all respects except in colour. One ball is drawn at random.



If E is the event of drawing a white ball, then, out of the six exhaustive ways of drawing, only two are favourable to the event E.

#### 14.10 Classical or Mathematical or a Priori definition of Probability due to Laplace

Out of  $n$  exhaustive, equally likely and mutually exclusive outcomes, if  $m$  are favourable to the event A, then the probability of the occurrence of the event A denoted by  $P(A)$  is the ratio  $m:n$  and we write  $P(A) = \frac{m}{n} = \frac{\text{Number of outcomes favourable to A}}{\text{Total number of possible outcomes}}$

From the definition, it is clear that  $0 \leq P(A) \leq 1$ .

When the probability of an event is 0, we say that the event will never occur. On the other hand, if the probability of an event is 1, then we say that the event is sure to happen.

Conversely, an impossible event has its probability 0 and a sure event has its probability 1.

We use the symbol  $P(\bar{A})$  to denote the probability of not happening of the event A.

In the definition, out of  $n$  exhaustive, equally likely and mutually exclusive outcomes,  $(n-m)$  are not favourable to the event A.

$$\therefore P(\bar{A}) = \frac{n-m}{n} = 1 - \frac{m}{n} = 1 - P(A)$$

$$\text{or, } P(A) + P(\bar{A}) = 1.$$

A and  $\bar{A}$  are called complementary events of each other. You are to note that A and  $\bar{A}$  are mutually exclusive events and one of them is sure to happen i.e. A and  $\bar{A}$  form an exhaustive system of events.

Similarly, if A, B, C are three mutually exclusive events forming an exhaustive system then,  $P(A) + P(B) + P(C) = 1$ .

**Example. 1** In an urn, there are 2 white, 3 black and 5 yellow balls. One ball is drawn at random. Find the probability that it is (i) white (ii) black (iii) yellow.

$$\begin{bmatrix} W & -2 \\ B & -3 \\ Y & -5 \end{bmatrix}$$

**Solution :** There are altogether  $(2+3+5)$  i.e. 10 balls and choosing any one of them can be done in 10 ways. These 10 ways are all equally likely, mutually exclusive in the sense that when one is chosen, the remaining will not be chosen and further, the 10 ways form an exhaustive system because all the possible ways have been included.

(i) If  $W$  represents the event of choosing 1 white ball out of the 2, there are

2 ways which are favourable to  $W$ . Then by definition,  $P(W) = \frac{2}{10}$ .

Similarly, if  $B$  and  $Y$  respectively denote the events of drawing a black

and a yellow ball then,  $P(B) = \frac{3}{10}$

&  $P(Y) = \frac{5}{10}$ , for cases (ii) & (iii)

Now,  $P(W)+P(B)+P(Y) = \frac{2}{10} + \frac{3}{10} + \frac{5}{10} = \frac{10}{10} = 1$ .

**Note 1.** If  $A$  and  $B$  are two events then, we can use the symbol  $P(A \cup B)$  to denote the happening of one of them. If they form an exhaustive system, then, surely,  $P(A \cup B) = 1$ . Further, if  $A$  and  $B$  are mutually exclusive, then  $P(A \cup B) = P(A) + P(B)$ .

Thus,  $P(A) + P(B) = 1$ , when  $A$  and  $B$  are both mutually exclusive and exhaustive events.

**Note 2.** If  $A$  and  $B$  are not mutually exclusive then,  $P(A \cup B) \neq P(A) + P(B)$

**Note 3.**  $P(A \cup B)$  exists even if  $A$  and  $B$  are not mutually exclusive and also when they do not form an exhaustive system of events. Even then,  $P(A \cup B)$  denotes the probability of happening of one of them.

However, we shall not discuss the case here.

**Example 2.** A fair die is tossed twice. Write the sample space. How many sample points are there? Find the probability that the sum of points obtained in the two tosses is (i) equal to 10. (ii) greater than 10. (iii) less than 10.

**Solution :** The sample space  $S$  is given by  
 $S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6),$   
 $(2,1), (2,2), (2,3), (2,4), (2,5), (2,6),$

(3,1),(3,2),(3,3),(3,4),(3,5),(3,6),  
 (4,1),(4,2),(4,3),(4,4),(4,5),(4,6),  
 (5,1),(5,2),(5,3),(5,4),(5,5),(5,6),  
 (6,1),(6,2),(6,3),(6,4),(6,5),(6,6)}

There are, in all  $6 \times 6$  i.e. 36 samples or outcomes.

Let A, B and C denote the events of getting the sum of points equal to greater than and less than 10 respectively.

The outcomes favourable to A are (4,6),(5,5) and (6,4) which are 3 in number.

$$\therefore P(A) = \frac{\text{Number of outcomes favourable to A}}{\text{Total number of possible outcomes}} = \frac{3}{36} = \frac{1}{12}$$

The outcomes favourable to B are (5,6),(6,5), and (6,6).

$$\therefore P(B) = \frac{3}{36} = \frac{1}{12}$$

Since A, B, C are mutually exclusive events forming an exhaustive system, therefore  $P(A) + P(B) + P(C) = 1$

$$\text{i.e. } \frac{1}{12} + \frac{1}{12} + P(C) = 1$$

$$\therefore P(C) = 1 - \frac{1}{6} = \frac{5}{6}$$

### Independent events

As stated earlier, two events A and B are said to be independent if the occurrence of A has no effect on the occurrence of B and vice-versa.

We use the symbol AB to denote the event of the combined occurrence of A and B.

Further, if A and B are independent, then we have  $P(AB) = P(A)P(B)$

**Example 3.** If a coin is tossed twice, what is the probability of getting head in both the cases?

**Solution:** We can look upon the problem from two stand points

**One :** If S is the sample space of the tossing of a coin twice, then,  $S = \{HH, HT, TH, TT\}$

Thus, out of the four exhaustive no. of outcomes only one is favourable to {HH}

$$\therefore P(HH) = \frac{1}{4}$$

**Two :** Probability of getting head in the 1st toss =  $\frac{1}{2}$  i.e.  $P(H) = \frac{1}{2}$

and Probability of getting head in the second toss =  $\frac{1}{2}$  i.e.  $P(H) = \frac{1}{2}$

Since, the outcome of the first toss has no effect on the outcome of the second toss, we get

$$P(HH) = P(H)P(H) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$





### 14.13 A note on playing cards

Playing cards mean those cards popularly used in many indoor games like Bridge, Rummy etc.

A deck or pack of cards consists of 52 cards divided into two sets of 26 cards each. One set is red set and the other set is black set. In each colour set, there are two suits of 13 cards each. In a suit, there are 3 face cards called King, Queen and Jack. In each suit, there is one card called Ace and nine numeral cards starting from 2 and ending at 10. When one says four powers in a suit, it means Ace, King, Queen and Jack of the suit. Thus, there are two colours, four suits, 12 face cards consisting of four Kings, four Queens and four Jacks. Before any draw, cards are well shuffled to ensure that each cards has equal chance of being drawn.

[ A demonstration of playing cards and dice to the students is desirable so that they become familiar with the names and terms associated with these pieces of games of chance]

The four suits have their names, colours, symbols and number of cards as follows:

Suit name	Colour	Symbol	No. of Cards			
			Ace	Face	Numeral	Total
Heart	Red		1	3	9	13
Diamond	Red		do	do	do	do
Club	Black		do	do	do	do
Spade	Black		do	do	do	do

**Example 4.** A die is tossed 4 times. Find the probability that 6 appears at least once.

**Solution :** The probability that 6 appears in a single toss =  $\frac{1}{6}$

$\therefore$  The probability that 6 does not appear in a toss =  $1 - \frac{1}{6} = \frac{5}{6}$

Now, the appearances in all the 4 tosses are all independent. Thus, the probability

that 6 does not appear in all the 4 tosses =  $\frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} = \left(\frac{5}{6}\right)^4$

Let us consider two events namely A, the event in which all the four tosses do not give any 6 and B, the event in which there is at least one 6 in the 4 tosses.

Now, these two events are mutually exclusive and they form an exhaustive system of events. We are to find P(B).

But,  $P(A) + P(B) = 1$

$$\Rightarrow P(B) = 1 - P(A)$$

$$= 1 - \left(\frac{5}{6}\right)^4$$

**Example 5.** From a well suffled pack of cards, one card is drawn at random. Find the probability that the drawn card is a power.

**Solution:** There are 52 cards in the pack and so one card can be drawn in 52 ways.

Further, there are 4 suits in the pack and each suit has four powers (viz. Ace, King, Queen and Jack). So, there are 16 powers in the pack and out of these, one may be drawn in 16 ways.

Thus, out of the 52 ways, 16 are favourable to the event.

Thus, the reqd. probability =  $\frac{16}{52} = \frac{4}{13}$ .

**Example 6.** Two dice are thrown. Find the probability that the sum of their points is 11.

**Solution :** The favourable cases of giving a sum of 11 points are shown below:

Point on the 1st die	Point on the second die
5	6
6	5

That is to say, there are only 2 favourable cases of getting the sum of 11 points, because no other pair of points will give 11 as sum of points. But there are  $6 \times 6$  ways in which the two dice can turn up giving different sums of points ranging from  $(1+1)$  i.e. 2 to  $(6+6)$  i.e. 12.

$$\text{Thus, the reqd. probability} = \frac{2}{6 \times 6} = \frac{1}{18}$$

**Example 7.** From a pack of cards, two cards are drawn at random (one after another without replacing). Find the probability that both are aces.

**Solution :** One card out of 52, can be drawn in 52 ways. And one ace out of 4, can be drawn in 4 ways.

$$\therefore \text{The probability that the first card drawn is an ace} = \frac{4}{52} = \frac{1}{13}$$

Having drawn one ace in the first draw, we are left with 51 cards of which 3 are aces.

$$\text{So, the probability that the second card drawn is also an ace} = \frac{3}{51} = \frac{1}{17}$$

The required probability that both the cards drawn are aces

$$= \frac{1}{13} \times \frac{1}{17} \quad [\because P(AB) = P(A) P(B) \text{ when A and B are independent}]$$

$$= \frac{1}{221}$$

#### 14.14 Reconciliation between the empirical and the classical definition of probability.

In the empirical definition, we take the probability of an event A as the ratio

$$\frac{\text{The no. of trials in which A appeared}}{\text{Total no. of trials}}$$

$$= \frac{m}{n} \text{ (say)}$$

Supposing in tossing a coin 1000 times if H appeared 507 times then,

$$P(H) = \frac{507}{1000} = 0.507. \text{ On the other hand, by the classical definition, } P(H) = \frac{1}{2} = 0.5.$$

In the tossing of a fair coin a large number of times, it has been observed that the ratio  $\frac{m}{n}$  approaches to 0.5. The larger the number of trials i.e.  $n$ , the nearer is the ratio  $\frac{m}{n}$  to 0.5.

Thus, like the case in any deterministic experiment, giving allowance to experimental errors, we take the theoretical or classical value as the probability of the event to which empirical values converge for large values of  $n$ .

### EXERCISE 14.1

1. Give some events in our day to day life that are mutually exclusive.
2. If two events A and B are such that  $P(A) + P(B) = 1$ , then, write  $P(B)$  in terms of  $P(\bar{A})$ .
3. A, B, C are three events which are equally likely, mutually exclusive but not forming an exhaustive system of events. Then, show that  $P(A) = P(B) = P(C) = \frac{1}{3}$ . Give one example of such a situation.
4. Distinguish between subjective and objective probabilities giving examples in each case.
5. When do you say that a die is fair ?
6. When a fair coin is tossed 3 times, how many outcomes can be there? Are the outcomes equally likely?
7. From an urn containing 4 white and 5 red balls, two balls are drawn at random. Find the probability that at least one is white.
8. From a pack of cards, two cards are drawn at random after a thorough shuffle. Find the probability that both are kings.
9. Two fair dice are rolled. Find the probability that the sum of the points is 7. Also, find the sum of points which is the most probable by showing all the possible sum of points in a chart or table.
10. Given that  $p$  is the probability that a person aged  $x$  years will die in a year, find the probability that none of the four persons all aged  $x$  years will die in a year.  
[This probability  $p$  can be obtained from a table called life table in the study of vital statistics]

11. In the above exercise if Mr. A is one of the four persons, find the probability that at least one of them will die in a year and Mr. A is the first person to die.
12. From a well shuffled pack of cards, two cards are drawn at random. Find the probability that both the cards are diamonds.
13. A letter is chosen at random from the word 'ASSASSINATION'. Find the probability that the letter is (i) a vowel (ii) a consonant.
14. In a 20-20 cricket match, a batsman hits a boundary 5 times out of 24 balls he faced. Find the probability that he did not hit a boundary in a ball he faced.
15. A die is thrown. Find the probability of the following events:
  - (a) A prime number will appear.
  - (b) A number less than 6 will appear.
  - (c) A number more than 6 will appear.
  - (d) A number less than or equal to 3 will appear.
16. There are four men and three ladies in a council. If two council members are selected at random for a committee, how likely is it that both are ladies?

### ANSWER

1. The result of a lottery, the result of game, the gender of a person, the result of a student's examination, the result of the competition for a gold medal among a number of contenders etc.
2.  $P(\overline{B}) = P(\overline{A})$
5. When the six faces have equal probabilities of turning up in every toss.
6.  $2^3$  i.e. 8. Yes
7.  $\frac{13}{18}$
8.  $\frac{1}{221}$
9.  $\frac{1}{6}, 7$
10.  $(1-p)^4$
11.  $\frac{1-(1-p)^4}{4}$
12.  $\frac{1}{17}$
13. (i)  $\frac{6}{13}$  (ii)  $\frac{7}{13}$
14.  $\frac{19}{24}$
15. (a)  $\frac{1}{2}$  (b)  $\frac{5}{6}$  (c) 0 (d)  $\frac{1}{2}$
16.  $\frac{1}{7}$



## SUMMARY

**In this chapter we have studied the following items:**

1. Classical definition of probability
2.  $P(A) + P(\bar{A}) = 1$   
Here,  $\bar{A}$  is the event of non-occurrence of the event A.
3. For mutually exclusive events A and B,  
 $P(A \cup B) = P(A) + P(B)$ .
4. If A and B are mutually exclusive and exhaustive then  
$$P(A \cup B) = P(A) + P(B)$$
and  $P(A) + P(B) = 1$
5. For independent events A and B  
 $P(AB) = P(A)P(B)$

\*\*\*\*\*