

னிரிப்புகு மூ நான் கான் குடி (யுல) DEPARTMENT OF EDUCATION (S) Government of Manipur

## CLASS – X MATHEMATICS CHAPTER – 2 POLYNOMIALS

# NOTES

#### Degree of a polynomial

If p(x) is a polynomial in x, the highest exponent of x in p(x) is called degree of p(x).

A polynomial is called linear, quadratic, cubic or biquadratic according as its degree is one, two, three or four respectively.

#### > Working rule to divide a polynomial by another polynomial

- 1. Write the dividend and divisor after arranging the term in the descending order of their degrees.
- 2. Divide the highest degree term (first term) of the dividend by the highest degree term (first term) of the divisor to get the first term of the quotient.
- **3.** Multiply the divisor by the first term of the quotient and subtract this product from the dividend to get the remainder.
- **4.** Taking the remainder as the new dividend, keeping the divisor same, find the quotient and remainder to get the next quotient term.
- **5.** Continue the process till the degree of the remainder is less than the degree of the divisor.

### > Division Algorithm for Polynomials

If p(x) and d(x) are any two polynomials with  $d(x) \neq 0$ , then we can find polynomials q(x) and r(x) such that  $p(x) = d(x) \times q(x) + r(x)$ , where either r(x) = 0 or degree of r(x) < degree of d(x).

In case degree of the dividend p(x) is less than that of the divisor d(x), then we take q(x) = 0 and r(x) = p(x).

Remainder Theorem

Let p(x) be any polynomial of degree greater than or equal to one and let a be any real number. If p(x) is divided by the linear polynomial x - a, then the remainder is p(a).

Factor Theorem: If p(x) is a polynomial of degree  $\ge 1$  and a is any real number, then x - a is a factor of p(x) if and only if p(a) = 0.